

MH-004 PHQ-9 English

PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

Patient's Name (<i>Print</i>):			D	ate:		
Over the last 2 weeks, how often have you been bothered by a (Please answer all numbered questions. <i>Use an "X" to indicate</i>			_	ies?		
			Not at all (0)	Several Days (1)	More than half the days (2)	Nearly every day (3)
Little interest or pleasure in doing things						
2. Feeling down, depressed, or hopeless						
3. Troubled falling or staying asleep						
4. Feeling tired or having little energy						
5. Poor appetite or overeating						
6. Feeling bad about yourself – or that you are a failure of let yourself or your family down	or hav	е				
7. Trouble concentrating on things, such as reading the newspaper or watching television						
8. Moving or speaking so slowly that other people could noticed. Or the opposite – being so fidgety or restless that have been moving around a lot more than usual						
9. Thoughts that you would be better off dead, or of hur yourself	ting					
	Office	Add	d columns			ł
	Use		Total	x 1 =	x 2 =	x 3 =
	only		Total			
•			TOTAL			
				Not difficult a	at all	
10. If you made and "X" for any of the above issues, how difficult for you to do your work, take care of things at home.				Somewhat difficult		
made it for you to do your work, take care of things at home, or get along with other people?			VVICII	Very difficult		
•				Extremely dif	ticult	

written: 10/13/22

rev. 10/13/22

Page 1 of 1

PHQ-9* Questionnaire for Depression Scoring and Interpretation Guide

For physician use only

Scoring:

Count the number (#) of boxes checked in a column. Multiply that number by the value indicated below, then add the subtotal to produce a total score. The possible range is 0-27. Use the table below to interpret the PHQ-9 score.

Not at all	(#)	x 0 =	
Several days	(#)	x 1 =	
More than half the days	(#)	x 2 =	
Nearly every day	(#)	x 3 =	
Total score:			

Interpreting PHQ-9 Scores					
Diagnosis	Total Score	For Score	Action		
Minimal depression	0-4	≤ 4	The score suggests the patient may not need depression treatment		
Mild depression Moderate depression	5-9 10-14	5 - 14	Physician uses clinical judgment about treatment, based on patient's duration of symptoms and functional impairment		
Moderately severe depression Severe depression	15-19 20-27	> 14	Warrants treatment for depression, using antidepressant, psychotherapy and/or a combination of treatment.		

^{*} The PHQ-9 is described in more detail at the Pfizer website: http://www.phqscreeners.com/