

Generalized Anxiety Disorder 7-item (GAD-7) scale

Over the last 2 weeks, how often have you been bothered by the following problems?	Not at all sure	Several days	Over half the days	Nearly every day
1. Feeling nervous, anxious, or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it's hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	1	2	3
<i>Add the score for each column</i>	+	+	+	
Total Score (<i>add your column scores</i>) =				

If you checked off any problems, how difficult have these made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all _____

Somewhat difficult _____

Very difficult _____

Extremely difficult _____

Scoring

Scores of 5, 10, and 15 are taken as the cut-off points for mild, moderate and severe anxiety, respectively. When used as a screening tool, further evaluation is recommended when the score is 10 or greater.

Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. It is moderately good at screening three other common anxiety disorders - panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%) and post-traumatic stress disorder (sensitivity 66%, specificity 81%).

Source: Spitzer RL, Kroenke K, Williams JBW, Lowe B. A brief measure for assessing generalized anxiety disorder. *Arch Intern Med.* 2006;166:1092-1097.

PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

NAME: _____

DATE: _____

Over the last 2 weeks, how often have you been bothered by any of the following problems?
(use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead, or of hurting yourself	0	1	2	3

add columns + +

(Healthcare professional: For interpretation of TOTAL, TOTAL: please refer to accompanying scoring card).

<p>10. If you checked off <i>any problems</i>, how <i>difficult</i> have these problems made it for you to do your work, take care of things at home, or get along with other people?</p>	<p>Not difficult at all _____</p> <p>Somewhat difficult _____</p> <p>Very difficult _____</p> <p>Extremely difficult _____</p>
--	--

PHQ-9 Patient Depression Questionnaire

For initial diagnosis:

1. Patient completes PHQ-9 Quick Depression Assessment.
2. If there are at least 4 ✓s in the shaded section (including Questions #1 and #2), consider a depressive disorder. Add score to determine severity.

Consider Major Depressive Disorder

- if there are at least 5 ✓s in the shaded section (one of which corresponds to Question #1 or #2)

Consider Other Depressive Disorder

- if there are 2-4 ✓s in the shaded section (one of which corresponds to Question #1 or #2)

Note: Since the questionnaire relies on patient self-report, all responses should be verified by the clinician, and a definitive diagnosis is made on clinical grounds taking into account how well the patient understood the questionnaire, as well as other relevant information from the patient.

Diagnoses of Major Depressive Disorder or Other Depressive Disorder also require impairment of social, occupational, or other important areas of functioning (Question #10) and ruling out normal bereavement, a history of a Manic Episode (Bipolar Disorder), and a physical disorder, medication, or other drug as the biological cause of the depressive symptoms.

To monitor severity over time for newly diagnosed patients or patients in current treatment for depression:

1. Patients may complete questionnaires at baseline and at regular intervals (eg, every 2 weeks) at home and bring them in at their next appointment for scoring or they may complete the questionnaire during each scheduled appointment.
2. Add up ✓s by column. For every ✓: Several days = 1 More than half the days = 2 Nearly every day = 3
3. Add together column scores to get a TOTAL score.
4. Refer to the accompanying **PHQ-9 Scoring Box** to interpret the TOTAL score.
5. Results may be included in patient files to assist you in setting up a treatment goal, determining degree of response, as well as guiding treatment intervention.

Scoring: add up all checked boxes on PHQ-9

For every ✓ Not at all = 0; Several days = 1;
More than half the days = 2; Nearly every day = 3

Interpretation of Total Score

Total Score	Depression Severity
1-4	Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression

PHQ9 Copyright © Pfizer Inc. All rights reserved. Reproduced with permission. PRIME-MD ® is a trademark of Pfizer Inc.

Ht:

Date:

Wt:

BMI:

10-12 YEAR VISIT

Development

1. Sex Education
 - It's time to talk birds and bees, but answer questions about sex factually, using actual language and terms.
 - Talk with your children about the changes his/her body will undergo during puberty. Prepare your daughter for the onset of menarche and her periods.
2. Actively discourage harmful habits like smoking, alcohol, and drugs. Talk with your child about peer pressure and how to say no.
3. Talk to your children about bullying. Be involved at their school and keep open communication with teachers to ensure your child is safe at school.
4. Organized sports are a great way to boost confidence and establish healthy habits. Be careful of year round sports, which increases injury risk. It's best to move from one sport to another rather than simply concentrating on one sport year round.

Injury Prevention

1. Teach your child to **ALWAYS** wear seat belts even if others in the car do not. By this age, buckling up should be an almost automatic reflex. No riding in the front seat until age 13! CDC rules, so blame them!
2. Teach your child about firearm safety. Your child should **NEVER** handle firearms, even pellet and BB guns, unless you are present. Teach your child that if another child asks him to "look at" or "play with" guns, he is to come home at once.
3. Continued used of protective equipment for high risk activities (HELMETS are cool!)

Good Health Habits and Self Care

1. Continue to maintain a regular bedtime routine. Children these ages need 10-12 hours a night.
2. Set limits on TV viewing/video games/social media. Establish these expectations now! Out children have access to limitless amounts of electronic information, and much of it is terrifying.
3. Your child should practice good dental care and self-hygiene.
4. Obesity is a significant health threat. If you are worried, PLEASE bring it up at your visit.
5. To stay healthy, here are some easy rules to remember: 5-2-1-0
 - FIVE servings of fruits/veggies a day
 - Less than TWO hours of total screen time a day
 - ONE hour of outdoor activity a day
 - ZERO calories from sugary drinks

Next Visit

Continue yearly visits, which also serve as sports and camp physicals. Some parents come around the birthday. Some come every fall break or other convenient time. It is a good idea to avoid late June until early August for your yearly check. These are especially busy times here at the office.

Your 11 year old will receive at least 2 vaccines as well as a finger poke and urine check. Let them know that all 11 year olds come in very nervous about this visit, but almost all of them leave realizing that they worried for no good reason.

All rising Teens need to get the HPV (Gardasil) vaccine. More people will die this year from HPV disease than died yearly from measles before that vaccine came out. It is a "no-brainer" vaccine. If started before age 15, your child will only need two shots to complete the series. After age 15, it is a three shot series.

Don't forget to get a flu vaccine.

CONCUSSION Information Sheet



This sheet has information to help protect your children or teens from concussion or other serious brain injury. Use this information at your children's or teens' games and practices to learn how to spot a concussion and what to do if a concussion occurs.

What Is a Concussion?

A concussion is a type of traumatic brain injury—or TBI—caused by a bump, blow, or jolt to the head or by a hit to the body that causes the head and brain to move quickly back and forth. This fast movement can cause the brain to bounce around or twist in the skull, creating chemical changes in the brain and sometimes stretching and damaging the brain cells.

How Can I Help Keep My Children or Teens Safe?

Sports are a great way for children and teens to stay healthy and can help them do well in school. To help lower your children's or teens' chances of getting a concussion or other serious brain injury, you should:

- Help create a culture of safety for the team.
 - › Work with their coach to teach ways to lower the chances of getting a concussion.
 - › Talk with your children or teens about concussion and ask if they have concerns about reporting a concussion. Talk with them about their concerns; emphasize the importance of reporting concussions and taking time to recover from one.
 - › Ensure that they follow their coach's rules for safety and the rules of the sport.
 - › Tell your children or teens that you expect them to practice good sportsmanship at all times.
- When appropriate for the sport or activity, teach your children or teens that they must wear a helmet to lower the chances of the most serious types of brain or head injury. However, there is no "concussion-proof" helmet. So, even with a helmet, it is important for children and teens to avoid hits to the head.



Plan ahead. What do you want your child or teen to know about concussion?

How Can I Spot a Possible Concussion?

Children and teens who show or report one or more of the signs and symptoms listed below—or simply say they just "don't feel right" after a bump, blow, or jolt to the head or body—may have a concussion or other serious brain injury.

Signs Observed by Parents or Coaches

- Appears dazed or stunned.
- Forgets an instruction, is confused about an assignment or position, or is unsure of the game, score, or opponent.
- Moves clumsily.
- Answers questions slowly.
- Loses consciousness (*even briefly*).
- Shows mood, behavior, or personality changes.
- Can't recall events *prior to* or *after* a hit or fall.

Symptoms Reported by Children and Teens

- Headache or "pressure" in head.
- Nausea or vomiting.
- Balance problems or dizziness, or double or blurry vision.
- Bothered by light or noise.
- Feeling sluggish, hazy, foggy, or groggy.
- Confusion, or concentration or memory problems.
- Just not "feeling right," or "feeling down."

Talk with your children and teens about concussion. Tell them to report their concussion symptoms to you and their coach right away. Some children and teens think concussions aren't serious or worry that if they report a concussion they will lose their position on the team or look weak. Be sure to remind them that *it's better to miss one game than the whole season.*

To learn more, go to www.cdc.gov/HEADSUP



Centers for Disease Control and Prevention
National Center for Injury Prevention and Control

Concussions affect each child and teen differently. While most children and teens with a concussion feel better within a couple of weeks, some will have symptoms for months or longer. Talk with your children's or teens' health care provider if their concussion symptoms do not go away or if they get worse after they return to their regular activities.



What Are Some More Serious Danger Signs to Look Out For?

In rare cases, a dangerous collection of blood (hematoma) may form on the brain after a bump, blow, or jolt to the head or body and can squeeze the brain against the skull. Call 9-1-1 or take your child or teen to the emergency department right away if, after a bump, blow, or jolt to the head or body, he or she has one or more of these danger signs:

- One pupil larger than the other.
- Drowsiness or inability to wake up.
- A headache that gets worse and does not go away.
- Slurred speech, weakness, numbness, or decreased coordination.
- Repeated vomiting or nausea, convulsions or seizures (shaking or twitching).
- Unusual behavior, increased confusion, restlessness, or agitation.
- Loss of consciousness (passed out/knocked out). Even a brief loss of consciousness should be taken seriously.

Children and teens who continue to play while having concussion symptoms or who return to play too soon—while the brain is still healing—have a greater chance of getting another concussion. A repeat concussion that occurs while the brain is still healing from the first injury can be very serious and can affect a child or teen for a lifetime. It can even be fatal.

Revised 5/2015


What Should I Do If My Child or Teen Has a Possible Concussion?

As a parent, if you think your child or teen may have a concussion, you should:

1. Remove your child or teen from play.
2. Keep your child or teen out of play the day of the injury. Your child or teen should be seen by a health care provider and only return to play with permission from a health care provider who is experienced in evaluating for concussion.
3. Ask your child's or teen's health care provider for written instructions on helping your child or teen return to school. You can give the instructions to your child's or teen's school nurse and teacher(s) and return-to-play instructions to the coach and/or athletic trainer.

Do not try to judge the severity of the injury yourself. Only a health care provider should assess a child or teen for a possible concussion. Concussion signs and symptoms often show up soon after the injury. But you may not know how serious the concussion is at first, and some symptoms may not show up for hours or days.

The brain needs time to heal after a concussion. A child's or teen's return to school and sports should be a gradual process that is carefully managed and monitored by a health care provider.



To learn more, go to www.cdc.gov/HEADSUP

You can also download the CDC *HEADS UP* app to get concussion information at your fingertips. Just scan the QR code pictured at left with your smartphone.

Discuss the risks of concussion and other serious brain injury with your child or teen and have each person sign below.

Detach the section below and keep this information sheet to use at your children's or teens' games and practices to help protect them from concussion or other serious brain injury.

I learned about concussion and talked with my parent or coach about what to do if I have a concussion or other serious brain injury.

Athlete Name Printed: _____ Date: _____

Athlete Signature: _____

I have read this fact sheet for parents on concussion with my child or teen and talked about what to do if they have a concussion or other serious brain injury.

Parent or Legal Guardian Name Printed: _____ Date: _____

Parent or Legal Guardian Signature: _____