

Primary Immunodeficiency Risk Panel

Patient		Specimen		Ordering Physician	
First Name:	Test	Specimen Type:	Buccal Swab	Physician:	Dr. Test
Last name:	Patient	Collection Date:	07-20-2022	Institution:	Medical Clinic
Date of Birth:	01-01-9977	Received Date:	07-21-2022	Reported Date:	08-05-2022
Gender:	Male	Panel Coverage :	>=50%	Ref Accession:	N/A
Accession ID:	E2207218829	Average Read Depth:	>50%		

SUMMARY OF RESULTS

NEGATIVE

Summary of Result : NEGATIVE

Recommendations

A negative result indicates that the individual does not have pathogenic or likely pathogenic variants known to be associated with Immunodeficiency risk from the list of genes evaluated through next generation sequencing (NGS). For more information, please contact the National Society of Genetic Counselors and locate a practitioner near you at https://www.nsgc.org/page/find-a-genetic-counselor or by phone at 312.321.6834.



Elite Clinical Laboratory 3600 S Gessner Road,Suite 110, Houston,Texas-77063 USA, Phone:281.378.2116, Fax:281.466.2483, CLIA#45D1061571, Lab Director:Dr.Albert Chen MD.

Patient Details

Name : Patient, Test

Date of Birth : 01-01-9977

Accession Number : E2207218829

Test Methodology

Immunodeficiency Risk Factor Screening 46 gene panel screening performed at Elite Clinical Laboratory LLC utilizes Next Generation Sequencing (NGS) technology using targeted gene enrichment chemistry on the Illumina® MiniSeq platform. Genomic DNA extracted from Buccal swabs (Dry and Wet) are tagemented and enriched for regions of interest using probes specific for each region specified in the BED (Browser Extensible Data) file. Positive (www.coriell.org) and Negative controls are included with each machine run to ensure the accuracy of amplicon preparation alongside targeted sequencing on coding regions and intronic/exonic boundaries of interest and precise finding of the variant on the positive control and negative control respectively. Exclusions from analysis are listed below Sequences obtained are then aligned against a human reference genome and variants such as SNVs (Small Nucleotide Variants) and Indels (Insertions and Deletions) are noted.For a detailed list of regions covered and comprehensive statistics by Elite Clinical Laboratory LLC Immunodeficiency screen, please contact us on Email:info@elitelabs.com

Computational analysis and variant calling is performed by Elite (www.elitelabs.com). Briefly, reads from the sequence output were aligned to the human reference genome (GRCh37) using the Burrows-Wheeler Aligner (BWA). Variants to the reference were called using the Genomic Analysis Tool Kit (GATK). The variants were annotated and filtered using the Elite Clinical Laboratory LLC analysis work flow implementing the ACMG guidelines for interpretation of sequence variants. This includes comparison against the gnomAD population catalog of variants in 123,136 exomes, the 1000 Genomes Project Consortium's publication of 2,500 genomes, the NCBI-ClinVar database of clinical assertions on variant's pathogenicity and multiple lines of computational evidence on conservation and functional impact. The following databases and insilico algorithms are used to annotate and evaluate the impact of the variant in the context of human disease: 1000 genomes, gnomAD, ClinVar, OMIM, dbSNP, NCIB Ref Seg Genes, ExAC Gene Constraints, VS-SIFT, VS PolyPhen2, PhyloP, GERP+, GeneSplicer, Max EntScan, NNSplice, PWM Splice Predictor. Analysis was using the HGVS nomenclature (www.hgvs.org/mutnomen)as implemented by custom reported transcript annotation algorithm. Only variants with an end ACMG classification of "Pathogenic" or "Likely Pathogenic" are reported and VUS/Benign/Likely Benign variants are not reported. Classification of pathogenicity is consistently updated by the National Institutes of Health and the information found within this report is consistent with the current knowledge base as the knowledge base changes, so may clinical interpretation of variants.

All reports are reviewed prior to release by either Elite Clinical Laboratory LLC's Technical Supervisor or the General Supervisor.

Genes Evaluated

ATM, BLM, BRCA2, BTK, CDX1, CFTR, CYBA, CYBB, F13B, F5, F7, F9, FANCC, FGB, G6PC, G6PD, IFNGR1, IFNGR2, ITGB2, JAGN1, JAK2, MEFV, MPL, MSH6, MST1, MYD88, NCF1, NFKB2, NRAS, PALB2, PIK3CD, PLCG2, PMS2, PTEN, PTPRC, RAG1, RAG2, RFXANK, RUNX1, SPINK5, STAT1, STAT3, STK4, TERT, TNFRSF13B, VPS13B.



Elite Clinical Laboratory 3600 S Gessner Road, Suite 110, Houston, Texas-77063 USA, Phone:281.378.2116, Fax:281.466.2483, CLIA#45D1061571, Lab Director:Dr.Albert Chen MD.

Patient Details

Date of Birth : 01-01-9977

Accession Number : E2207218829

Test Limitations

Name : Patient, Test

Some Genes/variations for the following genomic regions may not be reported such as: large genomic rearrangements greater than 50 bp in length, rare (low frequency) mutations, or structural (non-coding) variations and pseudogenes. MST1 gene and its variants detected on chromosome-3 are only considered for tertiary analysis and reporting. Pseudogene for the same gene on chromosome-1 is excluded from further analysis and reporting.

Excluded: MST1L macrophage stimulating 1 like (pseudogene) Location: 1p36.13; chr 1 : NC 000001.10:17,080,142 - 17,091,958

Rare diagnostics errors may occur if these mutations occur within the priming sequencing regions Presence of a pathogenic/likely pathogenic variant does not guarantee that an individual will develop a Immunodeficiency, nor is the absence of such variants a guarantee that an individual will not develop a 'Immunodeficiency'. The results of this genetics screening is meant strictly to guide a physician in the management of their patient's health. Any Likely pathogenic or Pathogenic variants detected in the report should be clinically correlated (As per Physicians Advice) and confirmed via the orthogonal testing platforms (Sanger Sequencing or PCR based tests).

Regulatory Disclosures

Genetic-based hereditary Immunodeficiency Risk Factor screening is intended as a tool to guide physicians in the management of their patients and should NOT be treated as a diagnostic tool NGS-based hereditary Immunodeficiency screening is considered a high-complexity laboratory-developed test (LDT) by CMS under the Clinical Laboratory ImprovementAmendment (CLIA) and is not FDA-cleared. The test and performance metrics were validated in house by Elite Clinical LaboratoryLLC technical personnel (or designated scientific advisors) and approved by their Laboratory Director The results are intended for use only by the ordering physician and/or designated healthcare provider. The ordering provider is responsible for 1) ascertaining the medical necessity of the ordered test, 2) resulting diagnoses, 3) management of the disease and/or decisions based on the data provided. Results rely on collection personnel following specified collection and shipment protocols.

REFERENCES

Richards, Sue, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genetics in medicine 17.5 (2015): 405.Exome Aggregation Consortium et al. Analysis of Protein-Coding Genetic Variation in 60,706 Humans. Nature 536.7616 (2016): 285–291. PMC. Web. 13 May 2018. The 1000 Genomes Project Consortium. A Global Reference for Human Genetic Variation. Nature 526.7571 (2015): 68–74. PMC. Web. 13 May 2018.

Testing was performed by Elite Clinical Laboratory LLC , 3600 S Gessner Rd, Houston, TX, 77 063, USA CLIA # 45D2229553

Laboratory Director: Dr. Albert Chen PhD.