

Dysport.  
(abobotulinumtoxinA)

READY TO TAKE ON THE LINES  
BETWEEN YOUR BROWS?

LIVE YOUR  
ACCOMPLISHMENTS  
*Not Your Lines*



Soften and smooth frown lines between the brows in a natural-looking way.

Actual patient. Individual results may vary.

**Indication**

*Dysport*<sup>®</sup> (abobotulinumtoxinA) is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

**Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.**

# SAY "see ya" TO FROWN LINES BETWEEN YOUR BROWS

*Dysport*<sup>®</sup> (*abobotulinumtoxinA*) for Injection may soften and smooth frown lines between the brows for natural-looking results that may last up to 5 months.<sup>1-3\*</sup>

98%  
OF USERS

## Natural-Looking

98% of users<sup>†</sup> say their results look natural, and 97% would get *Dysport* again.<sup>4††</sup>

2-3  
DAYS

## Fast-Acting

Over half of users<sup>†</sup> see improvement in **2-3 days** after one 10- to 20-minute treatment.<sup>1,2,5§</sup>

5  
MONTHS

## Long-Lasting

Smoothed lines between the eyebrows may last for **up to 5 months**.<sup>1-3\*</sup>

"DYSPORE GIVES ME  
*natural-looking results*

THAT AREN'T OVERDONE."

-Rebecca, 35, Actual *Dysport* Patient,  
Mom, Volunteer & Full-time Project Manager

\*GL-1 and GL-3 evaluated subjects for at least 150 days following treatment. Based on a  $\geq 1$ -grade improvement from baseline utilizing data from 2 double-blinded, randomized, placebo-controlled pivotal studies (GL-1, GL-3) in a post-hoc analysis.

†User = clinical trial subject.

††Subject's satisfaction and self-perception reported 3 weeks after treatment (n=531).

§A secondary endpoint based on Kaplan-Meier estimates of cumulative rate of time to onset of response. The median time to onset of response was 3 days in GL-1 (*Dysport*: 55/105 [52%]; placebo: 3/53 [6%]) and GL-2 (*Dysport*: 36/71 [51%]; placebo: 9/71 [13%]), and 2 days in GL-3 (*Dysport*: 110/200 [55%]; placebo: 4/100 [4%]).

Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 6 and 7.



"LIVING MY BEST LIFE MEANS  
*living my look*  
...INSIDE AND OUT."

-Alia, 37, Actual *Dysport* Patient,  
Entrepreneur and Small Business Owner

Actual patient. Individual results may vary.

## HERE'S HOW IT WORKS

*Dysport* helps soften and smooth frown lines between the eyebrows without changing the look or movement of the rest of your face.

*Dysport* blocks the signal from the nerve to the muscles, which temporarily prevents contraction of the muscles that cause frown lines between the eyebrows. Untreated muscles still work normally, so you can continue to express yourself naturally and confidently.

# ASPIRE

GALDERMA REWARDS

## REWARDS BEYOND THE RESULTS

Help your investment go further. Start earning points for each qualifying *Dysport*<sup>®</sup> (*abobotulinumtoxinA*) for Injection treatment and redeem certificates that can be used across the Galderma portfolio, including on *Restylane*<sup>®</sup> and *Sculptra*<sup>®</sup>.



Sign up and receive an instant reward. Learn more and join at [AspireRewards.com](https://www.aspire-rewards.com).


*"I had two frown lines*

BETWEEN MY EYEBROWS,  
ONE FOR EACH KID."

-Rebecca, 35, Actual *Dysport*<sup>®</sup> Patient,  
Mom, Volunteer & Full-time Project Manager


*DYSPORT* MAY HELP  
SOFTEN LINES BETWEEN  
THE BROW FOR UP  
TO **5 MONTHS**.<sup>1-3\*</sup>

BEFORE



Rebecca, 35, actual *Dysport* patient. Frown before *Dysport* treatment.

AFTER



Frown after 50 units, day 27. Individual results may vary.

AFTER JUST ONE 10- TO  
20-MINUTE TREATMENT,  
MANY SEE RESULTS IN  
AS LITTLE AS **2-3 DAYS**.<sup>1,2,5\*</sup>

\*A secondary endpoint based on Kaplan-Meier estimates of cumulative rate of time to onset of response. The median time to onset of response was 3 days in GL-1 (*Dysport*: 55/105 [52%]; placebo: 3/53 [6%]) and GL-2 (*Dysport*: 36/71 [51%]; placebo: 9/71 [13%]), and 2 days in GL-3 (*Dysport*: 110/200 [55%]; placebo: 4/100 [4%]).<sup>1,2,5</sup>

Please see full Important Safety Information,  
including Distant Spread of Toxin Effect Boxed  
Warning, on pages 6 and 7.

"I WANTED TO  
*smooth the frown lines*  
BETWEEN MY BROWS WITHOUT ADDING  
TO MY TO-DO LIST EVERY MONTH."

-Rebecca, 35, Actual *Dysport* Patient,  
Mom, Volunteer & Full-time Project Manager

# INDICATION

*Dysport*<sup>®</sup> (*abobotulinumtoxinA*) is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

# IMPORTANT SAFETY INFORMATION

**What is the most important information you should know about *Dysport*? Spread of Toxin Effects:** In some cases, the effects of *Dysport* and all botulinum toxin products may affect areas of the body away from the injection site. Symptoms can happen hours to weeks after injection and may include swallowing and breathing problems, loss of strength and muscle weakness all over the body, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, or loss of bladder control. Swallowing and breathing problems can be life threatening and there have been reports of death. You are at the highest risk if these problems are pre-existing before injection.

**These effects could make it unsafe for you to drive a car, operate machinery, or do other dangerous activities.**

**Do not have *Dysport* treatment if you:** are allergic to *Dysport* or any of its ingredients (see the end of the Medication Guide for a list of ingredients), are allergic to cow's milk protein, had an allergic reaction to any other botulinum toxin product, such as Myobloc<sup>®</sup>, Botox<sup>®</sup>, or Xeomin<sup>®</sup>, have a skin infection at the planned injection site, are under 18 years of age, or are pregnant or breastfeeding.

**The dose of *Dysport* is not the same as the dose of any other botulinum toxin product and cannot be compared to the dose of any other product you may have used.**

**Tell your doctor about any swallowing or breathing difficulties and all your muscle or nerve conditions,** such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis, or Lambert-Eaton syndrome, which may increase the risk of serious side effects including difficulty swallowing and difficulty breathing. Serious allergic reactions have occurred with the use of *Dysport*. Dry eye has also been reported.

**Tell your doctor about all of your medical conditions, including** if you have surgical changes to your face, very weak muscles in the treatment area, any abnormal facial change, injection site inflammation, droopy eyelids or sagging eyelid folds, deep facial scars, thick oily skin, wrinkles that can't be smoothed by spreading them apart, or if you are pregnant or breastfeeding or planning to become pregnant or breastfeed.

**Tell your doctor about all the medicines you take,** including prescription and nonprescription medicines, vitamins and herbal and other natural products. Using *Dysport* with certain other medicines may cause serious side effects. **Do not start any new medicines while taking *Dysport* without talking to your doctor first.**

**Especially tell your doctor if you:** have received any other botulinum toxin product, such as Myobloc<sup>®</sup> (*rimabotulinumtoxinB*), Botox<sup>®</sup> (*onabotulinumtoxinA*), or Xeomin<sup>®</sup> (*incobotulinumtoxinA*), in the last four months or any in the past (be sure your doctor knows exactly which product you received), have recently received an antibiotic by injection, take muscle relaxants, take an allergy or cold medicine, or take a sleep medicine.

## Common Side Effects

The most common side effects are nose and throat irritation, headache, injection site pain, injection site skin reaction, upper respiratory tract infection, eyelid swelling, eyelid drooping, sinus inflammation, and nausea.

Ask your doctor if *Dysport* is right for you.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see *Dysport* Full Prescribing Information including Medication Guide at [DysportUSA.com](http://DysportUSA.com).

## References

1. Data on file. GL-Y-97-52120-719 (GL-1) Clinical Study Report. Fort Worth, TX: Galderma Laboratories, L.P.; 2007.
2. Data on file. GL-Y-97-52120-718 (GL-3) Clinical Study Report. Fort Worth, TX: Galderma Laboratories, L.P.; 2007.
3. Data on file. MA-35497. Post Hoc Analysis. Fort Worth, TX: Galderma Laboratories, L.P.; 2017.
4. Molina B, Grangier Y, Male B, et al. Patient satisfaction after the treatment of glabellar lines with Botulinum toxin type A (Speywood Unit): a multi-centre, prospective, non-interventional European observational study. *J Eur Acad Dermatol Venereol*. 2015 JUL29(7):1382-1388.
5. Data on file. GL-Y-97-52120-085 (GL-2) Clinical Study Report. Fort Worth, TX: Galderma Laboratories, L.P.; 2007.
6. Deem M. *Dysport*: What You Should Know. Real Self Website. <https://www.realself.com/dysport>. August 23, 2019.

Dysport.  
(abobotulinumtoxinA)


ASK YOUR DOCTOR OR SPECIALIST  
ABOUT *DYSPORT* TODAY.

Actual patient. Individual results may vary.

“I WANTED TO FEEL CONFIDENT  
*about my look*  
WITHOUT THE FROWN LINES.”

-Deven, 39, Actual *Dysport* Patient,  
Director of Events & Marketing

**realself.**



RealSelf Worth It rating<sup>6</sup>

97 percent of *Dysport* users\* would get it again.<sup>4†</sup>

\*User = clinical trial subject.

†Subject's satisfaction and self-perception reported 3 weeks after treatment (n=531).

Please see full Important Safety Information,  
including Distant Spread of Toxin Effect Boxed  
Warning, on pages 6 and 7.

Please see *Dysport* Full Prescribing Information  
including Medication Guide at [DysportUSA.com](https://www.dysportusa.com).

© 2022 Galderma Laboratories, L.P. The *Dysport* trademark is used under license.  
All trademarks are the property of their respective owners. USMP/DYS/0297/1119

GALDERMA

EST. 1981