

# **Guideline**

# Cleaning of the Draeger neonatal incubator and Paediatric Paraid Trolley.

# 1 Scope

For use within the Paediatric and Neonatal Decision Support and Retrieval Service (PaNDR) for the East of England.

# 2 Purpose

To provide guidance for the PaNDR team in the safe and effective general and deep cleaning of the Neonatal Draeger Isolette transport incubator and the Paediatric Paraid Trolley.

#### 3 Neonatal

#### 4 Definitions and abbreviations

- **General cleaning** of the Neonatal transport incubator carried out after every transfer with Tristel Jet solution.
- **Deep cleaning** of the Neonatal transport incubator– carried out with Tristel Jet after transfer of every infant with known colonization or infection. Incubators should be deep cleaned on a monthly basis by the NICU assistants and recorded in the monthly check book and the board in the clinical room.
- **Tristel Jet** -a sporicidal disinfectant that can be used on surfaces and equipment located near to the patient. To use spray at least two doses on a wipe, wipe entire surface and ensure surface remains wet for 1 minute and leave to dry. Once open use within 6 months.

#### 5 Introduction

Cleaning is routinely undertaken by the neonatal clinical assistants on the Neonatal Unit for the Neonatal PaNDR team. However, cleaning may be required to be undertaken between transfers if the team is unable to return to base. This is the responsibility of all members of the PaNDR team.

#### Gloves and aprons must be worn.

• The transport incubator must be cleaned after every transfer to minimise the risk of spread of infection between infants.



#### Addenbrooke's Hospital

- Deep cleaning of the incubator is required after transferring an infant with MRSA, ESBL, CPE and/or COVID-19
- If necessary, general and deep cleaning can take place in the referring/receiving unit before the next transfer.

# 6 General cleaning

Turn off the incubator and unplug from the mains.

- Open the portholes and doors and remove the cuffs and silicone access inserts. Leave to soak in hot soapy water whilst the rest of the incubator is being cleaned with the Tristel Jet solution.
- Remove the entry port sleeve from the end door access port and discard.
- Remove the positioning bumper, Z-Flo mattress, clean and ensure these are completely dry.
- Wash the tray and perspex incubator hood inside and out and dry thoroughly.
- Wipe monitoring equipment, leads and infusion pumps with Tristel Jet solution and dry thoroughly.
- Take the door cuffs and silicone access inserts from the soapy water and leave to air dry.
- Reassemble and make up incubator with clean linen for next transfer.

# 7 Disassembling the incubator for deep cleaning

Turn the incubator off and unplug from the mains.

- Open the portholes and doors and remove cuffs and silicone inserts.
   Leave to soak as above.
- Remove the entry port sleeve from the end door access port and discard.
- Open the main door and remove the Z-Flo mattress and set aside for cleaning.
- Press the metal mattress tray-stop beneath the tray and slide the tray off the base through the end access door.
- Reclose and secure the access doors.
- Remove the four pumps and the Neopuff from above the incubator hood.
- Release the four black rubber hood clips located at the front and rear corners of the incubator.



#### Addenbrooke's Hospital

- Fully lift both hoods together off the base and set aside on a trolley.
- Unclip the four corners on the cream incubator base and lift off, revealing the fan and metal heat dispersers.
- Sterilization of any component of the transport incubator is generally not necessary, however cold liquid sterilization can be used if required.

## 8 The deep cleaning process

#### 8.1 The incubator base

- Remove any visible lint/ debris and wipe the base with Tristel solution.
- Take extra care to clean along the rubber seals around the edge of the base where the hood sits.

#### 8.2 The upper and outer hood shells

- Clean all surfaces thoroughly with the Tristel solution.
- Dry with a clean cloth or paper towel ensuring there are no smears.
- Inspect the 'feet' on the underside of the upper shelf for damage.

#### 8.3 Mattress tray, mattresses and rails

- Use Tristel Jet solution to clean all the surfaces and rails thoroughly.
- Dry with a clean cloth or paper towel.

#### 8.4 Trolley base and shelf

Wash the trolley base, rails and shelf and dry thoroughly.

# 9 Reassembling after cleaning

- Lower the cream base back on and clip it back in place. Carefully place the inner hood shell into the outer shell, ensuring the port-holes and inner shell doorway are aligned.
- Install both hoods together onto the base ensuring inner and outer shells are sitting securely in the grooves around the edge of the base.
- Secure the black hood clips.
- Replace the restraint rails on the mattress tray.
- Place both mattresses on the tray and slide the tray back in (clear plastic foot end first) until it clicks over the release tab and stops at the right end of the base.



#### Addenbrooke's Hospital

- Replace the tubing access inserts and door cuffs with a clean, previously washed and dried set.
- Install a new entry port sleeve on the end door access port.
- Replace the Draeger incubator filter located at the back of the incubator.
- Replace the pumps and Neopuff.
- Remake the incubator with clean linen.
- Plug the incubator back into the mains, turn it on and check the set temperature is 36°C.

#### **Paediatric**


The Paediatric trolley must be cleaned after each patient use by wiping down all surfaces with clinell wipes. A deep clean is required if transferring a child with MRSA, ESBL, CPE and/or COVID-19. Cleaning after use is the responsibility of the clinical team. A monthly deep clean should be arranged and it has been agreed that the neonatal assistant will assist with this.

## **Deep Cleaning of the Paediatric Paraid Transfer Trolley**

For a thorough deep clean of the Paediatric Paraid Transfer Trolley, follow these steps:

#### Remove the Mattress:

- The mattress consists of three parts: the patient head end, the main body, and the foot end.
- To remove the mattress, slide the rod underneath each section down from the smaller hole into the larger hole in the keyhole slot, then lift it out.





#### **Clean the Straps:**

 The straps attached to the main body of the trolley should be wiped down using a suitable cleaning solution, such as Tristel Jet. Ensure all parts are fully dried.

#### **Monitoring Shelf:**

 There is a monitoring shelf located at the foot end of the trolley, above the mattress. This shelf should be wiped down with the cleaning solution. No equipment needs to be removed from this shelf—just wipe around the area thoroughly.

#### **Shelf and Equipment Cleaning:**

- The bottom shelf of the trolley holds oxygen cylinders, pumps, humidifiers, and INomax®. These items should be carefully removed where possible to allow access for cleaning the parts of the shelf that they sit above.
- The INomax® does not need to be removed. It only needs to be unclipped and lifted up to allow someone to wipe underneath it.
- The humidifier is secured within a fixed cage attached to the trolley, which cannot be removed. Clean both the humidifier cage and the surrounding area thoroughly, ensuring all surfaces are wiped down and dried.
- Once the equipment has been lifted or removed, wipe down the shelf with a suitable cleaning solution, ensuring it is fully dried before reattaching or repositioning the equipment.

#### Reassembly:

- Once cleaning is complete, reattach the mattress by aligning and securing it using the keyhole slot locking mechanism.
- Reattach or reposition the pumps, INomax, humidifier, and other equipment back onto the trolley.

# 10 Monitoring compliance with and the effectiveness of this document

The PaNDR team will monitor compliance with this document by undertaking regular audits

The effectiveness of the document will be monitored by review of any reported incidents.

#### 11 Associated documents

- Draeger Isolette manual
- Tristel Jet Safety Data

Pandre & Recrieval Screen

Addenbrooke's Hospital

# **Equality and diversity statement**

This document complies with the Cambridge University Hospitals NHS Foundation Trust service equality and diversity statement.

#### **Disclaimer**

It is **your** responsibility to check against the electronic library that this printed out copy is the most recent issue of this document.

# **Document management**

Approval:	PaNDR Consultants 2024		
Owning department:	PaNDR		
Author(s):	Rhonda Young & Patrick Sheppard		
Pharmacist:	n/a		
File name:	Cleaning of the Draeger neonatal incubator version 3 October 2024.doc		
Supersedes:	Version 2 March 2021		
Version number:	3	Review date:	October 2027
Local reference:		Document ID:	100271

Cambridge University Hospitals NHS Foundation Trust

PanDR

Pactiatric & Neonatal Decision

Support & Retrieval Service

Addenbrooke's Hospital

# Appendix 1: Use of Tristel Fuse first aid measures/ Exposure controls/personal protection

Use of Tristel Fuse First aid measures:		
Inhalation:	Non-toxic Non-toxic	
Eye contact:	Rinse eyes with water	
Skin contact:	Wash affected area with water	
Ingestion:	Do not induce vomiting. Give water to drink	
	Seek medical advice where necessary	

Exposure controls/personal protection:		
Personal protection:	Avoid eye contact and prolonged skin contact	
	Gloves and apron recommended	
Skin contact:	Low risk	
Eye contact:	Low risk	
Inhalation:	Low risk	
Ingestion:	Low risk, substantial ingestion may cause	
	discomfort to mouth and digestive tissues	