



BROADLOOM SEDGE CONTROL

NOTE: Read the entire label directions for use, container disposal, conditions of warranties and limitations of liability before using this product.

The label is the law.

BROADLOOM® SEDGE CONTROL is a postemergence herbicide for residential and small area use to control broadleaf weeds, annual sedges and yellow nutsedge in established turfgrass and ornamentals. *Broadloom does not control purple nutsedge.*

Product Information

Broadloom controls more than 30 broadleaf weeds. Broadloom Sedge Control does not control grasses. Broadloom Sedge Control is effective mainly through contact; therefore, all plants must be thoroughly covered with spray.

Use Broadloom Sedge Control on **established** bluegrass, fescue, bentgrass, Bermudagrass, bahiagrass, centipedegrass, zoysiagrass, ryegrass, and St. Augustinegrass.

Broadloom Sedge Control is effective mainly through contact; therefore, all target plants must be thoroughly covered with spray

Low Application Rates for Established Turfgrass

Broadloom Sedge Control application rates 0.55 fl. oz. (3.5 tsp.) per 1,000 sq. ft. 0.75 fl. oz. (5 tsp.) per 1,000 sq. ft. Do not apply more than 1.5 fluid ounces of Broadloom Sedge Control per 1,000 square feet (4 pints per acre) in one season.

- Spray areas with Broadloom Sedge Control in water as the spray carrier. 200 sq. ft. 0.15 fl. oz. (1 teaspoon) 0.2 to 0.4 gallons of water (1.6 to 3.2 pints)
- 400 sq. ft. 0.3 fl. oz. (2 teaspoons) 0.4 to 0.8 gallons of water (3.2 to 6.4 pints)
- 1,000 sq. ft. 0.75 fl. oz. (5 teaspoons) 1 to 2 gallons.
- Broadloom is tank mix compatible with commonly used pesticides.
- Broadloom is rainfast or sprinkler irrigation resistant eight (8) hours post application.
- Do not cultivate or mow seven (7) days post application.
- Refer to product label for sensitive and susceptible ornamental plant types.
- Do not apply Broadloom Sedge Control to turf that has been under stress (drought, cold temperature or injury from other herbicides or pesticides).
- Do not apply Broadloom Sedge Control to any newly seeded or newly sprigged turf until seedlings or sprigs are well established, as injury may result.
- Broadloom Sedge Control may cause a speckling or bronzing but should outgrow these conditions in ten (10) days.
- In perennial ryegrass, apply no more than 0.75 fluid ounces per 1,000 square feet (2 pints per acre) at one time and make subsequent applications no less than 21 days later.

DIY Pest and Lawn - 386-774-2847

2417 Enterprise Rd., Orange City, FL, 32763

WARRANTY AND CONDITIONS OF SALE - Seller makes no warranty, expressed or implied, concerning the use and handling of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.