



Headway® G Fungicide

Group 3 & 11 fungicides

For the control of certain pathogens causing foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass plants.

*** See label for complete list of diseases controlled.**

Headway G may be used on turf on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields on warm and cool season grasses. **Do not apply above 90 degrees fahrenheit.** Headway G is a minimal dust formulation.

Apply Prior to Disease Development - Repeat applications at specified intervals for as long as required.

Do not apply more than 19.2 lbs. per 1,000 sq. ft. per year.

Applications may be made by ground only.

Headway G is a combination product of Group 33 (propiconazole) and a Group 11 (azoxystrobin) fungicide. Headway G displays systemic properties providing curative and preventative control. Reapply Headway G on a 14 - 28 treatment interval. Rotate to another fungicide Group after two consecutive applications of Headway G; such as T-Methyl 50 WSB fungicide. Consult your local or state turf authorities for resistance management strategies that are complementary to those in the label. Maximum of 3 pounds per 1,000 square on bermuda grass.

Water in Headway G with 1/4 to 1/2 inch or before a suitable rain to activate the product.

Average rate for Headway G is 2 - 4 pounds per 1,000 square feet. Refer to the product label for correct userates of Headway G fungicide.

Common spreader settings are 5 1/2 for established turf and 8 1/4 for bare ground applications. Refer to the product label for correct spreader settings.

IMPORTANT

- Read, understand, and follow manufacturers label instructions.
- Container Disposal: Follow manufacturer's label instructions for container disposal.

DIY Pest and Lawn - 386-774-2847

2417 Enterprise Rd., Orange City, FL 32763