

# PT Ultracide, Bifen I/T, Precor IGR Flea Kit





Flea & Flea larvae



## DIRECTIONS FOR USE IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

PT Ultracide at the rate of 14 oz for up to 1827 sq ft. Carpeted areas may be vacuumed before treatment and after treatment has dried. Treat infested areas or areas which could be infested; these include rugs, carpets, upholstered furniture, pet beds and pet resting areas. Hold can at arms length and direct spray toward the area to be treated. Use a sweeping motion to apply product and back away from the treated area while holding the can 36 inches away from the surface being treated. An area of 80 - 100 sq ft can be treated in approximately 10 seconds with PT Ultracide. When treating upholstered furniture, treat under cushions and areas where flea development occurs. Cover aguariums and fish bowls and remove birds from the area prior to treatment. Treated areas should be vacated during application. TO CONTROL TICKS AND FLEAS: Old bedding should be removed and replaced with fresh, clean bedding after treatment. Pets should be treated with a registered product for flea and tick control in conjunction with this product. SHAKE WELL BEFORE USE. PT Ultracide kills fleas and prevents flea infestations for a full season. Adult fleas may be seen in the treated areas when brought in on infested animals or when adults emerge from pupal cases; however, a population will not develop in the treated area. In transportation equipment, certain plastics may be whitened or dulled. Apply only to carpets and cloth upholstered areas, or test an inconspicuous area before treating.

# Except for the applicator, DO NOT PERMIT HUMANS OR PETS TO CONTACT TREATED SURFACES UNTIL THE SPRAY HAS DRIED.

DO NOT TREAT PETS WITH THIS PRODUCT. NOTICE.PT Ultracide has been tested on several types of carpet, fabric, and other household furnishings without adverse effects. In a few instances, waxed surfaces have been whitened or dulled.Because of the wide variety of floor types and finishes, treat a small inconspicuous area before treating the entire area. Avoid contact with antique finishes.In a few instances finishes have been whitened or dulled. Holding the can 36 inches away from the target surface normally prevents any whitening from occurring.



#### Bifen I/T and Precor IGR

1st time mix 2 oz of Bifen I/t + 1 oz Bottle of Precor IGR with 1 gallon of water into a pump sprayer. Treat the interior of home along baseboards under kick plates in the kitchen, behind the stove, under the dishwasher and behind the refrigerator. Treat the garage as well as the entire exterior perimeter of the home. (As for the exterior perimeter, spray 2ft up wall of home and 2ft off wall of home)also around the faucets, Dryer vents, windows, eves and anything that may be close to or touching the exterior of the home including; trees, brush and plants etc... Day 7, 14, use 1 oz of Bifen I/t - 1 gal of water treat the same locations.

Flea larvae: Immature fleas mature in shaded areas accessible to pets or other animals. When treating these areas use a higher volume treatment so that the insecticide penetrates into the soil. NOTE: If adult fleas on lawn areas are being controlled by applying Demon Max at a rate of 1 fl. oz. per 1000 square feet, then the rate of larval application can be accomplished by a two-fold increase in spray volume.

### Repeat on the days 7, and 14.

Eliminating a flea infestation will mean treating all places that can be infested, including your pet, your home and your yard. Failing to treat all of the areas will likely lead to treatment failure. We recommend starting with your pets and then treating the home and yard. The treatments should be done around the same time for the best results.



**DIY Pest and Lawn 386-774-2847** 

### 2417 Enterprise Rd. Orange City FL, 32763

WARRANTY AND CONDITIONS OF SALE Seller makes no warranty, expressed or implied, concerning the use and handling of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.