

APTDG25DM5.3

# **Dimension 0.25G**

### **FEATURES & USE TIPS:**

- · For use on Turf, Landscape Ornamentals and Nursery Ornamentals
- Provides preemergent control of crabgrass and other listed annual grasses and broadleaf weeds.
- Provides early postemergent crabgrass control in limited conditions.
- · Consistently sized particles reduce segregation potential

### PRODUCT SPECIFICATIONS:

Active Ingredient: Dithiopyr (0.25%)

PRODUCT
CHARACTERISTICS:

PRODUCT CODE:

Date: 060614

Supersedes: 000000

Product Type:
Dimension® Herbicide

Particle Size: SGN: 150

Net Weight: 50 lbs.

Coverage: 14,700 sq. ft. @ medium rate

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC

To order, contact your local Andersons Golf Products distributor.

For more information, contact Andersons at our website: www.andersonspro.com or call 1-800-225-2639.

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.



# he Professional ndersons Turf™ Products **Professional**

## Dimension® 0.25G

#### For use on Turf, Landscape Ornamentals, and Nursery Ornamentals On DG Pro<sup>®</sup> Carrier

For season long control or suppression of listed annual grasses and broadleaf weeds in omamental landscape areas, estab-lished lawns and omamental turf (including golf course fairways, roughs and tee boxes and sports turf) and container and field crown nurser ornaments<sup>1</sup>.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 4.6 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre); and is prohibited from use in Nassau and Stiffit (Yoursies

\*Dithiopyr OTHER INGREDIENTS... TOTAL... CAS No. 97886-45-8

\*This product contains 0.125 pounds of active ingredient, diffuory, per 50 pound bag.
Product protected by U.S. Patient No. 415(2):180. Other patients pending.
EPA Reg. No. 91982/13
PER St. No. 91980-14, 9199-01-28, 9198-NL-0014
Underlined letter is first letter used in nan code on bag
DDECALITIONAL NO. 600-11

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS KEEP OUT OF REACH OF CHILDREN

CAUTION

AU II UN ses sey initiation. Hamful if inhaled. Avoid contact with eyes or dothing. Avoid breathing dust. Wash thoroughly with and water after handling. Remove contaminated dothing and wash before reuse. Protonged or frequently repeated ontact while handling this material may cause allergic reaction in some individuals.

FIRST AID	
f in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
finhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
f swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
f on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.  Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Non-Agricultural Use Requirements used from product flat are NOT within the scope of the Worker Protection Sharked fit agricultural protection of the product flat are NOT within the scope of the Worker Protection Sharked fit agricultural protection is OCFR hear 170. The WPS agrices where the product is used to produce agricultural prints on them, mannesses or generobases to bot retainer of lower for the three belows used of dash to be selfect. In addition, of directions for one exquare vesterings, do not enter or the other three belows used of dash to selfect. In addition, of directions for one exquare vesterings, do not enter or low others to enter the feeting and except protections.

STORAGE AND DISPOSAL
STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water,
od, or feed by storage and disposal. Keep container dosed and away from food, feedstuffs, and domestic water supplies PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal in accordance with applicable Federal, state or local p

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if avails Completely empty beg into application equipment. Then dispose of empty beg in a sanitary landfill or by incinera allowed by State or local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to "pesticide disposal" directions above

\*\*TUNPFAMES SAFETY\*\*

This product may be used on seeded, sodied, or springed lawns and omamental larfs that are well-established. The gases must have developed a good not objecte mark a uniform stand, and have received it least two movestips of the seeders, sodiers, or storp in selection, sodiers, sodiers, or storp in selection, sodiers, sodi

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and Washington' may result in undesirable turforass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

\*\*\*DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

EDING, OVERSEEDING, OR SPRIGGING fing, overseeding, or sprigping of treated areas with this product should be delayed until 12 weeks from the firm colors. Research, overseeding or sprigging before 12 weeks after application may prevent establishment of its furgissess. When researching or overseeding, proper cultural practices such as soil culturation, ringition and four invalid be followed. If overseeding with perential regisses in the fail, overseed 2 weeks or later after a striple.

application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

OTHER USE PRECAUTIONS
The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass is faulted before crabgrass and continued in the leaf or first titler.

For best results, cultural practices that disturb the soil, such as core-spike- or hydro-serification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Covincut or conditional statements and any Prememergence and Early Postemergence and Early Postemergence Control of cabgass (including the large, smooth, and southern species) where applied prior to the emergence of car It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the first tillering of crabgrass, which generally corresponds to the time when you ca

Spring Applications:

Where split fertilizer applications are listed, the rates in Table 1 may be split across two applications made 6-10 weeks

Wee	Weeds Controlled						
Barley, Brome, Crabgra Dallisgrass (seedling), F	108 - 160 lbs./acre 2.4 - 3.6 lbs./1,000 sq ft 0.27 - 0.40 lbs. al/acre						
Bamyardgrass Chickweed Clover, hop Crabgrass, tropical Crowfootgrass Dichondra, Carolina Lespedeza, common Kikuyugrass Many's grass Marestail Medic, black	Muberry weed Mustard Cots, wild Oxals Buthercup Creeping Woodsorrel Yellow Woodsorrel Pineappleweed Ryegrass, annual Ryegrass, perennial Sandour	Smutgrass Speedwell, com Spurge Garden Prostrate Spotled	160 – 200 lbs/lacre 3.6 – 4.6 lbs/1,000 sq ft 0.40 – 0.50 lbs. al/acre				
Bluegrass, annual Bitlercress Carpetweed Chickweed Geranium Carolina Goosegrass	Henbit Knotweed, pr Parsley-piert Pigweed, red Purslane, cor Rocket, Lond Shemardsou	200 lbs./acre 4.6. lbs./1,000 sq ft 0.50 lbs. ai/acre					

Use the higher rates where weed infestations are dense. Also use the higher rates if applications are made early and longer residual control is desired.

Fall Applications
This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate pering application to provide season-horg control.

REGION	Preemergence
North	3.4
Transition	4.6*
South	4.6*
Coastal South	4.6*

ensure equipment is functioning properly and appropriate (appropriate property). ORNAMENTALS (Landscape, Field Grown and Container Grown)
This specially herbiding provides presempency control of lated annual grass and broadlesf weeds in areas plantend with tolerant ornamental plants lated on this label. It is intended for use on plants grown for seaffeite purposes
where the property of the pro

USE RATES
Apply his product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following lable for product recommendations when treating a small area.

DO NOT apply more than 200 lbs. per acre per application and no more than 3 applications (600 lbs.) per acre per year

ork State, this product may only be used by Professional Applicators at no more than 200 lbs. per acre ds per 1,00 sq. ft./ 0.5 lbs. active ingredient per acre) per year; and is prohibited from use in Nassau to Counties

APPLICATION TIMING AND RECOMMENDATIONS
This product is a preemingence herbicade that will not control established weeds. Make applications prior to week end germation for to be any pour. The herbicade that week end control is obtained when applied to sold that is free of dods, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the sold surface as the herbicide better will be broken.

- long agony has product to establishment of minimized.

- long agony has product to establishment of minimized.

- long agony has product to establishment of minimized to establish the product to the plant route of long and that seek as he no create that it is been and to be taken from the contract of this product to the plant route of period part may may contract the product to the plant route of period part may may contract the product to the plant route of period part may may contract may occur.

- In ord party amount or mannership prints that are used as the bid could be found to the product of period product of the product of period product of the prod

tapped for syrup).

Do not apply this product in enclosed structures and greenhouses.

Do not apply another preemergence herbicide within 4 weeks of application

Treatment of Omamental spacies not listed on this label:

Treatment of Omamental spacies not listed on this label:

Treatment of Omamental spacies or under all post
spacies or varieties or under all post
slabe governor conditions, blass who wish his use this product on ornamental spacies and listed or this label may
determine the sublishing for use by breating a real number of omamental points at listed real. Perior to tear
of larger ease, treated plants should be observed for any symptoms of herboical stry, such as fortal changes
readed vigor of stand exclusion, along 50 to 00 step of round groups condition to before the first
the standard plants should be observed for any symptoms of herboical stry, such as fortal changes
the standard plants should be observed for any symptoms of herboical stry, such as fortal changes
the standard plants and the standard plants are standard to the standard plants are standard to the standard plants are standard to the standa

IMPORTANT: READ BEFORE USE:
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product.
If terms are not acceptable, return the unogened product container at once. By using this product, user or buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCHOOMING INTERING ULCENTER OF LIABILITY.

MARRANTED SECTION TO THE DISCHARGE AND LIMITATION OF LIABILITY.

MANUFACTION OF THE DISCHARGE AND LIMITATION OF THE BRIDGE OF THE

LEGAL RIGHTS:
THE LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION TO STATEJURISDICTION.

Visit our website: www.AndersonsPro.com @2014 The Andersons, Inc. The Andersons logo, DG Pro and Professional Turf are tran Dimension is a registered trademark of DowAgroSciences

P.O. Box 119 Maumee, OH 43537

APTDG25DM5.3 APTDG25DM5 3B

Net Wt. 50 lbs. (22.68 kg)



#### Suggested Spreader Settings

These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product 50 lbs. treats 20,800 sq. ft. (0.48 acre) at the Low Rate (2.4 lbs/1,000 Sq. Ft.) 50 lbs. treats 14,700 sq. ft. (0.34 acre) at the Medium Rate (3.4 lbs. / 1,000 Sq. Ft.) 50 lbs. treats 13,800 sq. ft. (0.32 acre) at the High Rate (3.6 lbs./1,000 Sq. ft.) 50 lbs. treats 10,800 sq. ft. (0.25 acre) at the Heavy Rate (4.6 lbs./1,000 Sq. ft.)

			SPREADER SETTINGS							
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE 2.4 lbs/1,000 sq. ft.	MEDIUM RATE 3.4 lbs/1,000 sq. ft	HIGH RATE 3.6 lbs/1,000 sq.ft.	HEAVY RATE 4.6 lbs/1,000 sq. ft.				
Andersons Model 2000 / AccuPro 2000; (cone 6) Andersons 2000 SR / SR-2000 (cone 6)	3 mph	11 feet	К	L 1/4	L 3/4	M 1/2				
Scotts R-8, R-8A (cone 6)	3 mph	11 feet	J	K	K 1/4	L 1/2				
LESCO 020093 (0.25)	3 mph	7 Feet	D	E	E 1/4	F				
LESCO 705698 (0.25)	3 mph	7 Feet	10	12	12 1/2	14				
Lely Models WTR, WFR, HR, 125- (PTO at 450 rpm)	4.5 mph	34 feet	4 1/4 IIIC	5 1/4 IIIC	5 1/2 IIIC	6 IIIC				
Vicon (03 Series)	4.5 mph	30 feet	18	23	26	29				

180				ptable on Method oy an (x)	Field/Land Contain Ornamentals I	ation to dscapes or er Grown Noted by an (x)			Accer Application Noted b	on Method	Field/Lan Contain Ornamentals	Noted by an (x) Field/		1	Acces Applicatio Noted b	on Method	Containe Ornamentals N	Noted by an (x) Field/
Company			Over the Top		Container Grown	Landscape Grown		Tolerant Cultivars	Over the Top	Directed	Container Grown	Landscape Grown		Tolerant Cultivars	Over the Top		Container Grown	Landscape Grown
Mathematical	abelia, dwarf (Abslia x grandiflora)	nana		×		х				x		x	(Acer saccharinum)					x
Company		maureli	x	×	×	x	(Farana 6 days)	Auereo-		x		x	(Acer saccharum) marigold	honeycomb	-			
Company   Comp	acacia redolens		х	×	x	х	(Euonymus ionunei)	colorata		x		x	(rageres parura)	wheeleris				x
	Ajuga carpet bugle							emerald n' gold				x x		golden snowflake				
Company	(Aiuga genevensis)			x x		x x				x x		x x	mondo grass	double white	-			
State   Stat	almond, flowering			x				Manhattan	×	x	x	x		sunnyside				х
The content				×		х		silver king				x	mountainash (Sorbus aucuparia)			x		x
Part	arborvitae	pyramidalis					eurone green leaved			×		x x	myrtle, crape (Lagerstroemia indica)	langer				
Marchane		smaragh				x x	(Euryops pectinatus) fan palm. European	VIIIUIS	_ ^	×	^		myrtle wax					X
Company	arborvitae, dwarf golden		x	X X	×	x x	(Chamaerops humilis) fan palm, Mexican			x			(Myrica californica) harcissus		-			- x
Company	(Thuja orientalis)						(Washingtonia robusta) fem (various)			х		х	(Narcissus spp.) oak, laurel		_	×		х
Company	(Fraxinus pennsylvanica)	autumn numle								х		х	oak, pin					x
Company	(Fraxinus americana)												(Quercus rubra)					X
March   Marc	(Callistephus chinensis) azalea	brilliant						nadia					(Quercus virginiana) bak, willow					- x
Company		buccaneer		х		x	(Ables fraseri) Forsythia			^			(Quercus phellos) bleander	hardy red				
March   Marc				x		x	(Forsythia viridissima)	bronxensis					osteospermum		1			
March   Marc		Elsie Lee exbury		x x			(Forsythia xintermedia) (Forsythia x 'Meadowlark')	lynwood gold		x		x	(Osteospermum fruticosum) pachysandra	33	$\rightarrow$	×		x
Mary		fashion hardijzer		x x						x		х	(Pachysandra terminalist) palm, King (Archontophoenix			×		х
Mary State		beauty		×			(Forsythia suspensa)	spring glory	×	x	x	x	cuninghamiana) palm, queen		x	×	x	x
Second		higasa hinocrimson		x x		x x		weeping rubrum	×	X	x	X	(Syagrus romanzofflanum) pampas grass			x		x
March Marc		Holland (hybrid)		x		x	(Penniserum setaceum) fuchsia (Funholo con I			х		x	pansy (Viola x wittrockiana)			×		x
Company		Marion Lee		x x			galium (Ga(ium ordoratum)			х		x	(Bougainvilles glabra)	Barbara Karst	x		x	х
Martin   M		orange cup orchid lights		x			gardenia (Gardenia jasminoides)	radicans		x x		X X	(Prunus persica) periwinkle, dwaff		-			X
March   Marc		snow		x x		x		wnite gem variegata	X X	X	X		(Vinca minor) petunia	picoti		x		×
March 2006   Mar	azalea, flame			×		×	geranium (Pelargonium x hortorum)			X			photinia, red tip (Photinia x fraseri)			х		х
Commonweight   Comm	(Rhododendron						gum (Eucalyptus citriodora)	nockers 17					pieris (Pieris taiwanensis)					×
March 1999	azalea, kirishima			х		x	(Crataegus spp.)	crimson cloud		x		×	pieris, Japanese	mountain fire				x
March   Marc	bamboo heavenly			x		х		Jack Evans		x		x			×		×	×
Company	barberry, Japanese	aurea					hawthorn, Indian	white ballering					(Pinus nigra) pine, Japanese black		_ ^		_ ^	
Property	ons institution(s)	green		x		×	(Rhaphiolepsis indica) heather, twisted	Mediterranean					(Pinus Thunbergiia) pine, lobiolly			x		×
Description		pygmy red				x x	(Erica cinerea) hemlock, Canada	pink					pine, longleaf (Pinus palustris)			x		x
Mary				x		X	([Suga canadensis] hisbiscus (Hibiscus con I						pine, mugo or Swiss Mt. (Pinus mugo)					x
The content of the	atropurpurea)						holly	hula girl		x		×	(Pinus sylvestris) pine, slash	-	-			X X
March   Marc	(Gaillardia grandiffora)						(llex x 'Nellie R. Stevens')	Nellie R. Stevens	x	х	x	x	(Pinus elliotti) pine, Virginia		-	×		х
March Annew Company	(Ammophila breviligulata)							fosteri		X X		x x	(Pinus virginiana) pine, white (Pinus etmbus)		х	×	x	х
Manuface	(Arctostaphylos uva-ursi)	Massachusetts						blue girl		X X		x x	pineapple, guava † (Feljoa sellowiana)			x		х
March   Marc	(Monarda didyma)							China girl	×	X X	x	X	(Pittosporum tobira)		x	×	x	×
Management   Man	(Begonia spp.)						(flex cassine) holly, Chinese (flex comute)	burfordii needlennint		X		X	plumbago, cape (Plumbago auriculata)	royal cape	х	×	×	х
Processor   Proc	(Betula nigra)	100					holly, Japanese (llex crenata)	compacta hellerie		x x		x x	(Potentilla fruticosa)	abbotswood		x		х
Marchan and	(Rudbeckia hirta)	goldstrum						northern beauty		x		x		nolden viceny		x		×
Finance   Fina	(Gaillardia spp.)		$\sim$				holly, yaupon	steeds	×	X	×	×	(Ligustrum japonicum)	regal texanum	×	x	×	x
Worker or   War	(Festuca ovina)													yellow tipped		X		X
Part	(Vaccinium spp.)	blue jay		x x		x	(Lonicera xylosteum)					×		onome		Y		Y
March   Marc		north blue		x x		x v	(Lonicera japonica) (Lonicera tatarica)	Canadian white		x		x	(Pvricantha koldzumii)	lalandei victory	×	x x	×	X X
Second	bottlebrush (Collistemon citrious)		x	x	x	x	hosta (Hosta sieboldii)			×		×	(Arecastrum			x		x
Second	bougainvillea			x		х	(Hosta lancifolial)	albo marginata	×	X X	x	X X	quince, Japanese † (Chaenomeles Japonica)			х		х
Base contention			×	×	×	×	foe plant (Droşanthemum floribundum)							Cunningham		X X		X X
Manufacture	(Buxus microphylla)		¥		×				×	X	×			PJM		x		x
Control of the cont	(Buxus microphylla japonica)						(Impatiens spp.)	dwarf blue		х		х	rhododendron, Carolina		-	X X		X X
Common processing	(Buxus sempervirens)							Bulgaria thomdale		X X		X X	(Rhododendron					
Commission of the commission	(Cytisus scoparius)			X X			jasmine, Asian (Trachelospermum			x		x	(Rhododendron			*		
Commissioned   Comm	camelia						jasmine, står (Trachelospermum			х		х	rosebay			×		х
Commission and	(Camellia japonica)	debutante mathotiana	¥	X	×		jasminoides) juhiper	Arcadia		x		x	nbbon grass		_	x		x
March   Marc	(Camellia sasangua)		¥		×	×	(Juniperus spp.)	bar harbor				x	rockcress (Arabis caucaisca)					x
Content of the cont	candytuft	setsukgekka						blue Pacific		X X		x						X
Clase and	(Iberis sempervirens)							blue star		X		x	hybrid) rosemary †		<u> </u>		_ ^	X
Application   Section   Process	(Carex spp.)			<u> </u>		<u> </u>		buffalo calgary carpet		x x		x	(Rosmarinus officinalis)	nana				x
Separate process of the control of	(Ajuga reptans)							emerald sea emerald		x		x	salvia (Salvia farinacea)					х
Contents on protection	oedar, red							endora com		x		x		red				×
ordinate and analysis of the processor o	pelosia			x		х		fruitlandi		x		x	smoketree, royal pumla		-			X X
Content	centaura (Centaurea montana)							gold tip		X X		x	(Cotinus coggygria) snapdragon					х
Control   Cont	cockscomb, plumose (Celosia cristata)	plumosa						hughes		X y		x		sterile	х		х	х
Total	coleus (Coleus blumei)							parsoni pfitzeriana		x x		x x		fanali				x
confidence perspherence of the confidence perspherence perspherence of the confidence perspherence perspherence of the confidence perspherence perspective perspherence perspherence perspherence perspherence perspherence perspherence perspherence perspective perspherence perspective perspec	columbine (Aquilegia spp.)							plumosa Prince of					(Astilbe X arendsii) spiraea	Anthony	-			X
Comparison and	coneflower, purple (Echinacea purpurea)	magnus purple	x	x	×	×		procumbens						Waterer red dolchica		x		x
Composing   Comp	copper leaf (Acalypha wilkesiana)			x		x		San Jose		x		x		froebeli pink goldenflame		X		x x
Contracting of	coreopsis (Coreopsis spp.)	moonbeam						sargent green		x		x		snowmound		x		x
Contended	com flower (Centaurea spp.)							scopulorum						Van Houtte				x
Bacchant plants	cotoneaster (Cotoneaster apiculatus)							spartan		x			spiraea, garland (Spiraea x arguta)	willing				х
Cycle	coyotebrush (Baccharis pilularis)							weberi		x		x x	spruce, Black Hilfs (Picea glauca var densata)	nlauro				X
Cyperacy Indian   Space   Sp	cycad			x		x	juniper	Yukon belle gold coast	X		x			gunda	_ ^		_ ^	X
Comman Agent Profession   Special	cypress, bald (Taxodium distichum)						(Juniperus chinensis)	torulosa-Holly					(Pices ables) spruce, white	conica				×
Commandation   Comm	cypress, Italian	glauca		х		x		wood					(r/icea grauca) sweetflag grass (Acorus calimus)					х
Cuprescryopies lysered    Cuprescryopies l	cypress, Hinoki false	gracilis		x		x	juniper (Juniperus procumbnes nana)	Japanese	х	х	x	х	sweetgum (Liquidambar styraciflua)					х
Culpress opening   Culpress op	cypress, leyland			х		х	juniper		х	х	х	х	sycamore, American (Platanus occidentalis)	nuby olow				
Minchans spp.)	hybrid	Kina Atta- 1					(Juniperus sabina L'tamariscifolia')						(Leptospermum scoparium) tree fern (tiki fern)	,uuy giuW			_ ^	X
Prince of the	(Narcissus spp.)						(Juniperus scopulorum)		х	х	x	х	(Asparagus virgatus) trumpet flower or Carolina					×
Project Control (Note of Control (Note	ивутту (Hemerocallis spp.)	bright yellow					king palm (Archontonhoenix	anyrotitet	х	х	х	х		apeldoom	-	x		¥
Commanda		single gold		x		x	cunninghamiana) laurel, Carolina cherry	bright n' tiaht	x	x	x	x	(Tulip spp) tufted hairgrass					- x
Communication   Communicatio		Wilson's		x		x	(Prunus caroliniana) laural, mountain						(Leschampsia caespitosa) verbena, shrub					x
Controlling poly   Control poly	dianthus (sweet William)	yellow								х			Verbena, St. Paul's (Verbena peruviana.)					х
Communication   Communicatio	(Dianthus spp.)	magic fountain					(Leucothoe fontanesiana) ligustrum, Japanese			х		х		berry bush				х
Comus fonds	dogwood	+												European cran				
	(Cornus florida) dogwood, American	flavarimea						Peter Pan						linden				
Paudoting memberial	(Comus sericea) Douglas fir	-					(Liriope muscari)	giant Blacksouty	¥	x	¥	x	vinca (periwinkle)	wright				X X
yorasu-curiomissa yorasu-curio	(Pseudotsuga menziesii)		x		x			majestic monroe white		x x		x	(Vinca minor) windmill palm		-			x
Climas pare/folia		drako			<u> </u>			silvery sun proof		x		x	(Xvlosma congestum)					×
Mescuritus sinerals sinerals (fines controls)  Higher, and sinerals X X X  Person controls (fines controls)	(Senecio cineraria) elm							runzyaid		х		х	yarrow			х		х
	(Senecio cineraria) elm (Ulmus parvifolia)		x	x	x	x	(Liriope spicata)						Validou abby)	dwarf		-		
	(Senecio cineraria) elm (Ulmus parvifolia) eulaliagrass		x	x	x	х		emerald elf	х		x		(Nex vomitoria) yew	-		×		_ x