

# Specimen Label



Dow AgroSciences



## SPECIALTY HERBICIDE

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

- For control of annual and perennial broadleaf weeds including invasive and noxious weeds, certain annual grasses, and certain woody plants and vines, on:
  - rangeland, permanent grass pastures (including grasses grown for hay\*), Conservation Reserve Program (CRP)
  - non-crop areas for example, airports, barrow ditches, communication transmission lines, electric power and utility rights-of-way, fencerows, gravel pits, industrial sites, military sites, mining and drilling areas, oil and gas pads, non-irrigation ditch banks, parking lots, petroleum tank farms, pipelines, roadsides, railroads, storage areas, dry storm water retention areas, substations, unimproved rough turf grasses; and
  - natural areas (open space) for example, campgrounds, parks, prairie management, trailheads and trails, recreation areas, wildlife openings, and wildlife habitat and management areas;
  - including grazed areas in and around these sites.


\*Hay from grass treated with Milestone within the preceding 18-months can only be used on the farm or ranch where the product is applied unless allowed by supplemental labeling

It is permissible to treat non-irrigation ditch banks, seasonally dry wetlands (such as flood plains, deltas, marshes, swamps, or bogs) and transitional areas between upland and lowland sites only when dry. Milestone can be used to the waters edge. Do not apply directly to water and take precautions to minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize overspray to open water. Note: Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

### IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- Carefully read the section "**Restrictions in Hay or Manure Use .**"
- It is mandatory to follow the "**Use Precautions and Restrictions**" section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied unless allowed by supplemental labeling.
- Consult with a Dow AgroSciences representative if you do not understand the "Use Precautions and Restrictions". Call [1-(800) 263-1196] Customer Information Group.

### Forage and Manure Management



©Copyright 2011 Dow AgroSciences LLC

Not For Sale, Distribution, or Use in New York State.




|       |   |           |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

Active Ingredient:

|   |        |
|---|--------|
| Triisopropanolammonium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro-..... | 40.6%  |
| Other Ingredients .....   | 59.4%  |
| Total .....   | 100.0% |

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 21.1% - 2 lb/gal

**Container Use Directions**

|   |  |   |
|---|--|---|
| <p><b>1 - Tip</b></p>  <p>Tilt container to angle as shown and fill head to desired amount – use vertical scale for measuring. Container should be closed.</p> | <p><b>2 - Level</b></p>  <p>Hold container up-right and check the amount for accuracy. Add or subtract as needed, using pour-back scale as guide.</p> | <p><b>3 - Dispense</b></p>  <p>Remove cap on head and pour into sprayer or other devices. No fluid will pour from the main container. Replace cap for storage in sealed condition.</p> |
|---|--|---|

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**  
EPA Reg. No. 62719-519 EPA Est. \_\_\_\_\_

**CAUTION**

**Causes Moderate Eye Irritation**

**Avoid contact with eyes or clothing.**

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


Read all Directions for Use carefully before applying.

**This product is not intended for reformulation or repackaging into other end-use products.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Not For Sale, Distribution, or Use in New York State.**

**Not for use on pastures in Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont. All other labeled uses are permitted in these states including grazed areas in and around these sites.**



Light grey = states where use in pastures is not permitted  
Dark grey = NY where the product is not registered

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material as polyethylene or polyvinyl chloride
- Shoes plus socks
- Protective eyewear

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS does not pertain to non-agricultural use on sites, such as, rangeland, permanent grass pastures, or non-cropland. See the Agricultural Use Requirements section below for information where the WPS applies.

**Entry Restrictions for Non-WPS Uses:** For applications on rangeland and permanent grass pastures (not harvested for hay) and non-cropland areas, do not enter or allow worker entry into treated areas until sprays have dried.

**Storage and Disposal**

Do not contaminate water, food, feed or fertilizer by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** If this product is exposed to subfreezing temperatures, the active ingredient may crystallize and settle out of solution. Under these conditions the product should be warmed to at least 40°F and agitated well to dissolve any crystallized active ingredient prior to use.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Nonrefillable containers 5 gallons or less:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down

## Storage and Disposal (Cont.)

over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### Refillable containers larger than 5 gallons:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### Nonrefillable containers larger than 5 gallons:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## Resistance Management Guidelines

- Development of plant populations resistant to this herbicide mode of action is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications.
- In croplands, use an effective integrated pest management (IPM) program, integrating tillage or other mechanical methods, crop rotation or other cultural control methods into weed control programs whenever practical.
- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to a herbicide. Application of a herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where identified, spreading of resistant weeds to other fields may be prevented by cleaning harvesting and tillage equipment before moving to other areas and by planting weed-free seed.
- Contact your extension specialist, certified crop consultant, or Dow AgroSciences representative for the latest resistance management information.

## Use Precautions and Restrictions

**Consult with a Dow AgroSciences representative if you do not understand the "Use Precautions and Restrictions." Call (1-800-263-1196) for more information.**

### Pasture and Rangeland Restrictions

- **Do not use grasses treated with Milestone in the preceding 18-months for hay intended for export outside the United States.**
- **Hay from areas treated with Milestone in the preceding 18-months CAN NOT be distributed or made available for sale off the farm or ranch where harvested unless allowed by supplemental labeling.**

- **Hay from areas treated with Milestone in the preceding 18-months CAN NOT be used for silage, haylage, baylage and green chop unless allowed by supplemental labeling.**
- **Do not move hay made from grass treated with Milestone within the preceding 18-months off farm unless allowed by supplemental labeling.**
- **Do not use hay or straw from areas treated with Milestone within the preceding 18-months or manure from animals feeding on hay treated with Milestone in compost.**
- **Do not use grasses treated with Milestone in the preceding 18-months for seed production.**

**Maximum Application Rate:** On all labeled use sites do not broadcast apply more than 7 fl oz per acre of Milestone per year. The total amount of Milestone applied broadcast, as a re-treatment, and/or spot treatment cannot exceed 7 fl oz per acre per year. Spot treatments may be applied at an equivalent broadcast rate of up to 0.22 lb acid equivalent (14 fl oz of Milestone) per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 0.11 lb acid equivalent (7 fl oz) per acre of Milestone per annual growing season as a result of broadcast, spot or repeat applications.

Obtain Required Permits: Consult with appropriate state or local water authorities before applying this product around public waters. State or local public agencies may require permits.

- **Avoiding Injury to Non-Target Plants:** Do not aerially apply Milestone within 50 feet of a border downwind (in the direction of wind movement), or allow spray drift to come in contact with, any broadleaf crop or other desirable broadleaf plants, including, but not limited to, alfalfa, cotton, dry beans, flowers, grapes, lettuce, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes or other broadleaf or vegetable crop, fruit trees, ornamental plants, or soil where sensitive crops are growing or will be planted. Avoid application under conditions that may allow spray drift because very small quantities of spray may seriously injure susceptible crops. Read and consider the "Precautions for Avoiding Spray Drift and Spray Drift Advisory" at the end of this label to help minimize the potential for spray drift.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Do not contaminate water intended for irrigation or domestic purposes.** Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not apply this product to lawns, turf, ornamental plantings, urban walkways, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
- Trees adjacent to or in a treated area can occasionally be affected by root uptake of Milestone. Do not apply Milestone within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.
- Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of Milestone. Injury to crops may result if treated soil and/or runoff water containing Milestone is washed, or moved onto land used to produce crops. Exposure to Milestone may injure or kill susceptible crops and other plants, such as grapes, soybeans, tobacco, sensitive ornamentals. Do not treat frozen soil where runoff could damage sensitive plants.
- **Grass revegetation:**
  - Milestone can be used to control broadleaf plants in grass revegetation programs. Consult Dow AgroSciences' literature for more details about Milestone applications and grass stand establishment.
- **Application before seeding grasses**
  - Milestone can be applied to control broadleaf weeds prior to grass planting. Grass seed germination and seedling development can be adversely affected by many factors such as seed viability and seedling vigor, soil condition (sub-optimal soil temperatures or soil water content), weather after planting, seedbed preparation and seed placement, disease, insects, or animals. Milestone applications will help to reduce competition from weeds and improve the chance for successful grass stand establishment. Some grass species are more sensitive to Milestone; consult Dow AgroSciences' literature for more details.



- **Postemergence applications on grass:** During the season of establishment, Milestone should be applied only after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses are tolerant to Milestone at this stage of development. Milestone may suppress certain established grasses, such as smooth brome grass (*Bromus inermis*), especially when plants are stressed by adverse environmental conditions. Plants should recover from this transient suppression with the onset of environmental conditions favorable to grass growth and upon release from weed competition.
- **Seeding Broadleaf Plants (Forbs) and Wildflowers**
  - Milestone can be applied in the summer to control broadleaf weeds prior to forb planting. Forbs can be seeded 90 days after a summer application as a dormant fall planting or the following spring. Consult Dow AgroSciences literature for details.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or grass hay harvest following application of Milestone at labeled rates. Cutting hay too soon after spraying weeds will reduce weed control. Wait 14 days after herbicide application to cut grass hay to allow herbicide to work. Do not transfer grazing animals from areas treated with Milestone to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.
- **Restrictions in Hay or Manure Use:**
  - ◆ Do not use aminopyralid-treated plant residues, including grass, wood plants, trees, hay or straw from areas treated within the preceding 18-months, in compost, mulch wood chips, or mushroom spawn.
  - ◆ Do not use manure from animals that have eaten aminopyralid-treated forage or hay within the previous 3 days in compost, mulch or mushroom spawn. Livestock must have 3 days of eating non-aminopyralid-treated materials in order to clear their system of aminopyralid. Do not use aminopyralid-treated plants in areas where commercially grown mushrooms or susceptible broadleaf plants may be grown.
  - ◆ Do not spread manure from animals that have consumed aminopyralid-treated forage or hay within the previous 3 days on land used for growing susceptible broadleaf crops.
  - ◆ Manure from animals that have consumed aminopyralid-treated forage or hay within the previous 3 days may only be used on areas used for pasture, grass grown for seed, wheat and corn.
  - ◆ Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields or areas treated with aminopyralid or manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - ◆ Do not plant a broadleaf crop in fields or areas treated in the previous year with manure from animals that have consumed aminopyralid-treated forage or hay until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - ◆ To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- **Crop Rotation:** Do not rotate to any crop from rangeland, permanent pasture or CRP acres within one year following treatment. Cereals and corn can be planted one year after treatment. Broadleaf crops are sensitive to aminopyralid residues in the soil and prediction of crop safety by field bioassay (see instructions below) is the BEST way to determine planting options. Broadleaf crops such as canola, flax, and alfalfa can require **at least 2 to 3 years** depending on the crop and environmental conditions. More sensitive crops such as soybeans, tobacco, peanuts, potatoes, and peas may require a longer plant back interval and should not be planted until a field bioassay shows that the level of aminopyralid present in the soil will not adversely affect that broadleaf crop.

**Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern or drainage. The field bioassay can be initiated one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed

germination), chlorosis (yellowing), epinasty, and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, forage grasses, native grasses or grasses grown for hay.

#### Sprayer Clean-Out Instructions

**It is recommended to use separate spray equipment on highly sensitive crops such as tobacco, soybeans, potatoes, peanuts and tomatoes.**

Do not use spray equipment used to apply Milestone for other applications to land planted to, or to be planted to, broadleaf plants unless it has been determined that all residues of this herbicide have been removed by thorough cleaning of equipment.

Equipment used to apply Milestone should be thoroughly cleaned before reusing to apply any other chemicals as follows:

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
  2. Rinse a second time, adding 1 quart of household ammonia or tank cleaning agent for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
  3. Flush the solution out of the spray tank through the boom.
  4. Rinse the system twice with clean water, recirculating and draining each time.
  5. Spray nozzles and screens should be removed and cleaned separately.
- Do not apply this product with mist blower systems that deliver very fine spray droplets. Use of mist blower equipment can reduce control achieved with the herbicide and increase spray drift potential.

#### General Use Information

Apply the specified rate of Milestone as a coarse low-pressure spray. Do not apply this product with mist blower systems that deliver very fine spray droplets. Spray volume should be sufficient to uniformly cover foliage or intended application site. Increase spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. To enhance foliage wetting and coverage, a non-ionic agricultural surfactant or other adjuvant may be added to the spray mixture as specified by the adjuvant label.

Milestone may be applied by ground or aerial application equipment on any registered use site specified on this label.

**Ground Broadcast Application:** Higher spray volumes (greater than 10 gallons per acre) generally provide better coverage and better control, particularly in dense and/or tall foliage.

**Aerial Broadcast Application:** Do not apply less than 2 gallons per acre total spray volume. Five gallons per acre or greater will generally provide better coverage and better control, particularly in dense and/or tall foliage.

**High-Volume Foliar Application:** High volume foliar treatments may be applied at rates equivalent to a maximum of 7 fl oz per acre per annual growing season. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems.

For basal bark and cut stubble and all types of cut surface applications, see woody plant section.

#### Low-Volume Foliar Treatment

To control susceptible woody plants, use Milestone alone or in tank mixes with other herbicides in water. The spray concentration of Milestone tank mixes and total spray volume per acre should be adjusted according to the size and density of target woody plants and type of spray equipment used. With low-volume application, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems, and root collars.

For best results, an adjuvant should be added to all spray mixtures. Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 psi may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

**Spot Application:** Spot treatments may be applied at an equivalent broadcast rate of up to 0.22 lb acid equivalent (14 fl oz of Milestone) per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 0.11 lb acid equivalent (7 fl oz) per acre of Milestone per annual growing season as a result of broadcast, spot or repeat applications.) Spray volume should be sufficient to thoroughly and uniformly wet weed foliage, but not to the point of runoff. Repeat treatments may be made, but the total amount of Milestone applied must not exceed 7 fl oz per acre per year. To prevent

misapplication, spot treatments should be applied with a calibrated sprayer with a known volume per acre. Table 1 shows Milestone amount to mix for various sprayer outputs in gallons per acre (GPA)

**Table 1:** Amount of Milestone (in mL) to mix in 1 gallon of water

| Gallons per acre | Milestone amount (in mL) to mix to achieve target application rates |           |           |
|------------------|---|-----------|-----------|
|                  | GPA   | 5 fl oz/a | 7 fl oz/a |
| 20               | 7.5   | 10.5      | 21.0      |
| 30               | 5.0   | 7.0       | 14.0      |
| 40               | 3.8   | 5.3       | 10.5      |
| 50               | 3.0   | 4.2       | 8.4       |
| 60               | 2.5   | 3.5       | 7.0       |
| 70               | 2.1   | 3.0       | 6.0       |
| 80               | 1.9   | 2.6       | 5.3       |
| 90               | 1.7   | 2.3       | 4.7       |
| 100              | 1.5   | 2.1       | 4.2       |

Use a syringe to measure cc

Note: Table 1 above shows mixes for various sprayer outputs in gallons per acre (GPA).

**Conversions:**

1 tsp = 5 mL                      30 ml = 1 fluid ounce                      1 cc = 1 mL  
 3 tsp = 1 Tbsp                      2 Tbsp = 1 fluid ounce

**Mixing Instructions**

**Mixing with Water:** To prepare the spray, add about half the required amount of water in the spray tank. Then, with agitation, add the specified amount of Milestone and other herbicides, if tank mixing. Finally, with continued agitation, add the rest of the water and additives such as adjuvants, surfactants or drift control and deposition aids.

**Addition of Surfactants or Adjuvants on All Labeled Use Sites:** The addition of a high quality non-ionic surfactant (of at least 80% active principal) or adjuvant at 0.25 to 0.5 % volume per volume (1 to 2 quarts per 100 gallons of spray) is recommended to enhance herbicide activity under adverse environmental conditions (such as, high temperature, low relative humidity, drought conditions, dusty plant surfaces) or when weeds are heavily pubescent or more mature.

**Tank Mixing with Other Herbicides:** Milestone may be applied in tank mix combination with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the tank mix product(s), and (3) that the tank mix combination is physically compatible (see tank mix compatibility testing below). When tank mixing, use only in accordance with the restrictions, precautions and limitations on the respective product labels.

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed specified application rates. If products containing the same active ingredient are mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** Perform a jar test prior to mixing in a spray tank to ensure compatibility of Milestone and other pesticides or carriers. Use a clear glass jar with lid and mix ingredients in the same order and proportions as will be used in the spray tank. The mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily remix if agitated. An incompatible mixture is indicated by separation into distinct layers that do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film in the jar. Use of an appropriate compatibility aid may resolve mix incompatibility. If the mixture is incompatible do not use that tank mix partner in tank mixtures.

**Invert emulsion spray mixtures**

Milestone can be applied in an invert emulsion using oil and an appropriate inverting agent. Follow label directions of the inverting agent.

**Mixing with Sprayable Liquid Fertilizer Solutions:** Milestone is usually compatible with liquid fertilizer solutions. It is anticipated that Milestone will not require a compatibility agent for mixing with fertilizers; however, a compatibility test (jar test) should be made prior to mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank.

**Note:** The lower the temperature of the liquid fertilizer, the greater the likelihood of mixing problems. Use of a compatibility aid may be required if Milestone is mixed with a 2,4-D-containing product and liquid fertilizer. **Mixing Milestone and 2,4-D in N-P or N-P-K liquid fertilizer solutions is more difficult than mixing with straight nitrogen fertilizer and should not be attempted without first conducting a successful compatibility jar test.** Agitation in the spray tank must be vigorous to be comparable with jar test agitation. Apply the spray mixture the same day it is prepared while maintaining continuous agitation. Rinse the spray tank thoroughly after use.

**Note:** Foliar-applied liquid fertilizers themselves can cause yellowing of the foliage of forage grasses and other vegetation.

**Use Rates and Timing**

Milestone may be applied as a broadcast spray by ground or aerial equipment or as a spot application to control weeds including, but not limited to, those listed on this label. When a rate range is given use the higher rate to control weeds at advanced growth stages, or under less than favorable growing conditions, or for longer residual control. Best results are obtained when spray volume is sufficient to provide uniform coverage of treated weeds. For optimum uptake and translocation of Milestone, avoid mowing, haying, shredding, burning or soil disturbance in treated areas for at least 14 days following application.

Milestone provides post emergence control and premergence control of emerging seedlings of susceptible weeds, and re-growth of certain perennial weeds following application. Preventing establishment of weeds will depend upon application rate, season of application, and environmental conditions after application.

Milestone can provide long-term control of susceptible weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Long-term weed control is most effective where grass vegetation is allowed to recover from overgrazing, drought, etc., and compete with weeds.

Milestone can be an important component of integrated vegetation management programs designed to renovate or restore desired plant communities. To maximize and extend the benefits of weed control provided by Milestone, it is important that other vegetation management practices, including proper grazing management, biological control agents, replanting, fertilization, prescribed fire, etc., be used in appropriate sequences and combinations to further alleviate the adverse effects of weeds on desirable plant species and to promote development of desired plant communities. Agricultural and natural resources specialists with federal and state government agencies can provide guidance on best management practices and development of integrated vegetation management programs.

**Plants Controlled**

The following weeds and woody plants will be controlled with the rates of Milestone indicated below (table 2). For best results, most weeds and woody plants should be treated when they are actively growing and under conditions favorable for growth. Use a higher rate in the rate range when growing conditions are less than favorable or when weed foliage is tall and dense, or when optimal longer term residual control is desired. Milestone also provides premergence control of germinating seeds or seedlings of susceptible weeds following application.

**Table 2:** Weeds and Woody Plants Controlled**Note:** Numbers in parentheses (-) refer to specific use directions for a particular weeds species.

| Common Name                        | Scientific Name                    | Rate Range (fl oz/acre) | Life Cycle             | Plant Family     |
|------------------------------------|------------------------------------|-------------------------|------------------------|------------------|
| amaranth, spiny                    | <i>Amaranthus spinosus</i>         | 4 to 7                  | annual                 | Amaranthaceae    |
| bedstraw                           | <i>Galium spp.</i>                 | 4 to 7                  | perennial              | Rubiaceae        |
| beggarticks                        | <i>Bidens spp.</i>                 | 4 to 7                  | annual                 | Asteraceae       |
| broomweed, annual                  | <i>Amphiachyris dracunculoides</i> | 4 to 7                  | annual                 | Asteraceae       |
| burdock, common                    | <i>Arctium minus</i>               | 4 to 7                  | biennial               | Asteraceae       |
| buttercup, hairy                   | <i>Ranunculus sardous</i>          | 4 to 7                  | annual                 | Ranunculaceae    |
| buttercup, tall                    | <i>Ranunculus acris</i>            | 4 to 7                  | perennial              | Ranunculaceae    |
| buttercup spp                      | <i>Ranunculus spp</i>              | 4 to 7                  | various                | Ranunculaceae    |
| camelthorn                         | <i>Alhagi pseudalhagi</i>          | 5 to 7                  | perennial              | Fabaceae         |
| cat's ear, common                  | <i>Hypochaeris radicata</i>        | 5 to 7                  | perennial              | Asteraceae       |
| cat's ear                          | <i>Hypochaeris spp</i>             | 5 to 7                  | perennial              | Asteraceae       |
| chamomile, scentless               | <i>Matricaria inodora</i>          | 4 to 7                  | annual                 | Asteraceae       |
| chicory                            | <i>Cichorium intybus</i>           | 4 to 6                  | perennial              | Asteraceae       |
| chickweed                          | <i>Stellaria media</i>             | 7                       | annual                 | Caryophyllaceae  |
| cinquefoil, sulfur (1)             | <i>Potentilla recta</i>            | 4 to 7                  | perennial              | Rosaceae         |
| cocklebur                          | <i>Xanthium strumarium</i>         | 3 to 5                  | annual                 | Asteraceae       |
| clover                             | <i>Trifolium spp.</i>              | 5 to 7                  | perennial              | Fabaceae         |
| crazyweed                          | <i>Oxytropis</i>                   | 5 to 7                  | perennial              | Fabaceae         |
| croton, tropic                     | <i>Croton glandulosus</i>          | 3 to 5                  | annual                 | Euphorbiaceae    |
| crownvetch                         | <i>Securigera varia</i>            | 5 to 7                  | perennial              | Fabaceae         |
| cudweed, purple                    | <i>Gamochaeta purpurea</i>         | 4 to 7                  | annual                 | Asteraceae       |
| daisy, oxeye (1)                   | <i>Leucanthemum vulgare</i>        | 4 to 7                  | perennial              | Asteraceae       |
| dock, curly                        | <i>Rumex crispus</i>               | 4 to 7                  | perennial              | Polygonaceae     |
| evening primrose, cutleaf          | <i>Oenothera laciniata</i>         | 4 to 7                  | annual                 | Onagraceae       |
| fiddleneck                         | <i>Amsinckia spp</i>               | 4 to 7                  | annual                 | Boraginaceae     |
| fireweed                           | <i>Epilobium angustifolium</i>     | 5 to 7                  | perennial              | Onagraceae       |
| fleabane, flax-leaf                | <i>Conyza bonariensis</i>          | 4 to 7                  | annual                 | Asteraceae       |
| fleabane, hairy                    | <i>Conyza bonariensis</i>          | 5-7                     | annual/biennial        | Asteraceae       |
| hawkweed, orange (2)               | <i>Hieracium aurantiacum</i>       | 4 to 7                  | perennial              | Asteraceae       |
| hawkweed, yellow (2)               | <i>Hieracium caespitosum</i>       | 4 to 7                  | perennial              | Asteraceae       |
| henbane, black                     | <i>Hyoscyamus niger</i>            | 5 to 7                  | annual/biennial        | Solanaceae       |
| henbit                             | <i>Lamium amplexicaule</i>         | 5 to 7                  | annual/<br>biennial    | Lamiaceae        |
| hogweed, giant                     | <i>Heracleum mantegazzianum</i>    | 7                       | perennial              | Apiaceae         |
| horsenettle, Carolina              | <i>Solanum carolinense</i>         | 4 to 7                  | perennial              | Solanaceae       |
| horsetweed (marestail)             | <i>Conyza canadensis</i>           | 4 to 7                  | annual                 | Asteraceae       |
| ironweed, tall                     | <i>Vernonia gigantea</i>           | 5 to 7                  | perennial              | Asteraceae       |
| ironweed, western                  | <i>Vernonia baldwinii</i>          | 7                       | perennial              | Asteraceae       |
| knapweed, diffuse (3)              | <i>Centaurea diffusa</i>           | 5 to 7                  | biennial/<br>perennial | Asteraceae       |
| knapweed, meadow                   | <i>Centaurea debeauxii</i>         | 5 to 7                  | perennial              | Asteraceae       |
| knapweed, Russian (4)              | <i>Acroptilon repens</i>           | 5 to 7                  | perennial              | Asteraceae       |
| knapweed, spotted (3)              | <i>Centaurea stoebe</i>            | 5 to 7                  | biennial/<br>perennial | Asteraceae       |
| knapweed, squarrose                | <i>Centaurea virgata</i>           | 5 to 7                  | biennial/<br>perennial | Asteraceae       |
| knapweeds                          | <i>Centaurea spp.</i>              | 5 to 7                  | biennial/ perennial    | Asteraceae       |
| knotweeds, Japanese, bohemian (11) | <i>Reynoutria japonica</i>         | 7-14                    | perennial              | Polygonaceae     |
| kudzu                              | <i>Pueraria montana</i>            | 7                       | perennial              | Fabaceae         |
| lady's thumb                       | <i>Polygonum persicaria</i>        | 3 to 5                  | annual                 | Polygonaceae     |
| lambquarters                       | <i>Chenopodium album</i>           | 5 to 7                  | annual                 | Chenopodiaceae   |
| lespedeza, annual                  | <i>Lespedeza striata</i>           | 5 to 7                  | annual                 | Fabaceae         |
| licorice, wild                     | <i>Glycyrrhiza lepidota</i>        | 7                       | perennial              | Fabaceae         |
| locoweed                           | <i>Astragalus spp.</i>             | 5 to 7                  | perennial              | Fabaceae         |
| locust, black                      | <i>Robinia pseudoacacia</i>        | 7                       | woody perennial        | Fabaceae         |
| locust, honey                      | <i>Gleditsia triacanthos</i>       | 7                       | woody perennial        | Fabaceae         |
| loosestrife, purple (12)           | <i>Lythrum salicaria</i>           | 7-14                    | perennial              | Lythraceae       |
| mayweed, scentless                 | <i>Tripleurospermum perforate</i>  | 4 to 7                  | annual                 | Asteraceae       |
| mayweed, stinking                  | <i>Anthemis cotula</i>             | 7                       | annual                 | Asteraceae       |
| medic, black                       | <i>Medicago lupulina</i>           | 4 to 7                  | perennial              | Fabaceae         |
| mimosa                             | <i>Albizia julibrissin</i>         | 7                       | woody perennial        | Fabaceae         |
| mullein (5)                        | <i>Verbascum spp.</i>              | 7                       | biennial               | Scrophulariaceae |
| nightshade, silverleaf             | <i>Solanum elaeagnifolium</i>      | 4-7                     | perennial              | Solanaceae       |
| oxtongue, bristly                  | <i>Picris echioides</i>            | 5 to 7                  | biennial               | Asteraceae       |



**Table 2:** Weeds and Woody Plants Controlled (Cont.)**Note:** Numbers in parentheses (-) refer to specific use directions for a particular weeds species.

| Common Name                     | Scientific Name                 | Rate Range (fl oz/acre) | Life Cycle      | Plant Family   |
|---------------------------------|---------------------------------|-------------------------|-----------------|----------------|
| pea, Swainson                   | <i>Sphaerophysa salsula</i>     | 5-7                     | perennial       | Fabaceae       |
| povertyweed                     | <i>Iva axillaris</i>            | 5-7                     | perennial       | Asteraceae     |
| ragweed, common                 | <i>Ambrosia artemisiifolia</i>  | 3 to 5                  | annual          | Asteraceae     |
| ragweed, western                | <i>Ambrosia psilostachya</i>    | 4 to 7                  | perennial       | Asteraceae     |
| ragweed, giant                  | <i>Ambrosia trifida</i>         | 4 to 7                  | annual          | Asteraceae     |
| ragwort, tansy                  | <i>Senecio jacobaea</i>         | 5 to 7                  | perennial       | Asteraceae     |
| redbud                          | <i>Cercis Canadensis</i>        | 7                       | woody perennial | Fabaceae       |
| rush skeletonweed               | <i>Chondrilla juncea</i>        | 5 to 7                  | perennial       | Asteraceae     |
| sicklepod                       | <i>Cassia obtusifolia</i>       | 7                       | perennial       | Fabaceae       |
| smartweed, Pennsylvania         | <i>Polygonum pennsylvanicum</i> | 3 to 5                  | annual          | Polygonaceae   |
| sneezeweed, bitter              | <i>Helenium amarum</i>          | 4 to 7                  | annual          | Asteraceae     |
| soda apple, tropical (6)        | <i>Solanum viarum</i>           | 5 to 7                  | perennial       | Solanaceae     |
| sowthistle, annual              | <i>Sonchus oleraceae</i>        | 7                       | annual          | Asteraceae     |
| sowthistle, perennial           | <i>Sonchus arvensis</i>         | 3 to 5                  | perennial       | Asteraceae     |
| spanishneedles                  | <i>Bidens bipinnata</i>         | 4 to 7                  | annual          | Asteraceae     |
| St. Johnswort, common           | <i>Hypericum perforatum</i>     | 5 to 7                  | perennial       | Clusiaceae     |
| stiltgrass, Japanese            | <i>Microstegium vimineum</i>    | 5-7                     | annual          | Poaceae        |
| starthistle, Malta (7)          | <i>Centaurea melitensis</i>     | 3 to 5                  | annual          | Asteraceae     |
| starthistle, purple (7)         | <i>Centaurea calcitrapa</i>     | 3 to 5                  | biennial        | Asteraceae     |
| starthistle, yellow (7)         | <i>Centaurea solstitialis</i>   | 3 to 5                  | annual          | Asteraceae     |
| sunflower, common               | <i>Helianthus annuus</i>        | 4 to 7                  | annual          | Asteraceae     |
| sweetclover, white              | <i>Melilotus albus</i>          | 5 to 7                  | biennial        | Fabaceae       |
| sweetclover, yellow             | <i>Melilotus officinalis</i>    | 5 to 7                  | biennial        | Fabaceae       |
| teasel                          | <i>Dipsacus spp.</i>            | 4 to 7                  | biennial        | Dipsacaceae    |
| thistle, artichoke              | <i>Cynara cardunculus</i>       | 5 to 7                  | perennial       | Asteraceae     |
| thistle, blessed milk           | <i>Silybum marianum</i>         | 4-7                     | biennial        | Asteraceae     |
| thistle, bull (8)               | <i>Cirsium vulgare</i>          | 3 to 5                  | biennial        | Asteraceae     |
| thistle, Canada (9)             | <i>Cirsium arvense</i>          | 5 to 7                  | perennial       | Asteraceae     |
| thistle, woolly distaff         | <i>Carthamus lanatus</i>        | 4 to 7                  | annual          | Asteraceae     |
| thistle, Italian                | <i>Carduus pycnocephalus</i>    | 7                       | annual          | Asteraceae     |
| thistle, musk (8)               | <i>Carduus nutans</i>           | 3 to 5                  | biennial        | Asteraceae     |
| thistle, plumeless (8)          | <i>Carduus acanthoides</i>      | 3 to 5                  | biennial        | Asteraceae     |
| thistle, Scotch                 | <i>Onopordum acanthium</i>      | 5 to 7                  | biennial        | Asteraceae     |
| thistle, Russian (preemergence) | <i>Salsola spp</i>              | 7                       | annual          | Chenopodiaceae |
| tree of heaven                  | <i>Ailanthus altissima</i>      | 7                       | perennial       | Simaroubaceae  |
| vetch                           | <i>Vicia spp.</i>               | 3 to 7                  | perennial       | Fabaceae       |
| willoweed, panicle              | <i>Epilobium brachycarpum</i>   | 5-7                     | annual          | Onagraceae     |
| wisteria                        | <i>Wisteria brachybotris</i>    | 7                       | woody perennial | Fabaceae       |
| wormwood, absinth(10)           | <i>Artemisia absinthium</i>     | 6 to 7                  | perennial       | Asteraceae     |
| yarrow, common                  | <i>Achillea millefolium</i>     | 7                       | perennial       | Asteraceae     |

- (1) **Sulfur cinquefoil or oxeye daisy:** Apply Milestone at 4 to 6 fl oz per acre to plants in the prebud stage of development.
- (2) **Orange or yellow hawkweeds:** Apply Milestone at 4 to 7 fl oz per acre to plants in the bolting stage of development.
- (3) **Diffuse, spotted, and squarrose knapweeds:** Apply Milestone at 5 to 7 fl oz per acre when plants are actively growing with the optimum time of application occurring from rosette to the bolting stages of development or in the fall. Plants will be controlled by mid-summer and fall applications even though plants may not show any changes in form or stature the year of application.
- (4) **Russian knapweed:** Apply Milestone at 5 to 7 fl oz per acre to plants in the spring and summer at early bud to flowering stages and to dormant plants in the fall.
- (5) **Mullein:** Apply to the rosette stage
- (6) **Tropical soda apple:** Apply Milestone at 5 to 7 fl oz per acre at any growth stage, but application by flowering will reduce seed production potential.
- (7) **Malta, purple, and yellow starthistle:** Apply Milestone at 3 to 5 fl oz per acre to plants at the rosette through bolting growth stages.
- (8) **Bull, musk, and plumeless thistles:** Apply Milestone at 3 to 5 fl oz per acre in the spring and early summer to rosette or bolting plants or in the fall to seedlings and rosettes. Apply at 4 to 5 fl oz when plants are at the late bolt through early flowering growth stages. 2,4-D at 1 lb ae/acre should be tank-mixed with Milestone starting at the late bud stages
- (9) **Canada thistle:** Apply Milestone at 5 to 7 fl oz per acre in the spring after all plants have fully emerged (some may be budding) until the oldest plants are in full flower stage. Use the higher rate when applying to the flower stage. Applications are also effective in the fall before a killing frost. Use higher rates for older/dense stands or for longer residual control.
- (10) **Absinth wormwood:** Apply 6 to 7 fl oz per acre before wormwood is 12 inches tall. When applying by air on CRP, coverage is important and a minimum of 3 GPA is specified. Remove old duff and litter by fire or mowing for best results
- (11) **Invasive knotweeds:** Japanese, Bohemian, giant knotweeds: Optimum suppression of invasive knotweeds with Milestone herbicide is obtained when applications are made to plants that are at least 3 to 4 feet tall. Results of field trials conducted in the western U.S. indicate that high volume applications (100 gpa or greater) of Milestone at 7 fl oz/A or a spot treatment rate up to 14 fl oz/A applied in summer will provide good control of invasive knotweeds. In the upper Midwest, mowing in summer followed by fall application of Milestone (prior to frost) provided the best control. Infestations of invasive knotweed that are mowed should be allowed to regrow to at least 3 feet in height prior to herbicide treatment. Monitoring and follow-up herbicide treatments on regrowth will be necessary to control resprouts and achieve long-term control.
- (12) **Purple loostrate:** For optimum control apply Milestone at 7 fl oz per acre plus 1 pt to 1 qt of 2,4-D amine or 1 to 2 qts of Garlon 3A. Spot treatments may also be made by applying Milestone at 14 fl oz (see Spot treatment section of the label) with or without the addition of 2,4-D or Garlon 3A.
- (13) **Fiddleneck:** For optimum control apply Milestone at 4 to 7 fl oz per acre when the plants are young and before flowering. Use higher rates if the plants are older and larger. In California optimal application timing is November through March.

## For Control or Suppression of Medusahead Rye

Milestone applied broadcast at 7 to 14 fl oz/A can suppress or control medusahead rye (*Taeniatherum caput-medusae*) and downy brome (*Bromus tectorum*, also called cheatgrass). The key to optimum results is the timing of application. Applications should be made in late summer prior to rains and seed germination in order to provide the best possibility of suppression or control. In general, control or suppression will be poor if any of the seeds have germinated prior to application even if they have not yet emerged through the soil surface. Tank mixes with Accord XRT II at 12 fl oz/A, where a non-selective herbicide can be used or where desired grasses are dormant and will not be harmed, and will aid in control. Spot treatment restrictions (see spot treatment section) apply for rates above 7 fl oz/A for broadcast applications.

## Woody Plant Control

Milestone may be applied to control woody plants by any application method listed on the label on any site listed.

Milestone may be applied alone or in tank-mix combinations with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products. Use as directed in the Directions of Use section of the tank-mix partner. Follow Mixing Instructions under the General Mixing and Application Instructions section.

Add Milestone to tank mixes for improved brush control on species such as alder, aspen, blackberry, boxelder, cherry, coyote brush, conifers, cottonwood, elm, maple, poplar, oak, brooms (Scotch, Spanish, French, Portuguese), gorse, hackberry, Russian and Autumn olive, salt-cedar.

### Low or High Volume Foliar Applications:

For broad spectrum brush control using a foliar application, Milestone may be added to tank mixes with Accord® XRT II, Arsenal Powerline, DMA® 4IVM, Garlon 4 Ultra, Remedy Ultra, Tordon 101M, Transline, Forestry Garlon XRT, or Garlon 3A, Rodeo®, Tordon® K, Tordon 22K or other products labeled for use on the intended site.

### Low Volume Basal Bark Applications:

To control susceptible woody plants with stems less than 6 inches in basal diameter, apply herbicide mix (see below for rates) with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Spray the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground in a manner that thoroughly wets the lower stems but not to the point of runoff. The use of a Spraying Systems Y2 nozzle or similar nozzle is recommended, which will narrow the spray pattern to target individual stems. Herbicide concentration should vary with tree diameter, bark thickness, volume used per acre, and susceptibility of species treated. Apply anytime, including the winter months, except when snow or water prevent spraying to the ground line or when stem surfaces are saturated with water.

Milestone may be used as a low volume basal treatment alone, for sensitive woody species in the Fabaceae family (legumes), or in combination with other products such as Garlon 4 Ultra, Forestry Garlon XRT, Remedy Ultra for broader control of other sensitive woody species. Applications should not exceed the maximum use rate per acre for the site.

Mix Milestone at 0.5 to 5% v/v alone, or with Garlon 4 Ultra or Forestry Garlon XRT in a commercially available basal diluent (or other oils or basal diluents as recommended by the manufacturer); the basal oil should be compatible with a water soluble herbicide such as Milestone. See table 3 to calculate the amount of Milestone that can be applied per acre at the various volumes and rates. Make a stable tank mixture for basal bark application by first combining each product with a compatibility agent prior to final mixing in the desired ratio. If using a tank mix, mix the oil-based products such as Garlon 4 Ultra thoroughly with basal oil and add any other oil-based products before adding the water based products. If the mixture stands for more than 30 minutes, reagitation may be required.

Oil and water based mixtures can separate over time. Long-term storage is not recommended without vigorous agitation prior to use or without a recommended compatibility agent.

Use caution when treating areas adjacent to susceptible and desirable species to avoid root uptake and possible injury when using Milestone or other soil active herbicides



### Low Volume Stem Bark Band Treatment

To control susceptible woody plants (see table 2) with stems less than 6 inches in basal diameter, mix 0.5 to 5 gallons of Milestone in enough oil to make 100 gallons of spray mixture. Apply with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Apply the spray in a 6- to 10-inch wide band that completely encircles the stem. Spray in a manner that completely wets the bark, but not to the point of runoff. The treatment band may be positioned at any height up to the first major branch. For best results apply the band as low as possible.

Spray mixture concentration should vary with size and susceptibility of species to be treated. Applications may be made anytime, including winter months.

**Table 3:**

| % of Milestone in Basal Mix | Fluid ounces of Milestone by GPA (gallons per acre) |      |      |      |      |      |      |
|-----------------------------|---|------|------|------|------|------|------|
|                             | (1)   | (2)  | (3)  | (4)  | (5)  | (6)  | (7)  |
| 1.0                         | 1.3   | 2.6  | 3.8  | 5.1  | 6.4  | 7.7  | 9.0  |
| 1.5                         | 1.9   | 3.8  | 5.8  | 7.7  | 9.6  | 11.5 | 13.4 |
| 2.0                         | 2.6   | 5.1  | 7.7  | 10.2 | 12.8 |      |      |
| 2.5                         | 3.2   | 6.4  | 9.6  | 12.8 |      |      |      |
| 3.0                         | 3.8   | 7.7  | 11.5 |      |      |      |      |
| 3.5                         | 4.5   | 9.0  | 13.4 |      |      |      |      |
| 4.0                         | 5.1   | 10.2 |      |      |      |      |      |
| 5.0                         | 6.4   | 12.8 |      |      |      |      |      |

 within spot treatment labeled rate  
 in excess of spot treatment labeled rate

NOTE: Avoid treating high density of stems adjacent to desirable trees with roots in the treatment zone. See table 4 for guidance on estimated volume per acre by treated stem density. Trees adjacent to or in a treated area can occasionally be affected by root uptake of Milestone. Applications of Milestone within the root zone of desirable trees should not be made unless injury can be tolerated. Severe injury or plant death can occur if used near roses, or leguminous trees such as locusts, redbud, mimosa, and caragana.

**Table 4:**

| Estimated gallons of spray solution per acre for basal bark applications on various stem densities per acre |              |                          |
|---|--------------|--------------------------|
| Number of Stems/Acre  | Volume Range | Target Spacing           |
|   | (gal/acre)   | (ft between brush/trees) |
| 250   | 1.0 - 1.7    | 8.4                      |
| 500   | 2.0 - 3.3    | 5.9                      |
| 750   | 3.0 - 5.0    | 4.9                      |
| 1000  | 4.0 - 6.6    | 4.2                      |
| 1250  | 5.0 - 8.3    | 3.8                      |
| 1500  | 5.9 - 9.9    | 3.4                      |

## Cut surface

Apply Milestone in the cut surface applications listed below for control of susceptible tree species such as legumes like Albezia, mimosa, locust, etc. Mixtures of Milestone and Garlon 3A or Garlon 4 Ultra may be effective on species other than legumes such as elm, maple, oak and conifers.

Cut surface applications may be used successfully at any season except during periods of heavy sap flow of certain species - for example, maples in the spring.

### Cut-Stump Treatment

Apply Milestone as a 10% dilution v/v in water, by spraying or painting all the exposed cambium layer on the freshly cut surface. The cambium area next to the bark is the most vital area to wet.

### With Tree Injector Method

Apply by injecting 1 milliliter of 10% v/v Milestone in water through the bark at intervals of 3 to 4 inches between centers of the injector wound. The injections should completely surround the tree at any convenient height. Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is injected directly into plants.

### With Hack and Squirt Method

Make cuts around the tree trunk at a convenient height with a hatchet or similar equipment so that the cuts overlap slightly and make a continuous circle around the trunk. Spray 1 milliliter of 10% v/v Milestone in water into the pocket created between the bark and the inner stem/trunk by each cut.

### With Frill or Girdle Method

Make a single girdle through the bark completely around the tree at a convenient height. The frill should allow for the herbicide to remain next to the inner stem and absorb into the plant. Wet the cut surface with 10% v/v Milestone in water.



#### For use in Hawaii only:

#### Incision Point Application (IPA) also known as Tree Injection or Hack and Squirt

For control of susceptible tree species such as *Albizia*, and other legumes and susceptible tree species, make cuts around the tree trunk at a convenient height with a machete, hatchet or similar equipment so that the cuts are about 6 inches apart between centers. Inject 1/2 to 1 milliliter of undiluted Milestone into the pocket created between the bark and the inner stem/trunk by each cut as soon as possible after cutting. The cambium area next to the bark is the most vital area to wet.

#### Preemergent Weed Control

Typically Milestone is used as a post emergent herbicide but it has preemergent activity on susceptible weeds. Use Milestone as a preemergence spray prior to weed seed germination. Control will depend upon species susceptibility, application timing, and environmental conditions, such as precipitation, following application. When applied at rates lower than 7 fl oz per acre, Milestone can provide short-term control of some susceptible weeds but when applied at 7 fl oz (broadcast) or 14 fl oz (spot treatment), weed control is extended.

Best results for use as a preemergent application for total vegetation control are obtained if Milestone at 7 fl oz per acre is tank mixed with other herbicides to broaden the weed spectrum and to control grasses. If grasses and broadleaf weeds tolerant to Milestone are present at the time of application or will germinate on the site, then tank mixtures with other herbicides, such as Accord® XRT II, Rodeo®, Dimension® 2EW or EC (annual grasses), Oust XP, Esplanade, flumioxazin, diuron, or other herbicides labeled for total vegetation control applications.

#### SPOT TREATMENTS FOR AREAS SUCH AS SUBJECT POLES, SUBSTATIONS, AND OTHER SMALL AREAS

Spot treatments may be applied at an equivalent broadcast rate of up to 0.22 lb acid equivalent (14 fl oz of Milestone) per acre per annual growing season to small spots for clearing around utility subject poles to help prevent fire damage, on small substations and other spot areas. To prevent misapplication, spot treatments should be applied with a calibrated sprayer.

#### Precautions for Avoiding Spray Drift

Avoid application under conditions that may allow spray drift because very small quantities of spray, which may not be visible, may injure susceptible crops. This product should be applied only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, non-target crops and other plants) is minimal (e.g., when wind is blowing away from the sensitive areas). A drift control aid may be added to the spray solution to further reduce the potential for drift. If a drift control aid is used, follow the use directions and precautions on the manufacturer's label. Do not use a thickening agent with Microfoil, Thru-Valve booms, or other spray delivery systems that cannot accommodate thickened spray solutions.

**Ground Equipment:** With ground equipment spray drift can be lessened by keeping the spray boom as low as possible; by applying 10 gallons or more of spray per acre; by keeping the operating spray pressures at the manufacturer's specified minimum pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); and by spraying when the wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to thermal inversions. Direct sprays no higher than the tops of target vegetation and keep spray pressures low enough to provide coarse spray droplets to minimize drift.

**Aerial Application:** Avoid spray drift at the application site. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The boom length must not exceed 75% of the fixed wing span and must be located at least 8 -10 inches below the trailing edge of the fixed wing; the boom length must not exceed 85% of the rotary blade.
2. Nozzles should be pointed backward parallel with the air stream or not pointed downwards more than 45 degrees.

State regulations must be followed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory**. This information is advisory in nature and does not supersede mandatory label requirements.

#### Aerial Drift Reduction Advisory

**Information on Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control.

Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

#### Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that will provide uniform coverage.
- **Nozzle Orientation** - Orient nozzles so that the spray is released parallel to the airstream to produce larger droplets than other orientations. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Boom Length:** The distance of the outer most operating nozzles on the boom must not exceed 75% of wingspan or 85% of rotor diameter.

**Application Height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain such as valleys and ravines can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

---

#### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

#### Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

## Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

## Limitation of Remedies

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

™Harmony, Express and Ally are trademarks of E. I. DuPont de Nemours and Company

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

Label Code: D02-879-006  
Replaces Label: D02-879-005  
LOES Number: 010-02112

EPA accepted 04/15/14

## Revisions:

1. Added additional uses and application sites throughout the label.
  - a. annual and perennial broadleaf
  - b. vines
  - c. non-crop areas for example
  - d. airports
  - e. barrowditches
  - f. fencerows
  - g. gravel pits
  - h. military sites
  - i. mining and drilling areas
  - j. oil and gas pads
  - k. parking lots
  - l. petroleum tank farms
  - m. pipelines
  - n. storage areas
  - o. storm water retention areas
  - p. substations
  - q. unimproved rough turf grasses
  - r. open space for example parks, prairie management
2. Added the word "dry" before "storm water retention areas"
3. Added the phrase "only when dry" after "upland and lowland sites"
4. Updated the statement "The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the intended rotational crop" to "The field bioassay can be initiated one year after the last application of aminopyralid in that field" throughout the label and supplemental label.

5. Updated and renumbered Tables.
6. Changed Rotational Interval (Months) from 3 to 4 for barley, grasses, field corn, grain sorghum, millet, oats, rye, triticale, sweet corn under Crop Rotation Intervals tables.
7. Changed "Do not rotate to unlisted crops prior to...from 18 to 24 months ...following application without a field bioassay."
8. Added the statement "**This product is not intended for reformulation or repackaging into other end-use products**" to the **Direction for Use** section.
9. Added and update the "**Seeding Broadleaf Plants (Forbs) and Wildflowers**" section under **Pasture and Rangeland Restrictions** section.
10. Removed the "**Seeding Legumes**" under **Pasture and Rangeland Restrictions** section.
11. Removed the "**Forbs and wildflowers in restoration programs**" section and language under **Pasture and Rangeland Restrictions** section. Language moved to new "**Seeding Broadleaf Plants (Forbs) and Wildflowers**" section.
12. Added "aminopyralid" or "aminopyralid-treated" throughout the "**Restrictions in Hay or Manure Use**" section.
13. Added "epinasty" to **Field Bioassay Instructions** section.
14. Added "potatoes" to **Sprayer Clean-Out Instructions** section.
15. Changed "**Application Methods**" to "**General Use Information**"
16. Added "adjuvant" throughout **General Use Information** section and other parts of label.
17. Added the statement "Milestone may be applied by ground or aerial application equipment on any registered use site specified on this label" under **General Use Information** section.
18. Added the statement "For basal bark and cut stubble and all types of cut surface applications, see woody plant section" and "**Low Volume Foliar Treatment**" section.
19. Deleted the statement "Milestone at rate of up to 7 fl oz per acre may be mixed with labeled rates of other herbicide..." from the **Tank Mixing with Other Herbicides** section under **Mixing Instructions** section.
20. Added **Invert emulsion spray mixtures** section under **Mixing Instructions** section.
21. Updated language under **Use Rates and Timing**
22. Added "**For Control or Suppression of Medusahead Rye and Other Winter Annual Grasses**" section.
23. "**Woody Plant Control**" section:
  - a. Added the statement: Milestone may be applied to control woody plants by any application method listed on the label on any site listed.
  - b. Added the following species: alder, coyote brush, cottonwood, brooms (Scotch, Spanish, French, Portuguese), gorse.
  - c. Renamed and updated language under **Low or High Volume Foliar Applications** section.
24. Updated language under **Low Volume Basal Bark Application**
25. Added **Low Volume Stem Bark Band Treatment** section
26. Added "**Chemical Side Trimming**" and "**Cut Stubble Application**" sections.
27. Updated language under **Cut-Stump Treatment**.
28. Added **For use in Hawaii only: Incision Point Application (IPA) also known as Tree Injection or Hack and Squirt** section.
29. Added **Preemergent Weed Control, BROADCAST APPLICATIONS and SPOT TREATMENTS FOR AREAS SUCH AS SUBJECT POLES, SUBSTATIONS, AND OTHER SMALL AREAS** sections.
30. Changed the statement "The distance of the outer most operating nozzles on the boom must not exceed 75% of wingspan or 85% of rotor diameter" to "The boom length must not exceed 75% of the fixed wing span and must be located at least 8 -10 inches below the trailing edge of the fixed wing; the boom length must not exceed 85% of the rotary blade" under **Precautions for Avoiding Spray Drift** section.
31. Updated trademark line.