KRYSTEXXA(pegloticase)	□ New Prescription/Referral□ Prescription Refill# of Refills:
Patient Name:	Patient Weight: DOB:
Rx: Patient Name:	kg
KRYSTEXXA:	<u>-</u>
Administer 8mg via IV infusion for at least 120mins	Q2wks on a 250ml NS at room temperature
Observe for 1 hour after infusion	•
☐ Discontinue treatment if uric acid level to > 6mg/dL	
Other Orders:	
Pre-Medication: (30 mins prior)	☐ Benadrylmg ANA Kit Protocol:
☐ Solumedrol 125mg IVP ☐ Tylenol	, =
☐ Solu-Cortef 100mg IVP ☐ 650mg	
Others:	5 50mg
Gout (M10.9)	
Dx: Others:	
	NPI#:
Physician Signature	Date: (Valid for 1 year)
By affixing my signature. I confirm the medical necessity of	of the aforementioned therapy, products, and services, as well as my
	ent to disclose the mentioned details and any relevant medical or patient
information concerning this treatment. I grant the pharma	acy permission to contact the insurance company on my behalf to secure
authorization for the patient.	
PHYSICIAN INFORMATION	
Physician Name:	CLINIC:
Contact Information: Phone:	Email:
Fax:	Other:
Office Mailing Address:	
Please include	ance attached Diagnosis (supporting) History and Physical
the following	al progress notes