

SUTAB Colonoscopy Instructions

Failure to comply with these instructions may result in the cancellation of your procedure. Arrange for a ride home from a licensed driver after your procedure. Don't take a bus, cab, or ridesharing service unless an adult rides with you.

ONE WEEK PRIOR TO PROCEDURE:

- Do NOT inject the following medications for 7 days: Dulaglutide (Trulicity), Tirzepatide (Mounjaro, Zepbound), Exenatide (Bydureon) and Semaglutide (Ozempic, Wegovy).
- Discontinue taking: Vitamin E, Ginkgo, Omega3, Fish oil, Flax seed, Ginseng, & NSAIDs (Ibuprofen, Motrin, Advil, Excedrin, Aleve). You may take Tylenol for aches or pain if necessary. ONLY if you have heart disease, you may continue your daily Aspirin.
- If taking Coumadin, Eliquis, Plavix, Pradaxa, or Xarelto, please do NOT discontinue these medications without proper authorization from our office.
- Continue taking all other prescribed medications including heart and blood pressure medications.

TWO DAYS PRIOR TO PROCEDURE:

- Discontinue taking: Semaglutide (Rybelsus), Liraglutide (Victoza, Saxenda, Xultophy), Lixisenatide (Soliqua) and Exenatide (Byetta).
- Do NOT eat any seeded foods (seeded fruits, seeded breads, nuts, popcorn, or corn).

ONE DAY PRIOR TO PROCEDURE:

• NO SOLID FOOD Clear liquid diet for breakfast, lunch, and dinner.

DO's	DON'TS
Apple Juice, Clear Soda,	SOLID FOOD
Beef, Chicken or Vegetable Broth (Liquid Only)	Apple Sauce, Eggs, Soup, Oatmeal,
Coconut Water (No Shavings),	Orange, Grapefruit, or Pineapple Juice,
Black Coffee, or Tea (No Cream),	Milk or Dairy Products,
Jell-O or Gatorade (No Red or Purple).	Malt, Alcoholic Drinks.

- At 4 PM open a bottle of SUTAB, swallow ONE tablet every 2 mins with a sip of water from the 16oz container. (All 12 tablets & all the water must be consumed over 15-20 mins, swallowing tablets too quickly may result in nausea or vomiting).
- Approximately 1 hr after the last tablet is ingested, drink 2 more cups of water (32oz) within 30 mins.
- You may continue a clear liquid diet.
- At 9PM repeat with the other bottle of SUTAB. Approximately 1 hr after the last tablet is ingested, drink 2 more cups of water (32oz) within 30 mins. DO NOT DRINK OR EAT ANYTHING AFTER MIDIGHT.

ON THE DAY OF THE PROCEDURE:

- Please take all heart or blood pressure medications with a small sip of water.
- Do NOT take diabetic medication the morning of your procedure.
- Please bring all your medications with you to the facility.

Memorial Hospital Pembroke 7800 Sheridan St Pembroke Pines, FL 33024

(954)883-8282

SurgiCare of Miramar 14601 SW 29th St Suite 301 Miramar, FL 33027 (954)266-3801 Weston Outpatient
Surgical Center
2229 N. Commerce Pkwy
Weston, FL 33326
(954)703-3000

Procedure Date	Time of Arrival
	It will be determined by a nurse who will call you one day
	prior. We advise all patients to take the day off
	work/school.

IF YOU HAVE ANY QUESTIONS, PLEASE CONTACT OUR OFFICE AT (954)874-7900 OR EMAIL US AT HELPDESK@DRMOURA.COM