Initial Mental Health Evaluation

* Date of Assessment:

* Start Time:

* End time:

* Rendering Clinician NPI:

Supervisor License Number and NPI (if applicable):

Supervising Clinician Name and Credentials (if applicable):

Referred by:

* Rendering Clinician License Number and Type:

* Name of Client:

* Date of Birth:

* Marital Status:
- M
- S

* Employment Status:
- E
- U

CHIEF COMPLAINT & HISTORY OF PRESENT ILLNESS (HPI):

Rapid Typing Tests

PRESENTING SYMPTOMS

Check all that apply:
- frequent crying
- feeling sad, empty, or down
- loss of interest
- fatigue
- loss of interest
- decreased appetite
- hopelessness
- helplessness
- worthlessness
- anxiousness
- difficulty concentrating
- difficulty making decisions
- recurrent thoughts about death, dying
- insomnia
- excessive sleep
- loss of appetite (without weight loss)
- loss of appetite (with weight loss)
- increased appetite (without weight gain)
- increased appetite (with weight gain)
- social withdrawal, agitation

PHYSICAL Symptoms:
- feeling of edge or tense
- excessive worry
- difficulty controlling worry, difficulty concentrating
- shortness
- difficulty falling or staying asleep
- irritability

Other Behaviors:
- stories
- hallucinations
- delusions
- talking about harm
- self-harm behaviors
- visual, auditory, or tactile

OCD Compulsions:
- cleaning
- washing
- checking
- ordering and arranging
- repeating counting
- hoarding
- inappropriate sobbing

PHOBIAL Specific:

Rapid Typing Tests

PHOBIAL Generalized:

Rapid Typing Tests

Other:
- accelerated heart rate
- sounding heart
- pounding heart
- sweating
- shortness of breath
- difficulty breathing
- sensation of choking
- trembling or shaking
- feeling faint or dizzy
- nausea or abdominal distress
- abdominal pain or discomfort
- chills or heat sensations, dehydrated or flush
- pins or hot sensations
- paresthesia
- derealization
- depersonalization
- fear of losing control or "going crazy"
- fear of dying
- persistent concern or worry about additional panic attacks or their consequences
- significant, maladaptive change in behavior related to the attacks

Other:
- unpredictable sudden threat
- persistently elevated mood
- persistently irritable mood
- increased energy
- inflated self-esteem
- grandiosity
- decreased need for sleep
- more talkative than usual
- racing thoughts
- pressured speech
- flight of ideas
- distractibility
- psychomotor agitation
- increase in goal-directed activity
- excessive involvement in risky activities
- increased involvement in activities that have a high potential for painful consequences
- diminished judgment
- impulsivity

Panic Disorders (DSM):
- direct experience, witnessing, or learning of a traumatic event(s)
- recurrent involuntary and intrusive distressing memories of the event(s)
- recurrent nightmares about the event(s) distressing memories of the event(s)
- dissociative reactions (e.g. flashbacks) during which the event(s) are relived
- recurrent distressing dreams related to the event(s) - dissociative reactions (e.g. flashbacks, re-enactment) of trauma
- intense or prolonged psychological distress at exposure to internal or external cues
- physiological reactivity on exposure to internal or external cues
- persistent avoidance of stimuli associated with the event(s)
- behaviors, difficulty falling or staying asleep
- negative affectations in cognition and mood (e.g. memory)

PSYCHOTIC Hallucinations:
- auditory
- command
- visual (simple)
- visual (complex)
- tactile
- olfactory
- gustatory

PSYCHOTIC Delusion:
- of control
- of thought
- of guilt or sin
- of reference
- of persecution
- of grandiosity
- of love (erotic)
- somatic
- somatic
- thought broadcasting
- thought insertion
- bizarre
- mood congruent
- mood incongruent
- mood neutral
- disorganized speech
- disorganized behavior

ADHD
- inattention
- distractibility
- hyperactivity and impulsivity

EATING DISORDERS (BINGE/ED)
- refusing or restricting food
- burning skin
- refusing or limiting eating
- puffing out hair
- falling head
- losing body hair

Non-ABUSIVE BEHAVIOR Assertions/Ingestions of Aggression:
- verbal
- in objects
- in self
- bedwetting

EATING DISORDERS (BINGE/ED)
- debarking
- biting
- pinning
- excessive exercise
- use of laxatives or diuretics
- use of appetite suppressants

Additional:

Rapid Typing Tests

FUNCTIONAL IMPAIRMENT

Occupational Functioning:

1
2
3
4
5

Academic Functioning:

1
2
3
4
5

Family Functioning:

1
2
3
4
5

Marital Functioning:

1
2
3
4
5

Is the Child, Adolescent(s):

Social Functioning - Describe Specific Issues:

Rapid Typing Tests

PSYCHOSOCIAL HISTORY

Family of Origin (household composition, key relationships, family roles, significant dynamics):

Rapid Typing Tests

Developmental History (birth complications, developmental milestones, early attachment):

Rapid Typing Tests

Legal History (arrests, incarcerations, current legal involvement, probation or parole):

Rapid Typing Tests

Vocational History:

Rapid Typing Tests

School History:

Rapid Typing Tests

Family History of Mental Illness and/or Substance Abuse:

Rapid Typing Tests

History of Problems:

Rapid Typing Tests

Current Outpatient Mental Health Treatment:

Rapid Typing Tests

Current Psychotropic Medications:

Rapid Typing Tests

Psychiatric Hospitalizations

Total number:

Where and when admitted?

Rapid Typing Tests

Current Outpatient Mental Health Treatment (provider, frequency, modality):

Rapid Typing Tests

Current Psychotropic Medications (name, dose, prescriber - include OTC and supplements):

Rapid Typing Tests

Total Number of Psychiatric Hospitalizations:

Rapid Typing Tests

Psychiatric Visualization History (facility, date, duration, reason – must match final):

Rapid Typing Tests

Prior Outpatient / PHP / IOP Treatment History:

Rapid Typing Tests

History of Suicide Attempt(s)?

Rapid Typing Tests

History of Non-Lethal Self-Injury?

Rapid Typing Tests

History of Violence Toward Others?

Rapid Typing Tests

Harm to Cruelty to Animals?

Rapid Typing Tests

Details (describe any No responses above – incidents, dates, lethality, circumstances):

Rapid Typing Tests

Outpatient Treatments:

Rapid Typing Tests

Violent Behaviors: Self-Harm

Recent suicidal attempts?
- Yes
- No

Recent use of non-lethal self-injury?
- Yes
- No

Harm or Cruelty to animals?
- Yes
- No

Additional Psychosocial History Information:

Medical History

Current Medical Conditions (include all chronic conditions and approximate onset date):

Rapid Typing Tests

Current Non-Psychiatric Medications (name, dose, prescriber – include OTC and supplements):

Rapid Typing Tests

Primary Care Provider (PCP) Name and Contact:

Rapid Typing Tests

Head Trauma?
- Yes
- No

Seizures?
- Yes
- No

Sleep Disorder?
- Yes
- No

SUBSTANCE USE HISTORY

Substance Use:
- Alcohol
- Tobacco
- Cannabis
- Opioids
- Cocaine
- Amphetamines
- Benzodiazepines
- Other
- Other Substances - Use Status, Frequency, Quantity, Last Use

Substance Abuse Treatment History:

Rapid Typing Tests

Additional Medical History:

Rapid Typing Tests

ADVERSE CHILDHOOD EXPERIENCES (ACE) & TRAUMA HISTORY

- Physical Abuse
- Emotional
- Physical
- Sexual
- Neglect
- Household Dysfunction

Adverse Childhood Experiences:
- Mental Illness in Household
- Substance Abuse in Household
- Parental Separation or Divorce (Parental Absence)
- Incarcerated Household Member
- Parental separation or divorce

Additional Trauma History (childhood, community violence, medical trauma, other):

Rapid Typing Tests

Trauma-Informed Care Considerations (sensitivity areas, triggers identified, care adaptations):

Rapid Typing Tests

ACE SCORE:

Risk Factors:
- Male gender
- Adolescent, young adult, or elderly age*
- Single, divorced, or widowed
- Unemployed or retired
- Access to lethal means (firearms, stockpiled medications)
- Recent discharge from psychiatric hospitalization
- Prior suicide attempts
- Family history of suicide or close friend
- Current or recent substance use
- Recent loss (relationship, job, housing)
- Stress identified
- Poor coping skills
- Chronic pain or serious medical illness
- Social isolation
- Lack of social support
- Domestic violence or unsafe living situation

Protective Factors:
- Responsibility to family or children
- Reasons for living identified
- Perceived social support
- Other
- Family support
- Future orientation / reasons for living
- Currently engaged in treatment
- Effective coping skills
- Moral or religious objection to suicide
- Positive coping skills
- Family support present
- Employment, structured daily routine

SAFETY, RISK DETERMINATION
- Current Suicidal Ideation or Intoxication
- Current Homicidal Ideation

CULTURAL & LINGUISTIC ASSESSMENT

Primary Language:

Is an Interpreter Needed?

Was an Interpreter Used This Session?

Race / Ethnicity (client-reported, optional):

Cultural Considerations Relevant to Assessment and Treatment:

Health Literacy Level:

COORDINATION OF CARE

ROI Released for PCP Communication?

ROI Released for Psychiatry / Prescribing Provider?

Coordination Completed This Session?

Coordination of Care Notes (other providers, referrals made, referrals received):

Rapid Typing Tests

MENTAL STATUS EXAMINATION (MSE)

General Appearance and Behavior:
- Neat
- Well-kept
- clean, dirty
- neatly dressed
- appropriately dressed
- inappropriately dressed
- disheveled
- unkempt
- overweight
- slight build
- well-nourished
- underweight
- obese
- poor hygiene, unkempt appearance
- stiff posture
- slumped posture
- tremors
- fidgeting
- functioning leg
- nail biting
- arm crossing
- pulling hair
- hand wringing
- hand tapping
- rocking self

Eye Contact:
- Poor
- Fair
- Good

Manner:
- Normal
- dramatic
- exaggerated
- aloof
- guarded
- shy
- seductive
- ingratiating
- hostile
- friendly
- inappropriately familiar
- defensive
- easily irritated
- cooperative
- suspicious

Relationship to Examiner:
- Attentive
- Distracted
- Interested