



Lifestyle and Clinical Correlates of Hepatocellular Carcinoma in South Texas: A Matched Case-control Study

Amelie G. Ramirez,^{*,‡} Edgar Muñoz,^{*} Dorothy Long Parma,^{*} Joel E. Michalek,[‡] Alan E. C. Holden,^{*} Timothy D. Phillips,[§] and Bradley H. Pollock^{||}

^{*}Institute for Health Promotion Research, University of Texas Health Science Center at San Antonio, San Antonio, Texas;

[‡]Department of Epidemiology and Biostatistics, University of Texas Health Science Center at San Antonio, San Antonio, Texas;

[§]Department of Veterinary Integrative Biosciences, Texas A&M University, College Station, Texas; and ^{||}Department of Public Health Sciences, School of Medicine, University of California, Davis, Davis, California

The incidence of hepatocellular carcinoma (HCC) in the United States is rising even as overall cancer rates are declining, and Latinos are disproportionately affected, especially in Texas. Our case-control study sought to determine relative etiologic contributions of lifestyle-related risk factors in South Texas.

Methods

Between October 2012 and July 2014, we identified, consented, and interviewed in person 51 HCC adult cases (diagnosed within the past 12 months and residing in Bexar or any of the 7 surrounding counties) from clinics at University Hospital and the Cancer Therapy and Research Center at UT Health San Antonio and 104 adult controls, 32 from a related study^{1,2} and 72 randomly selected residents in the study counties. From these 51 cases and 104 controls we created 42 matched pairs, with exact matching on sex, ethnicity (Hispanic, non-Hispanic), and age category (18–57 years, >57 years). Urine samples (10 mL each) and serum samples (150 μ L each) were assessed for biomarkers of previous aflatoxin exposure by using procedures previously described.³ The primary aflatoxin exposure serum biomarker was an aflatoxin B₁ (AFB₁)-lysine DNA adduct. We also assessed urinary aflatoxin M₁. We used matched logistic regression to estimate odds ratios and 95% confidence intervals. The protocol was approved by a local Institutional Review Board.

Results

Most matched cases and controls were Latino (67%). Compared with controls, HCC cases were more likely to report Medicare or Medicaid, lower income, less education, more lifetime alcohol use, and more smoking (Figure 1). Cases were less likely to report hypercholesterolemia (0.11 [0.02–0.51]), more likely to report hepatitis C (183.74 [27.37– ∞]), cirrhosis (2.17 [33.3– ∞]), and transfusions (4.35 [1.60–11.84]) and less likely to be taking aspirin (0.31 [0.11–0.85]), statins

(0.03 [0–0.20]), and omega-3/fish oil (0.10 [0.01–0.78]). Cases did not differ significantly from controls with regard to reported consumption of products made from corn (data not shown). Relative to controls, cases were more likely to have hepatitis C virus (HCV) antibodies (174.3 [26.2– ∞]) and have detectable aflatoxin levels in blood (6.09 [1.10–33.71]) and urine (3.42 [1.07–10.91]).

Discussion


This is an epidemiologic study relating HCC and aflatoxin exposure in a U.S. population, particularly a Latino-majority study population. A recent review of non-U.S. studies⁴ reported that aflatoxin may play a causative role in 4.6%–28.2% of all global HCC cases, and that the combined effect of AFB₁ exposure and hepatitis B virus infection appears additive rather than multiplicative. In the current study, serum (AFB₁) and urine (aflatoxin M₁) aflatoxin levels and HCV infection were significantly higher in cases than in controls. Our results also support the observations of others that HCV infection is a major risk factor for HCC incidence in the United States. Although HCV infection is a strong risk factor for development of HCC, rising trends in HCC incidence among Latinos cannot be attributed solely to HCV infection. Our data on a majority-Latino cohort suggest that HCC cases were poor and controls were more affluent, and cases smoked more with a higher prevalence of cirrhosis than controls. The lack of case-control differences for corn-based foods, a suspected risk factor for HCC, may be due to information bias from use of a 12-month dietary recall assessment. We hypothesized that contaminated corn products contributed to the increased risk of HCC and expected to

Abbreviations used in this paper: AFB₁, aflatoxin B₁; HCC, hepatocellular carcinoma; HCV, hepatitis C virus.

Most current article

© 2017 by the AGA Institute
1542-3565/\$36.00

<http://dx.doi.org/10.1016/j.cgh.2017.03.022>



*median unbiased estimate (MUE) from exact logistic regression

Figure 1. Association between lifestyle and clinical factors and HCC risk in South Texas. Odds ratios (95% confidence interval [CI]) derived from matched logistic regression. HBsAg, hepatitis B surface antigen.

observe increased consumption of corn-based foods in cases relative to controls; a possible reason may be that cases and controls are consuming corn products differentially contaminated with aflatoxin (eg, commercial vs home-grown). Random assays of corn products purchased by cases and controls should be done in a future study to verify this.

Study strengths include case-control matching on sex, ethnicity, and age and in-person interviewing of subjects in their language of choice. A possible weakness is that cases and controls were not randomly selected with the same inclusion criteria; therefore, selection bias may exist. Cases were selected as a convenience sample from local medical facilities, whereas controls were selected from an existing prevention study and randomly from the surrounding county populations. A larger study will be needed to address sources of aflatoxin exposure as a risk factor for HCC.

References

1. ClinicalTrials.gov. Phase 2 reduction of dietary mycotoxin exposure by ACCS100 (RDMEACCS100). Available from: <https://clinicaltrials.gov/ct2/show/NCT01677195?term=NCT01677195&rank=1>. Accessed March 24, 2015.
2. Pollock BH, Elmore S, Romoser A, et al. Intervention trial with calcium montmorillonite clay in a south Texas population

exposed to aflatoxin. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 2016;33:1346-1354.

3. Johnson NM, Qian G, Xu L, et al. Aflatoxin and PAH exposure biomarkers in a U.S. population with a high incidence of hepatocellular carcinoma. *Sci Total Environ* 2010;408:6027-6031.
4. Wu HC, Santella R. The role of aflatoxins in hepatocellular carcinoma. *Hepat Mon* 2012;12:e7238.

Reprint requests

Address requests for reprints to: Amelie G. Ramirez, DrPH, Institute for Health Promotion Research, Department of Epidemiology and Biostatistics, University of Texas Health Science Center at San Antonio, 7411 John Smith Drive, Suite 1000, San Antonio, Texas 78229. e-mail: ramirezag@uthscsa.edu; fax: (210) 562-6545.

Acknowledgments

The authors thank Amanda Sintes for her assistance with interviews of participants and Allen Trevino, Holiday Harris, and Norma Ketchum for their assistance with data management and their comments on an earlier version of the manuscript.

Conflicts of interest

The authors disclose no conflicts.

Funding

Supported by the Cancer Prevention Research Institute of Texas (CPRI) RP120462, the Institute for Health Promotion Research (IHPR) U01 CA86117 and U54 CA153511, and the Cancer Therapy and Research Center (CTRC) at The University of Texas Health Science Center at San Antonio, an NCI-designated Cancer Center (P30 CA054174).