


AP Physics 2: Algebra-Based Free-Response Questions SOLUTION

FORMULAE



Que-1

$$\mu = \frac{v_a}{v_m} = \frac{f \lambda_a}{f \lambda_m}$$

$f \rightarrow$ constant

$$\mu = \frac{5m}{1m}$$

Que-2

$$V = IR$$

$$Q = CV$$

$$C_{eq} = C_1 + C_2 \text{ (series)}$$

$$\frac{1}{C_{eq}} = \frac{1}{C_1} + \frac{1}{C_2} \text{ (parallel)}$$

Que-3

$$f = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2}$$

$$PE = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r}$$

Que-4

$$F = q(\vec{E} \times \vec{B})$$

$$B = \frac{\mu_0}{4\pi} \frac{2I}{r}$$

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

+91-9958461445

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

**Online Physics
Classes/Tutor
AP Physics (2)
Algebra-Based 2022
Paper Solution**

**AP, IB DP HL/SL, IGCSE, A-LEVEL, O-
LEVEL, MCAT, ACT, NEET, IIT**

Kumar Physics classes - www.kumarphysicsclasses.com

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

AP[®] PHYSICS 2 TABLE OF INFORMATION

CONSTANTS AND CONVERSION FACTORS

Proton mass, $m_p = 1.67 \times 10^{-27}$ kg Neutron mass, $m_n = 1.67 \times 10^{-27}$ kg Electron mass, $m_e = 9.11 \times 10^{-31}$ kg Avogadro's number, $N_0 = 6.02 \times 10^{23}$ mol ⁻¹ Universal gas constant, $R = 8.31$ J/(mol·K) Boltzmann's constant, $k_B = 1.38 \times 10^{-23}$ J/K	Electron charge magnitude, $e = 1.60 \times 10^{-19}$ C 1 electron volt, $1 \text{ eV} = 1.60 \times 10^{-19}$ J Speed of light, $c = 3.00 \times 10^8$ m/s Universal gravitational constant, $G = 6.67 \times 10^{-11}$ m ³ /kg·s ² Acceleration due to gravity at Earth's surface, $g = 9.8$ m/s ²
1 unified atomic mass unit, Planck's constant, Vacuum permittivity, Coulomb's law constant, $k = 1/4\pi\epsilon_0 = 9.0 \times 10^9$ N·m ² /C ² Vacuum permeability, Magnetic constant, $k' = \mu_0/4\pi = 1 \times 10^{-7}$ (T·m)/A 1 atmosphere pressure,	$1 \text{ u} = 1.66 \times 10^{-27}$ kg = 931 MeV/c ² $h = 6.63 \times 10^{-34}$ J·s = 4.14 × 10 ⁻¹⁵ eV·s $hc = 1.99 \times 10^{-25}$ J·m = 1.24 × 10 ³ eV·nm $\epsilon_0 = 8.85 \times 10^{-12}$ C ² /N·m ² $\mu_0 = 4\pi \times 10^{-7}$ (T·m)/A $1 \text{ atm} = 1.0 \times 10^5$ N/m ² = 1.0 × 10 ⁵ Pa

UNIT SYMBOLS	meter, m	mole, mol	watt, W	farad, F
	kilogram, kg	hertz, Hz	coulomb, C	tesla, T
	second, s	newton, N	volt, V	degree Celsius, °C
	ampere, A	pascal, Pa	ohm, Ω	electron volt, eV
	kelvin, K	joule, J	henry, H	

PREFIXES		
Factor	Prefix	Symbol
10 ¹²	tera	T
10 ⁹	giga	G
10 ⁶	mega	M
10 ³	kilo	k
10 ⁻²	centi	c
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES							
θ	0°	30°	37°	45°	53°	60°	90°
$\sin \theta$	0	1/2	3/5	$\sqrt{2}/2$	4/5	$\sqrt{3}/2$	1
$\cos \theta$	1	$\sqrt{3}/2$	4/5	$\sqrt{2}/2$	3/5	1/2	0
$\tan \theta$	0	$\sqrt{3}/3$	3/4	1	4/3	$\sqrt{3}$	∞

The following conventions are used in this exam.

- I. The frame of reference of any problem is assumed to be inertial unless otherwise stated.
- II. In all situations, positive work is defined as work done on a system.
- III. The direction of current is conventional current: the direction in which positive charge would drift.
- IV. Assume all batteries and meters are ideal unless otherwise stated.
- V. Assume edge effects for the electric field of a parallel plate capacitor unless otherwise stated.
- VI. For any isolated electrically charged object, the electric potential is defined as zero at infinite distance from the charged object

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

AP[®] PHYSICS 2 EQUATIONS

MECHANICS

$$v_x = v_{x0} + a_x t$$

$$x = x_0 + v_{x0} t + \frac{1}{2} a_x t^2$$

$$v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$$

$$\vec{a} = \frac{\sum \vec{F}}{m} = \frac{\vec{F}_{net}}{m}$$

$$|\vec{F}_f| \leq \mu |\vec{F}_n|$$

$$a_c = \frac{v^2}{r}$$

$$\vec{p} = m\vec{v}$$

$$\Delta\vec{p} = \vec{F} \Delta t$$

$$K = \frac{1}{2} m v^2$$

$$\Delta E = W = F_{\parallel} d = F d \cos \theta$$

$$P = \frac{\Delta E}{\Delta t}$$

$$\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$$

$$\omega = \omega_0 + \alpha t$$

$$x = A \cos(\omega t) = A \cos(2\pi f t)$$

$$x_{cm} = \frac{\sum m_i x_i}{\sum m_i}$$

$$\vec{\alpha} = \frac{\sum \vec{\tau}}{I} = \frac{\vec{\tau}_{net}}{I}$$

$$\tau = r_{\perp} F = r F \sin \theta$$

$$L = I \omega$$

$$\Delta L = \tau \Delta t$$

$$K = \frac{1}{2} I \omega^2$$

$$|\vec{F}_s| = k |\vec{x}|$$

a = acceleration
 A = amplitude
 d = distance
 E = energy
 F = force
 f = frequency
 I = rotational inertia
 K = kinetic energy
 k = spring constant
 L = angular momentum
 ℓ = length
 m = mass
 P = power
 p = momentum
 r = radius or separation
 T = period
 t = time
 U = potential energy
 v = speed
 W = work done on a system
 x = position
 y = height
 α = angular acceleration
 μ = coefficient of friction
 θ = angle
 τ = torque
 ω = angular speed

$$U_s = \frac{1}{2} k x^2$$

$$\Delta U_g = mg \Delta y$$

$$T = \frac{2\pi}{\omega} = \frac{1}{f}$$

$$T_s = 2\pi \sqrt{\frac{m}{k}}$$

$$T_p = 2\pi \sqrt{\frac{\ell}{g}}$$

$$|\vec{F}_g| = G \frac{m_1 m_2}{r^2}$$

$$\vec{g} = \frac{\vec{F}_g}{m}$$

$$U_G = -\frac{G m_1 m_2}{r}$$

ELECTRICITY AND MAGNETISM

$$|\vec{F}_E| = \frac{1}{4\pi\epsilon_0} \frac{|q_1 q_2|}{r^2}$$

$$\vec{E} = \frac{\vec{F}_E}{q}$$

$$|\vec{E}| = \frac{1}{4\pi\epsilon_0} \frac{|q|}{r^2}$$

$$\Delta U_E = q \Delta V$$

$$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$$

$$|\vec{E}| = \left| \frac{\Delta V}{\Delta r} \right|$$

$$\Delta V = \frac{Q}{C}$$

$$C = \kappa \epsilon_0 \frac{A}{d}$$

$$E = \frac{Q}{\epsilon_0 A}$$

$$U_C = \frac{1}{2} Q \Delta V = \frac{1}{2} C (\Delta V)^2$$

$$I = \frac{\Delta Q}{\Delta t}$$

$$R = \frac{\rho \ell}{A}$$

$$P = I \Delta V$$

$$I = \frac{\Delta V}{R}$$

$$R_s = \sum_i R_i$$

$$\frac{1}{R_p} = \sum_i \frac{1}{R_i}$$

$$C_p = \sum_i C_i$$

$$\frac{1}{C_s} = \sum_i \frac{1}{C_i}$$

$$B = \frac{\mu_0 I}{2\pi r}$$

A = area
 B = magnetic field
 C = capacitance
 d = distance
 E = electric field
 \mathcal{E} = emf
 F = force
 I = current
 ℓ = length
 P = power
 Q = charge
 q = point charge
 R = resistance
 r = separation
 t = time
 U = potential (stored) energy
 V = electric potential
 v = speed
 κ = dielectric constant
 ρ = resistivity
 θ = angle
 Φ = flux

$$\vec{F}_M = q\vec{v} \times \vec{B}$$

$$|\vec{F}_M| = |q\vec{v}| |\sin \theta| |\vec{B}|$$

$$\vec{F}_M = I\vec{\ell} \times \vec{B}$$

$$|\vec{F}_M| = |I\vec{\ell}| |\sin \theta| |\vec{B}|$$

$$\Phi_B = \vec{B} \cdot \vec{A}$$

$$\Phi_B = |\vec{B}| \cos \theta |\vec{A}|$$

$$\mathcal{E} = -\frac{\Delta \Phi_B}{\Delta t}$$

$$\mathcal{E} = B \ell v$$

KUMAR PHYSICS CLASSES


E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

AP[®] PHYSICS 2 EQUATIONS

FLUID MECHANICS AND THERMAL PHYSICS	WAVES AND OPTICS
$\rho = \frac{m}{V}$ $P = \frac{F}{A}$ $P = P_0 + \rho gh$ $F_b = \rho Vg$ $A_1 v_1 = A_2 v_2$ $P_1 + \rho g y_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2} \rho v_2^2$ $\frac{Q}{\Delta t} = \frac{kA \Delta T}{L}$ $PV = nRT = Nk_B T$ $K = \frac{3}{2} k_B T$ $W = -P \Delta V$ $\Delta U = Q + W$	$\lambda = \frac{v}{f}$ $n = \frac{c}{v}$ $n_1 \sin \theta_1 = n_2 \sin \theta_2$ $\frac{1}{s_i} + \frac{1}{s_o} = \frac{1}{f}$ $ M = \left \frac{h_i}{h_o} \right = \left \frac{s_i}{s_o} \right $ $\Delta L = m\lambda$ $d \sin \theta = m\lambda$
$A = \text{area}$ $F = \text{force}$ $h = \text{depth}$ $k = \text{thermal conductivity}$ $K = \text{kinetic energy}$ $L = \text{thickness}$ $m = \text{mass}$ $n = \text{number of moles}$ $N = \text{number of molecules}$ $P = \text{pressure}$ $Q = \text{energy transferred to a system by heating}$ $T = \text{temperature}$ $t = \text{time}$ $U = \text{internal energy}$ $V = \text{volume}$ $v = \text{speed}$ $W = \text{work done on a system}$ $y = \text{height}$ $\rho = \text{density}$	$d = \text{separation}$ $f = \text{frequency or focal length}$ $h = \text{height}$ $L = \text{distance}$ $M = \text{magnification}$ $m = \text{an integer}$ $n = \text{index of refraction}$ $s = \text{distance}$ $v = \text{speed}$ $\lambda = \text{wavelength}$ $\theta = \text{angle}$
MODERN PHYSICS	GEOMETRY AND TRIGONOMETRY
$E = hf$ $K_{\max} = hf - \phi$ $\lambda = \frac{h}{p}$ $E = mc^2$	$A = \text{area}$ $C = \text{circumference}$ $V = \text{volume}$ $S = \text{surface area}$ $b = \text{base}$ $h = \text{height}$ $\ell = \text{length}$ $w = \text{width}$ $r = \text{radius}$
$E = \text{energy}$ $f = \text{frequency}$ $K = \text{kinetic energy}$ $m = \text{mass}$ $p = \text{momentum}$ $\lambda = \text{wavelength}$ $\phi = \text{work function}$	$A = bh$ $A = \frac{1}{2}bh$ $A = \pi r^2$ $C = 2\pi r$ $V = \ell wh$ $V = \pi r^2 \ell$ $S = 2\pi r \ell + 2\pi r^2$ $V = \frac{4}{3}\pi r^3$ $S = 4\pi r^2$ $c^2 = a^2 + b^2$ $\sin \theta = \frac{a}{c}$ $\cos \theta = \frac{b}{c}$ $\tan \theta = \frac{a}{b}$
	

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

Begin your response to **QUESTION 1** on this page.

PHYSICS 2 SECTION II Time—1 hour and 30 minutes 4 Questions

Directions: Questions 1 and 4 are short free-response questions that require about 20 minutes each to answer and are worth 10 points each.

Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each.

Show your work for each part in the space provided after that part.

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445,01141032244

www.kumarphysicsclasses.com



www.kumarneetphysicsclasses.com

1. (10 points, suggested time 20 minutes)

Students are investigating electromagnetic wave phenomena in transparent media. They use a string to support a stationary thin, rectangular block of mass m_b , volume V_b and density ρ_b . The block has two narrow slits in its center and is submerged in a glass tank containing water with density ρ_w , as shown above.


(a)

i. On the dot below, which represents the block, draw and label the forces that are exerted on the block. Each force must be represented by a distinct arrow starting on, and pointing away from, the dot.



ii. Derive an expression for the force exerted on the block by the string in terms of the given quantities and physical constants, as appropriate.

Since the block is
under equilibrium
position



$$F_b + F_T - mg = 0$$

$$F_T = mg - F_b$$

$$= m_b g - \rho_w V_b g$$

(b) A monochromatic laser beam is incident perpendicular to the wall of the tank. The beam passes through the slits in the block. An interference pattern is formed on the screen inside the tank. The water is then replaced with a clear fluid with a greater index of refraction than that of water. In a coherent, paragraph-length response, describe how the greater index of refraction of the new fluid affects the interference pattern. Explain your reasoning in terms of speed, frequency, and wavelength of the light.

<p>WATER</p> <p>μ_g</p> <p>f - constant</p> <p>$\mu_g = \frac{\lambda_a}{\lambda_g}$</p> <p>$\lambda_g = \frac{\lambda_a}{\mu_g}$</p>	<p>MEDIUM</p> <p>$\mu_m > \mu_g$</p> <p>f - constant</p> <p>$\mu_m = \frac{\lambda_a}{\lambda_m}$</p> <p>$\lambda_m = \frac{\lambda_a}{\mu_m}$</p>	<p>$\mu_m > \mu_g$</p>
---	---	--------------------------------------

$\lambda_m < \lambda_a$

we know that $d \sin \theta = n \lambda$ since λ decreases
 $d \rightarrow$ constant

hence θ decreases then interference pattern.
 will decrease in width

(c) The block is replaced by a triangular prism, as shown above. The path of the beam is indicated by the dotted line, and the beam reaches the screen at point P. The fluid is then removed from the tank, and the prism is surrounded by air. Predict whether the beam will reach the side of the tank above point P, at point P, or below point P when the prism is surrounded by air. Support your answer using physics principles.

CASE - I

$$\mu_g \sin i$$

$$= \mu_f \sin r_1$$

$$\sin r_1 = \frac{\mu_g \sin i}{\mu_f}$$


CASE - 2

$$\mu_g \sin i$$

$$= \mu_a \sin r_2$$

$$\sin r_2 = \frac{\mu_g \sin i}{\mu_a}$$

Laser



Both the cases $\sin i \rightarrow$ constant

μ_g - constant

But $\mu_f > \mu_a$

then $\sin r_1 < \sin r_2$

$$r_1 < r_2 \Rightarrow r_2 > r_1$$

Hence the beam below to point P.

KUMAR PHYSICS CLASSES


E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

Begin your response to QUESTION 2 on this page.



2. (12 points, suggested time 25 minutes)

Students perform an experiment with a battery and four resistors, A, B, C, and D. The resistance of resistors A and C is $R_A = R_C = R$. The resistance of resistors B and D is $R_B = R_D = 2R$. The students create the two circuits shown above and measure the potential differences ΔV_A , ΔV_B , ΔV_C , and ΔV_D across resistors A, B, C, and D, respectively.


(a) From greatest to least, rank the magnitudes of the potential differences across the resistors. Use "1" for the greatest magnitude, "2" for the next greatest magnitude, and so on. If any potential differences have the same magnitude, use the same number for their ranking.

1 ΔV_A 1 ΔV_B 2 ΔV_C 3 ΔV_D Justify your answer.

$$\Delta V_A = V, \quad \Delta V_B = \left(\frac{V}{2R}\right)(2R) = V$$

$$\Delta V_C = \left(\frac{V}{3R}\right)(R) = \frac{V}{3} = \Delta V_C$$

$$\Delta V_D = \left(\frac{V}{3R}\right)(2R) = \frac{2V}{3} = \Delta V_D$$



$$\begin{aligned} \Delta V_A &= V \\ \Delta V_B &= V \\ \Delta V_C &= \frac{V}{3} \\ \Delta V_D &= \frac{2V}{3} \end{aligned} \quad \Delta V_D > \Delta V_C > \Delta V_B = \Delta V_A$$

In another experiment, the students have a capacitor with unknown capacitance C_U . They want to determine C_U by using a battery of potential difference 4.5 V and several other capacitors of known capacitance. They create circuits with the battery, the unknown capacitor, and one of the capacitors of known capacitance. The students wait until the capacitors are fully charged and then record the potential difference ΔV_{known} across the known capacitor and the potential difference ΔV_U across the unknown capacitor. Their data are shown in the table on the following page.

↓ ΔV_U

(b)

i. Calculate the amount of charge on the capacitor of known capacitance of 200 μF in the students' experiment.

$$Q = C \cdot \Delta V$$

$$= (200 \mu\text{F}) (0.91)$$

$$= 181 \text{ C}$$

Known Capacitance of Capacitors (μF)	ΔV_{known} (V)	ΔV_U (V)		
200	0.91	3.53		
300	0.65	3.74		
400	0.51	3.95		
500	0.42	4.06		
600	0.36	4.17		

ii. Briefly explain why the data in the table provide evidence that the capacitors are connected in series.

For series $\frac{1}{C_{\text{TOTAL}}} = \frac{1}{C_{\text{KNOWN}}} + \frac{1}{C_{\text{UNKNOWN}}}$

FROM TABLE C_{KNOWN} starts increasing, then C_{TOTAL} decreasing.
 $\Delta V_{\text{KNOWN}} = \frac{Q}{C_{\text{KNOWN}}}$ starts decreasing, then Q decreases.

KUMAR PHYSICS CLASSES
 E-2B BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI
 9958461445, 01141032244
 www.kumarphysicsclasses.com
 www.kumarneetphysicsclasses.com

iii. Briefly explain why connecting the capacitors in parallel would not provide enough information to determine the capacitance of the unknown capacitor if the only measuring device available is a voltmeter.

when parallel $C_{eq} = C_1 + C_2$

Hence $C_{equivalent}$ increased

$$\Delta V = \frac{Q}{C_{equivalent}}$$

make potential very small
therefore difficult to
measure by Voltmeter

Continue your response to QUESTION 2 on this page.


(c) The students want to produce a linear graph of the data so that the capacitance C_U of the unknown capacitor can be determined from the slope of the best-fit line for the data.

i. Indicate two quantities that could be plotted to produce the desired graph. Use the empty columns of the data table in part (b) to record any values that you need to calculate.

Vertical axis _____ Horizontal axis _____

ii. Label the axes below and provide an appropriate scale with units. Plot the data points for the quantities indicated in part (c)(i) on the axes and draw a best-fit line.

Known Capacitance of Capacitors (μF)	ΔV_{known} (V)	ΔV_U (V)	Q_{KNOWN} $= (\Delta V_{\text{KNOWN}}) \times C_{\text{KNOWN}}$
200	0.91	3.53	182 μC
300	0.65	3.74	195 μC
400	0.51	3.95	204 μC
500	0.42	4.06	210 μC
600	0.36	4.17	216 μC




iii. Using your best-fit line, determine the capacitance of capacitor C_U .

$C_U = \frac{\Delta Q_{\text{KNOWN}}}{\Delta V_{\text{UNKNOWN}}}$
 $= \frac{(207 - 187) \mu\text{C}}{(4 - 3.6) \text{V}} = \frac{20 \mu\text{C}}{0.4 \text{V}} = 50 \mu\text{F} = C_U$

Series hence charge constant

Begin your response to QUESTION 3 on this page.

3. (12 points, suggested time 25 minutes)



Note: Figure not drawn to scale.

A hydrogen atom can be modeled as an electron in a circular orbit of radius r about a stationary proton, as shown above. The gravitational force between the proton and electron is negligible compared to the electrostatic force between them.

(a) Derive an equation for the speed v of the electron in terms of r and physical constants, as appropriate.

$$\frac{mv^2}{r} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \Rightarrow v = \sqrt{\frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r(m)}} = \sqrt{\frac{1}{4\pi\epsilon_0} \frac{e^2}{r(me)}}$$

(b) Derive an equation for the total energy of the atom in terms of r and physical constants, as appropriate.

$$\begin{aligned} TE &= PE + KE \\ &= \frac{1}{4\pi\epsilon_0} \frac{(e)(-e)}{r} + \frac{1}{2}mv^2 \\ &= -\frac{1}{4\pi\epsilon_0} \frac{e^2}{r} + \left(\frac{1}{4\pi\epsilon_0} \frac{e^2}{r}\right) \left(\frac{1}{2}\right) \\ &= -\frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{e^2}{r}\right) \end{aligned}$$

$$\begin{aligned} \frac{mv^2}{r} &= \frac{1}{4\pi\epsilon_0} \frac{e \cdot e}{r^2} \\ \frac{1}{2}mv^2 &= \frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{e^2}{r}\right) = KE \end{aligned}$$

(c) When the hydrogen atom absorbs a photon, the electron moves to an orbit with a larger radius and the total energy of the atom increases. Is your equation for the energy derived in part (b) consistent with this description of the model of a hydrogen atom absorbing a photon? Explain why the equation is or is not consistent.

The equation in (b)

$$TE = -\frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{e^2}{r} \right)$$

If $r \uparrow \uparrow$ then it becomes less negative
It means total energy increases. This
is consistent with the given description
of the atom absorbing a photon.

(d) Experiments show that a hydrogen atom can absorb a photon of frequency 3.2×10^{15} Hz.

i. Calculate the energy of a photon with this frequency.

$$\begin{aligned} E &= h f = 6.6 \times 10^{-34} \times 3.2 \times 10^{15} \\ &= 2.12 \times 10^{-18} \text{ J} \end{aligned}$$

ii. A student claims that when a hydrogen atom absorbs a photon at this frequency, the energy could be converted into mass, adding an electron to the atom. Calculate the amount of energy needed to create a particle with the mass of an electron and determine whether or not there is sufficient energy gained by the atom to add another electron.

$$\begin{aligned}
 E_{\text{electron}} &= m c^2 \\
 &= (9.1 \times 10^{-31}) (3 \times 10^8)^2 \\
 &= 8.20 \times 10^{-14} \text{ J}
 \end{aligned}$$

From d (i) — energy of photon

$$\begin{aligned}
 E &= h f = 6.6 \times 10^{-34} \times 3.2 \times 10^{15} \\
 &= 2.12 \times 10^{-18} \text{ J}
 \end{aligned}$$

$$\frac{E_{\text{electron}}}{E_{\text{photon}}} = \frac{8.20 \times 10^{-14}}{2.12 \times 10^{-18}} \approx 4 \times 10^4 \approx 4000$$

↓
 Around this much
 photon need to be
 absorbed to produce
 1 electron.


KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI


9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com



iii. The left bar chart in the figure above is complete and represents the initial electric potential energy $U_{E,i}$ of the atom and the initial kinetic energy K_i of the electron before the photon is absorbed. In the space provided on the right, draw a bar chart to represent a possible final electric potential energy of the atom and final kinetic energy of the electron.



$$PE = - \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r}$$

$$KE = \frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r} \right)$$

$$TE = - \frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r} \right)$$

When photon is absorbed then electron will reach to upper orbit $r \uparrow \uparrow$

- PE (Less -ve) \rightarrow magnitude \downarrow
- KE (Less +ve) \rightarrow magnitude \downarrow
- TE (Less -ve) \rightarrow magnitude \downarrow

Begin your response to QUESTION 4 on this page.


4. (10 points, suggested time 20 minutes)


At the instant shown above, a negatively charged object is moving to the left with constant velocity v near a long, straight wire that has a current I directed to the left. The region contains a uniform electric field of magnitude E , and the charged object is at a distance d from the wire. The figure shows the electric and magnetic forces, F_E and F_M respectively, exerted on the charged object.

(a) Derive an expression for v in terms of E , d , I , and physical constants, as appropriate.

$$F_M = F_E$$
$$\cancel{q} v B = \cancel{q} E$$
$$v \left(\frac{\mu_0}{4\pi} \frac{2I}{d} \right) = E$$


$$v = \frac{2\pi d E}{\mu_0 I}$$





(b) The charged object is removed, and a square coil with side length $2L$ is placed near the long, straight wire, as shown above. The bottom of the coil is a distance L from the wire. The magnitude of the magnetic field due to the current in the wire is $3B_0$ at point P_1 and B_0 at point P_2 .

i. Write an "X" at a location on the figure where the magnitude of the magnetic field is $2B_0$. Briefly justify your reasoning.



$$B_0 = \frac{\mu_0}{4\pi} \frac{2I}{3L} \quad \text{--- (1)}$$

$$3B_0 = \frac{\mu_0}{4\pi} \frac{2I}{L} \quad \text{--- (2)}$$

Equation (2) - Equation (1)

$$3B_0 - B_0 = \frac{\mu_0}{4\pi} 2I \left(\frac{1}{L} - \frac{1}{3L} \right)$$

$$= \frac{\mu_0}{4\pi} \frac{2I}{L} \left(1 - \frac{1}{3} \right)$$

$$= \frac{\mu_0}{4\pi} \frac{2I}{L} \left(\frac{2}{3} \right) = \frac{\mu_0}{4\pi} \frac{2I}{\left(\frac{3L}{2} \right)}$$

KUMAR PHYSICS CLASSES

F 281 BASEMENT 3rd BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI


9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

ii. Over a time interval of 2.0 s, the current in the wire is decreased. The initial magnetic flux through the coil is $5 \times 10^5 \text{ Tm}^2$ and the final magnetic flux through the coil is $1.0 \times 10^5 \text{ Tm}^2$. The coil has a total resistance of 10Ω . Calculate the magnitude of the average current in the coil during the 2.0 s time interval.

$$\begin{aligned} \mathcal{E} &= -N \frac{d\phi}{dt} = -1 \frac{(\phi_f - \phi_i)}{(t_2 - t_1)} = - \frac{(1.0 \times 10^5 - 5 \times 10^5)}{(2 - 0)} \\ &= \frac{4 \times 10^5}{2} = 2 \times 10^5 \text{ Volt}, \quad I = \frac{V}{R} = \frac{2 \times 10^5}{10} = 2 \times 10^4 \text{ A} \end{aligned}$$



The wire is removed and the square coil is positioned so that the coil is directly above and concentric with a round coil of wire connected to a power supply. A part of the square coil is removed and a lightbulb is connected to the coil, as shown above.

(c) During a short time interval, the current in the power supply is constantly increasing. Use physics principles to explain why the lightbulb is lit during the entire time interval.

$B \rightarrow$ is produced by coil, if $I \uparrow$, $B \uparrow$, $\phi \uparrow$

hence more rate of change of flux linked with the coil and hence emf will be induced in the coil, current flows. therefore light bulb is lit during the entire time interval.

Physics Tutor ,Physics Classes In Delhi,Physics Tutor In South Delhi,physics tutor in486

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

+91-9958461445

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

**Online Physics
Classes/Tutor**

AP Physics (2)

Algebra-Based 2022

Paper Solution

**AP,IB DP HL/SL,IGCSE,A-LEVEL,O-
LEVEL,MCAT,ACT,NEET,IIT**

Kumar Physics classes-www.kumarphysicsclasses.com

Physics Tutor ,Physics Classes In Delhi,Physics Tutor In South Delhi,physics tutor in488
Delhi

Physics Tutor ,Physics Classes In Delhi,Physics Tutor In South Delhi,physics tutor in489
Delhi