

PHYSICS

JEE-MAIN (July-
Attempt)

29 July (Shift-1) Paper
Solution

$$r_{\text{com}} = \frac{m_1 r_1 + m_2 r_2 + \dots + m_n r_n}{m_1 + m_2 + \dots + m_n}$$

$$\oint \vec{E} \cdot d\vec{s} = q / \epsilon_0$$

$$R = \frac{\mu^2 \sin 2\theta}{q}$$

$$M = I A$$

$$I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \phi$$

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

IIT JEE PHYSICS PAPER

SOLUTION

29 JULY 2022

MORNING SHIFT

QUESTIONS

BASED ON

SHM,

GAUSS THEOREM, AC,

VERNIER CALLIPERS, SCREW

GAUGE ARE TRICKY

SECTION - A

1. Given below are two statements: One is labelled as Assertion (A) and other is labelled as Reason (R).

Assertion (A): Time period of oscillation of a liquid drop depends on surface tension (S), if density of the liquid is ρ and radius of the drop is r , then $T = K \sqrt{\frac{\rho r^3}{S}}$ is dimensionally correct, where K is dimensionless.

Reason (R): Using dimensional analysis we get R.H.S. having different dimension than that of time period. In the light of above statements, choose the correct answer from the options given below.

(A) Both (A) and (R) are true and (R) is the correct explanation of (A)

(B) Both (A) and (R) are true but (R) is not the correct explanation of (A)

(C) (A) is true but (R) is false

(D) (A) is false but (R) is true

$$T = K \sqrt{\frac{\rho r^3}{S^{3/2}}}$$

$$\text{R.H.S.} = \frac{M^{1/2} L^{-3/2} L^{3/2}}{[M T^{-2}]^{3/4}} = M^{1/8} L^0 T^{3/2}$$

$$\text{R.H.S.} \neq \text{L.H.S.}$$

2. A ball is thrown up vertically with a certain velocity so that, it reaches a maximum height h . Find the ratio of the times in which it is at height $h/3$ while going up and coming down respectively.

(A) $\frac{\sqrt{2}-1}{\sqrt{2}+1}$

(B) $\frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}+\sqrt{2}}$

(C) $\frac{\sqrt{3}-1}{\sqrt{3}+1}$

(D) $\frac{1}{3}$

B

Motion between O & A

$$v^2 = u^2 + 2as$$

$$(0)^2 = u^2 + 2(-g)h$$

$$u = \sqrt{2gh}$$

Consider up motion between O & P.


$$\frac{h}{3} = \sqrt{2gh} t_1 - \frac{1}{2} g t_1^2$$

$$\frac{g t_1^2}{2} - \sqrt{2gh} t_1 + \frac{h}{3} = 0$$

$$t_1^2 - \frac{2}{g} \sqrt{2gh} t_1 + \frac{2h}{3g} = 0$$

$$t_1 = \frac{\left(-\frac{2}{g} \sqrt{2gh}\right) \pm \sqrt{\frac{4}{g^2} (2gh) - 4(1) \left(\frac{2h}{3g}\right)}}{2(1)}$$







$$\begin{aligned} \frac{t_1}{t_2} &= \frac{\sqrt{2gh} + \sqrt{2gh - 4(g/2)(h/3)}}{\sqrt{2gh} - \sqrt{2gh - 4(g/2)(h/3)}} \\ &= \frac{\sqrt{2gh} + \sqrt{\frac{4gh}{3}}}{\sqrt{2gh} - \sqrt{\frac{4gh}{3}}} \\ &= \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}} \end{aligned}$$



3. If $t = \sqrt{x} + 4$, then $\left(\frac{dx}{dt}\right)_{t=4}$ is :
(A) 4 (B) zero (C) 8 (D) 16

$$\begin{aligned}t &= \sqrt{x} + 4 \\ \sqrt{x} &= t - 4 \\ x &= (t - 4)^2 \\ x &= t^2 - 8t + 16 \\ \frac{dx}{dt} &= 2t - 8 \\ \left. \frac{dx}{dt} \right|_{t=4} &= 2 \times 4 - 8 \\ &= 8 - 8 \\ &= 0\end{aligned}$$

4. A smooth circular groove has a smooth vertical wall as shown in figure. A block of mass m moves against the wall with a speed v . Which of the following curve represents the correct relation between the normal reaction on the block by the wall (N) and speed of the block v ?




$$N = \frac{m v^2}{r}$$

$$N \propto v^2$$

PARABOLA

5. A ball is projected with kinetic energy E , at an angle of 60° to the horizontal. The kinetic energy of this ball at the highest point of its flight will become:

- (A) Zero
- (B) $E/2$
- (C) $E/4$
- (D) E



$$E = \frac{1}{2} m u^2$$

At highest point, $v = u \cos 60^\circ$

KE | at top most point

$$= \frac{1}{2} m (u \cos 60^\circ)^2 = \frac{E}{4}$$

6. Two bodies of mass 1 kg and 3 kg have position vectors $i + 2j + k$ and $-3i - 2j + k$ respectively. The magnitude of position vector of centre of mass of this system will be similar to the magnitude of vector:

- ✓ (A) $i + 2j + k$
- (B) $-3i - 2j + k$
- (C) $-2i + 2k$
- (D) $-2i - j + 2k$

$$\begin{aligned}
 r_{com} &= \frac{m_1 \bar{r}_1 + m_2 \bar{r}_2}{m_1 + m_2} = \frac{1(\hat{i} + 2\hat{j} + \hat{k}) + 3(-3\hat{i} - 2\hat{j} + \hat{k})}{1 + 3} \\
 &= \frac{\hat{i} + 2\hat{j} + \hat{k} - 9\hat{i} - 6\hat{j} + 3\hat{k}}{4} \\
 &= \frac{-8\hat{i} - 4\hat{j} + 4\hat{k}}{4} = -2\hat{i} - \hat{j} + \hat{k}.
 \end{aligned}$$

$$\begin{aligned}
 |r_{com}| &= \sqrt{(-2)^2 + (-1)^2 + (1)^2} \\
 &= \sqrt{6} \text{ - similar to option A}
 \end{aligned}$$

7. Given below are two statements : One is labelled as Assertion (A) and other is labelled as Reason (R).

Assertion (A) : Clothes containing oil or grease stains cannot be cleaned by water wash.

Reason (R): Because the angle of contact between the oil/grease and water is obtuse.

In the light of the above statements, choose the correct answer from the option given below.


- ✓ (A) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (B) Both (A) and (R) are true but (R) is not the correct explanation of (A)
- (C) (A) is true but (R) is false
- (D) (A) is false but (R) is true



8. If the length of a wire is made double and radius is halved of its respective values. Then, the Young's

modulus of the material of the wire will:


- ✓ (A) remain same
- (C) become $\frac{1}{8}$ of its initial value
- (B) become 8 time its initial value
- (D) become 4 times its initial value



depends on material
of wire

9. The time period of oscillation of a simple pendulum of length L suspended from the roof of a vehicle, which moves without friction down an inclined plane of inclination α is given by:

- (A) $2\pi\sqrt{L/g \cos\alpha}$ (B) $2\pi\sqrt{L/g \sin\alpha}$ (C) $2\pi\sqrt{L/g}$ (D) $2\pi\sqrt{L/g \tan\alpha}$




$$g_{\text{eff}} = g \cos\alpha$$

$$T = 2\pi \sqrt{\frac{L}{g \cos\alpha}}$$

10. A spherically symmetric charge distribution is considered with charge density varying as

$$\rho(r) = \begin{cases} \rho_0 \left(\frac{3}{4} - \frac{r}{R} \right) & \text{for } r \leq R \\ \text{zero} & \text{for } r > R \end{cases}$$

Where, r ($r < R$) is the distance from the centre O (as shown in figure). The electric field at point P will be



(A) $\frac{\rho_0 r}{4\epsilon_0} \left(\frac{3}{4} - \frac{r}{R} \right)$

(B) $\frac{\rho_0 r}{3\epsilon_0} \left(\frac{3}{4} - \frac{r}{R} \right)$

(C) $\frac{\rho_0 r}{4\epsilon_0} \left(1 - \frac{r}{R} \right)$

(D) $\frac{\rho_0 r}{5\epsilon_0} \left(1 - \frac{r}{R} \right)$

sol. C


Apply Gauss theorem

$$\int \vec{E} \cdot d\vec{L} = \frac{Q}{\epsilon_0}$$

$$E \cdot 4\pi r^2 = \int_0^r \rho_0 \left(\frac{3}{4} - \frac{r}{R} \right) \cdot 4\pi r^2 dr$$

$$E r^2 = \frac{\rho_0 r^3}{4\epsilon_0} \left\{ 1 - \frac{r}{R} \right\}$$

$$E = \frac{\rho_0 r}{4\epsilon_0} \left\{ 1 - \frac{r}{R} \right\}$$



11. Given below are two statements.

Statement I : Electric potential is constant within and at the surface of each conductor. Statement II : Electric field just outside a charged conductor is perpendicular to the surface of the conductor at every point.

In the light of the above statements choose the most appropriate answer from the options given below.

- (A) Both statement I and statement II are correct
 (B) Both statement I and statement II are incorrect
 (C) Statement I is correct but statement II is incorrect
 (D) Statement I is incorrect but statement II is correct

STATEMENT-1

↳ Body of the conductor acts as a equipotential surface

STATEMENT-2

↳ since conductor acts as equipotential surface then tangential component of electric field should be zero hence electric field is \perp to the surface.

12. Two metallic wires of identical dimensions are connected in series. If σ_1 and σ_2 are the conductivities of these wires respectively, the effective conductivity of the combination is :

(A) $\frac{\sigma_1 \sigma_2}{\sigma_1 + \sigma_2}$

(B) $\frac{2\sigma_1 \sigma_2}{\sigma_1 + \sigma_2}$

(C) $\frac{\sigma_1 + \sigma_2}{2\sigma_1 \sigma_2}$

(D) $\frac{\sigma_1 + \sigma_2}{\sigma_1 \sigma_2}$



$$R_{eq} = R_1 + R_2$$

$$\frac{2l}{\sigma_{eq} A} = \frac{l}{\sigma_1 A} + \frac{l}{\sigma_2 A}$$

$$\sigma_{eq} = \frac{2\sigma_1 \sigma_2}{\sigma_1 + \sigma_2}$$

13. An alternating emf $E = 440 \sin 100\pi t$ is applied to a circuit containing an inductance of $\frac{\sqrt{2}}{\pi}$ H. If an a.c. ammeter is connected in the circuit, its reading will be :

(A) 4.4 A (B) 1.55 A (C) 2.2 A (D) 3.11 A

Sol. C

$$E = 440 \sin 100\pi t \quad L = \frac{\sqrt{2}}{\pi}$$

$$X_L = \omega L = 100 \cancel{\pi} \frac{\sqrt{2}}{\cancel{\pi}}$$
$$= 100\sqrt{2} \text{ ohm}$$

$$\text{Peak current } I_0 = \frac{E_0}{X_L} = \frac{400}{100\sqrt{2}}$$
$$= 2.2\sqrt{2} \text{ A}$$

Ammeter reads R.M.S value therefore reading will be I_{rms}

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}} = 2.2 \text{ Amp}$$

14. A coil of inductance 1 H and resistance 10 Ω is connected to a battery of 6 V. Determine approximately :

(a) The time elapsed before the current acquires half of its steady-state value.

(b) The energy stored in the magnetic field associated with the coil at an instant 15 ms after the circuit is

switched on. (Given $\ln 2 = 0.693$, $e^{-3/2} = 0.25$)

(A) $t = 10$ ms; $U = 2$ mJ (B) $t = 10$ ms; $U = 1$ mJ

(C) $t = 7$ ms; $U = 1$ mJ (D) $t = 7$ ms; $U = 2$ mJ




$$i(15 \text{ ms}) = \frac{E}{R} (1 - e^{-15/10})$$

$$i = \frac{6}{100} (1 - \frac{1}{4}) = \frac{3}{4} \times \frac{6}{100}$$

$$U = \frac{1}{2} L I^2$$

$$U = 1 \text{ mJ}$$



$$I = \frac{E}{R} (1 - e^{-t/\tau})$$

$$\frac{E}{2R} = \frac{E}{R} (1 - e^{-t/\tau})$$

$$\frac{1}{2} = 1 - e^{-t/\tau} \Rightarrow e^{-t/\tau} = 1 - \frac{1}{2}$$

$$e^{t/\tau} = 2 \Rightarrow \frac{t}{\tau} \log e = \log e^2$$

$$t = \tau \log e^2$$

$$= \frac{1}{100} \times 0.693$$

$$= 0.00693$$

$$= 7 \text{ ms}$$

15. Match List - I with List - II :

List - I

- (a) UV rays
- (b) X-rays
- (c) Microwave
- (d) Infrared wave

List - II

- (i) Diagnostic tool in medicine
- (ii) Water purification
- (iii) Communication, Radar
- (iv) Improving visibility in foggy days

Choose the correct answer from the options given below :

- (A) (a)-(iii), b-(ii), (c) - (i), d-(iv)
- (C) (a)-(ii), b-(iv), (c) - (iii), d-(i)

- (B) (a)-(ii), b-(i), (c) - (iii), d-(iv)
- (D) (a)-(iii), b-(i), (c) - (ii), d-(iv)

UV - Rays

↓
Water purification

Xray → diagnosing
fracture

Microwave

↳ Radar and mobile
communication

Infrared → less scattering
↓
used in foggy days

16. The kinetic energy of emitted electron is E when the light incident on the metal has wavelength λ . To double the kinetic energy, the incident light must have wavelength:

(A) $\frac{hc}{E\lambda - hc}$

(B) $\frac{hc\lambda}{E\lambda + hc}$

(C) $\frac{h\lambda}{E\lambda + hc}$

(D) $\frac{hc\lambda}{E\lambda - hc}$

$$E = \frac{hc}{\lambda} - \phi$$

$$2E = \frac{hc}{\lambda'} - \phi$$

subtract

$$E = hc \left(\frac{1}{\lambda'} - \frac{1}{\lambda} \right)$$

$$\lambda' = \frac{hc\lambda}{E\lambda + hc}$$

17. Find the ratio of energies of photons produced due to transition of an electron of hydrogen atom from its (i) second permitted energy level to the first level, and (ii) the highest permitted energy level to the first permitted level.

✓ (A) 3:4

(B) 4:3

(C) 1:4

(D) 4:1

$$E_n = - \frac{13.6}{n^2} \text{ eV}$$

$$\frac{E_2 - E_1}{E_\infty - E_1} = \frac{13.6 \left(1 - \frac{1}{4}\right)}{13.6}$$
$$= \frac{3}{4}$$

18. Find the modulation index of an AM wave having 8 V variation where maximum amplitude of the AM wave is 9V.

✓ (A) 0.8

(B) 0.5

(C) 0.2

(D) 0.1

Modulation index

$$\mu = \frac{A_m}{A_c}$$

$$2 A_m = 8$$

$$A_m = 4$$

$$A_c = 5$$

$$\mu = \frac{4}{5} = 0.8$$

19. A travelling microscope has 20 divisions per cm on the main scale while its vernier scale has total 50 divisions and 25 vernier scale divisions are equal to 24 main scale divisions, what is the least count of the travelling microscope?

(A) 0.001 cm (B) 0.002 mm (C) 0.002 cm (D) 0.005 cm

$$1 \text{ M.S.D} = \frac{1}{20} \text{ cm}$$

$$1 \text{ V.S.D} = \frac{24}{25} \text{ M.S.D}$$


$$= \frac{24}{25} \times \frac{1}{20} \text{ cm.}$$

$$\text{Least count} = \frac{1}{20} \left(1 - \frac{24}{25} \right) \text{ cm}$$

$$= \frac{1}{20} \times \frac{1}{25} = \frac{1}{500} \text{ cm}$$

$$= 0.002 \text{ cm}$$

20. In an experiment to find out the diameter of wire using screw gauge, the following observations were noted :



- (a) Screw moves 0.5 mm on main scale in one complete rotation
(b) Total divisions on circular scale = 50
(c) Main scale reading is 2.5 mm
(d) 45th division of circular scale is in the pitch line
(e) Instrument has 0.03 mm negative error
Then the diameter of wire is :

(A) 2.92 mm (B) 2.54 mm (C) 2.98 mm (D) 3.45 mm

$$MSR = 2.5 \text{ mm}$$

$$CSR = 45 \times \frac{0.5}{50} \text{ mm}$$

$$= 0.45 \text{ mm}$$

Diameter reading

$$= MSR + CSR - \text{Zero error}$$

$$= 2.5 + 0.45 - (-0.03)$$

$$= 2.98 \text{ mm}$$

SECTION - B

21. An object is projected in the air with initial velocity u at an angle θ . The projectile motion is such that the horizontal range R , is maximum. Another object is projected in the air with a horizontal range half of the range of first object. The initial velocity remains same in both the case. The value of the angle of projection, at which the second object is projected, will be _____ degree.

Sol. 15°

$$R_{\max} = \frac{u^2 \sin 2(45^\circ)}{g} = \frac{u^2}{g}$$

$$\frac{R}{2} \Rightarrow \frac{u^2}{2g} = \frac{u^2 \sin 2\theta}{g}$$

$$\sin 2\theta = \frac{1}{2}$$

$$2\theta = 30^\circ, 150^\circ$$

$$\theta = 15^\circ, 75^\circ$$

22. If the acceleration due to gravity experienced by a point mass at a height h above the surface of earth is same as that of the acceleration due to gravity at a depth d ($h \ll R_e$) from the earth surface. The value of α will be 2.
(use $R_e = 6400 \text{ km}$)

$$g \left(1 - \frac{2h}{R}\right) = g \left(1 - \frac{d}{R}\right)$$

$$\frac{2h}{R} = \frac{d}{R}$$

$$d = 2h = \alpha h.$$

$$\underline{\alpha = 2}$$

23. The pressure P and density d of diatomic gas $\gamma = \frac{7}{5}$ changes suddenly to $P_2 > P_1$ and d_2 respectively during an adiabatic process. The temperature of the gas increases and becomes 4 times of its initial temperature. (given $\frac{d_2}{d_1} = 32$)

$$P V^\gamma = \text{constant}$$

$$P \left(\frac{m}{d} \right)^\gamma = \text{constant}$$

$$\frac{P}{d^\gamma} = \text{constant}$$

$$\frac{P_1}{P_2} = \left(\frac{d_1}{d_2} \right)^\gamma = \left(\frac{1}{32} \right)^{\frac{7}{5}} = \frac{1}{128}$$

$$\frac{d_2}{d_1} = 32$$

$$\frac{T_1}{T_2} = \frac{P_1 V_1}{P_2 V_2} = \frac{32}{128} = \frac{1}{4}$$




24. One mole of a monoatomic gas is mixed with three moles of a diatomic gas. The molecular specific heat of mixture at constant volume is $\frac{\alpha^2}{4}$ RJ/mol K; then the value of α will be

Sol. ³ ----- (Assume that the given diatomic gas has no vibrational mode).



$$\begin{aligned}C_{V_{\text{mix}}} &= \frac{n_1 C_{V_1} + n_2 C_{V_2}}{n_1 + n_2} \\&= \frac{1 \left(\frac{3R}{2} \right) + 3 \left(\frac{5R}{2} \right)}{1 + 3} \\&= \frac{9R}{4} = \frac{\alpha^2}{4} R.\end{aligned}$$

$$\alpha = 3$$

25. The current I flowing through the given circuit will be 2 A.



A, L } same point
 K, B } same point



$$I = \frac{6}{3} = 2 \text{ A}$$

26. A closely wound circular coil of radius 5 cm produces a magnetic field of 3768×10^{-4} T at its center. The current through the coil is 3 A.

[Given, number of turns in the coil is 100 and $\pi = 3.14$]

$$B_{\text{centre}} = \frac{N \mu_0 I}{2R}$$

$$37.68 \times 10^{-4} = \frac{100 \times 4\pi \times 10^{-7} \times I}{2 \times 5 \times 10^{-2}}$$

$$I = 3 \text{ Amp}$$

27. Two light beams of intensities $4I$ and $9I$ interfere on a screen. The phase difference between these beams on the screen at point A is zero and at point B is π . The difference of resultant intensities, at the point A and B, will be 24 I.

$$I_{\text{net}} = I_1 + I_2 + 2\sqrt{I_1}\sqrt{I_2}\cos\phi$$

$$I_{\text{max}}$$

$$\phi = 0$$

$$I_{\text{max}} = (\sqrt{I_1} + \sqrt{I_2})^2$$

$$= (\sqrt{9I} + \sqrt{4I})^2$$

$$= 25I$$

$$I_{\text{min}}$$

$$\phi = \pi$$

$$I_{\text{min}} = (\sqrt{I_1} - \sqrt{I_2})^2$$

$$= (\sqrt{9I} - \sqrt{4I})^2$$

$$= I$$

$$I_{\text{max}} - I_{\text{min}} = 25I - I$$

$$= 24I$$

28. A wire of length 314 cm carrying current of 14 A is bent to form a circle. The magnetic moment of the coil is 11 A-m². [Given $\pi = 3.14$]

$$2\pi R = \frac{314}{100}$$

$$R = 0.5 \text{ m}$$

$$\text{Magnetic moment} = IA$$


$$= 14 \times \pi R^2$$

$$= 14 \times 3.14 \times \frac{1}{4}$$

$$= 10.99 \approx 11$$

29. The X-Y plane be taken as the boundary between two transparent media M_1 and M_2 . M_1 in $Z \geq 0$ has a refractive index of $\sqrt{2}$ and M_2 with $Z < 0$ has a refractive index of $\sqrt{3}$. A ray of light

travelling in M_1 along the direction given by the vector $P = 4\sqrt{3}\hat{i} - 3\sqrt{3}\hat{j} - 5\hat{k}$, is incident on the plane of separation. The value of difference between the angle of incident in M_1 and the angle of refraction in M_2 will be 15 degree.



$$\mu_1 \sin i = \mu_2 \sin r$$

$$i = \cos^{-1} \left(\frac{AZ}{A} \right) = \cos^{-1} \left(\frac{5}{\sqrt{(4\sqrt{3})^2 + (3\sqrt{3})^2 + 5^2}} \right)$$

$$i = \cos^{-1} \left(\frac{5}{10} \right)$$

$$i = 60^\circ$$

$$\sqrt{2} \sin 60 = \sqrt{3} \sin r$$

$$\sin r = \frac{1}{\sqrt{2}} \quad r = 45^\circ$$

$$Li - Lr = 60 - 45 = 15$$

30)- If the potential barrier across a p-n junction is 0.6 V. Then the electric field intensity, in the depletion region having the width of 6×10^{-6} m, will be _____ $\times 10^5$ N/C.

Sol. 1

$$E = \frac{V}{d} = \frac{\text{potential barrier across junction}}{\text{width of depletion layer}}$$
$$= \frac{0.6}{6 \times 10^{-6}} = 1 \times 10^5 \text{ V/m.}$$

KUMAR PHYSICS CLASSES

E 281 BASEMENT M BLOCK MAIN ROAD GREATER KAILASH 2 NEW DELHI

9958461445, 01141032244

www.kumarphysicsclasses.com

www.kumarneetphysicsclasses.com

IIT JEE PHYSICS PAPER

SOLUTION

29 JULY 2022

MORNING SHIFT

QUESTIONS

BASED ON

SHM,

GAUSS THEOREM, AC,

VERNIER CALLIPERS, SCREW

GAUGE ARE TRICKY

Physics Tutor ,Physics Classes In Delhi,Physics Tutor In South Delhi,physics tutor in479
Delhi

Physics Tutor ,Physics Classes In Delhi,Physics Tutor In South Delhi,physics tutor in480
Delhi