LEVEL 2—Anxiety—Parent/Guardian of Child Age 6-17^{*}

*Adapted from PROMIS Emotional Distress—Anxiety—Parent Item Bank

Child's Name:	Age:	Sex: 🛛 Male 🖵 Female	Date:
---------------	------	----------------------	-------

What is your relationship with the child receiving care?_____

<u>Instructions to parent/guardian</u>: On the DSM-5 Level 1 cross-cutting questionnaire that you just completed, you indicated that *during the past 2 weeks* your child receiving care has been bothered by "feeling nervous, anxious, or scared", "not being able to stop worrying", and/or "couldn't do things he/she wanted to or should have done because they made him/her feel nervous" at a mild or greater level of severity. The questions below ask about these feelings in more detail and especially how often your child receiving care has been bothered by a list of symptoms <u>during the past</u> <u>7 days</u>. Please respond to each item by marking (\checkmark or x) one box per row.

							Clinician Use
In the past SEVEN (7) DAYS, my child said that he/she					ltem		
		Never	Almost Never	Sometimes	Often	Almost Always	Score
1.	Felt like something awful might happen.	1	2	3	4	5	
2.	Felt nervous.	1	2	3	• 4	D 5	
3.	Felt scared.	1	2	3	• 4	5	
4.	Felt worried.	1	2	3	4	5	
5.	Worried about what could happen to him/her.	1	2	3	4	D 5	
6.	Worried when he/she went to bed at night.	1	2	3	4	5	
7.	Got scared really easy.	1	2	3	4	5	
8.	Was afraid of going to school.	1	2	3	4	5	
9	Worried when he/she was at home.	1	2	3	4	5	
10.	Worried when he/she was away from home.	1	2	3	4	5	
Total/Partial Raw Score:							
Prorated Total Raw Score:							
T-Score:							

 ${\small @2008-2012} \ {\it PROMIS} \ {\it Health} \ {\it Organization} \ ({\it PHO}) \ {\it and} \ {\it PROMIS} \ {\it Cooperative} \ {\it Group}.$

This material can be reproduced without permission by clinicians for clinical use with their patients.

Any other use, including electronic use, requires written permission of the PHO.

Instructions to Clinicians

The DSM-5 Level 2—Anxiety—Parent/Guardian of Child Age 6–17 measure is the 10-item PROMIS Anxiety Form that assesses the pure domain of anxiety in children and adolescents. The measure is completed by the parent or guardian about the child prior to a visit with the clinician. Each item asks the parent or guardian to rate the severity of his or her child's anxiety during the past 7 days.

Scoring and Interpretation

Each item on the measure is rated on a 5-point scale (1=almost never; 2=rarely; 3=sometimes; 4=often; and 5=almost always) with a range in score from 10 to 50 with higher scores indicating greater severity of anxiety. The clinician is asked to review the score on each item on the measure during the clinical interview and indicate the raw score for each item in the section provided for "Clinician Use." The raw scores on the 10 items should be summed to obtain a total raw score. Next, the T-score table should be used to identify the T-score associated with the total raw score and the information entered in the T-score row on the measure.

Score	T-Score	SE	Score	T-Score	SE
10	34.4	5.6	31	66.9	2.9
11	39.4	4.3	32	67.9	2.9
12	42.3	3.9	33	68.9	2.9
13	44.6	3.5	34	70	2.9
14	46.5	3.2	35	71	2.8
15	48.2	3.1	36	72	2.8
16	49.7	2.9	37	73	2.8
17	51.1	2.9	38	73.9	2.8
18	52.4	2.8	39	74.9	2.8
19	53.6	2.8	40	75.9	2.8
20	54.8	2.7	41	76.9	2.8
21	55.9	2.7	42	77.9	2.8
22	57.1	2.7	43	79	2.8
23	58.2	2.8	44	80	2.8
24	59.3	2.8	45	81.2	2.9
25	60.4	2.8	46	82.4	3
26	61.5	2.8	47	83.6	3.1
27	62.6	2.8	48	85	3.2
28	63.7	2.9	49	86.6	3.2
29	64.8	2.9	50	88.8	3.3
30	65.8	2.9			

Note: This look-up table works only if <u>all items</u> on the form are answered. If 75% or more of the questions have been answered, you are asked to prorate the raw score and then look up the conversion to T-Score. The formula to prorate the partial raw score to Total Raw Score is:

(Raw sum x number of items on the short form) Number of items that were actually answered

If the result is a fraction, round to the nearest whole number. For example, if 9 of 10 items were answered and the sum of those 9 responses was 28, the prorated raw score would be 28 X 10/9 = 31, after rounding. The T-score in this example would be 66.9.

The T-scores are interpreted as follows:

Less than 55	= None to slight
55.0—59.9	= Mild
60.0—69.9	= Moderate
70 and over	= Severe

If more than 25% of the total items (in this case more than 2) are missing a response, the scores should not be used. Therefore, the parent or guardian should be encouraged to complete all of the items on the measure.

©2008-2012 PROMIS Health Organization (PHO) and PROMIS Cooperative Group.

Frequency of Use

To track change in the severity of the child's anxiety over time, the measure may be completed at regular intervals as clinically indicated, depending on the stability of the child's symptoms and treatment status. For consistency, it is preferred that completion of the measures at follow-up appointments is by the same parent or guardian. Consistently high scores on a particular domain may indicate significant and problematic areas for the child that might warrant further assessment, treatment, and follow-up. Your clinical judgment should guide your decision.