

8760JANE ST. SUITE B05B CONCORD, ON L4K0E8 CANADA

(905) 237-4623 (905) 237-5098 reception@tipsmed.com www.tipsmed.com

Date:	
RE:	
DOB:	
Dear Doctor,	
I am currently treating the above patient and injection.	consider performing a fluoroscopically- guided
The patient has informed me of being on an anticoagulant. Please indicate if there are any contraindications to withholding this medication prior to the procedure to minimize the possibility of bleeding or hematoma formation.	
- Yes, it is ok to hold the anticoagulant as per table below. - No, do NOT hold the anticoagulant. Due to:	
Physician Signature:	
☐ Apixaban(Eliquis) [3d] ☐ Rivaroxaban(Xarelto)[3d] ☐ Dabigatran(Pradaxa) [5d] ☐ Warfarin(Coumadin) [~3d] ☐ Ticlopidine(Ticlid) [7d] ☐ Edoxaban (Lixiana) [3d]** **Per Updates to the ASRA	 □ ASA/Dypiridamol (Aggrenox)[7d] □ Clopidogrel (Plavix) [7d] □ Prasugrel (Effient) [7d] □ Ticagrelor (Brilinta) [7d] □ all NSAIDs(except Piroxicam)[7d] □ Absciximab (Reopro) [2d] Guidelines for Interventional Pain Procedures- ASRA Aug2019
Regional Anesthesia in the Patient Receiving Antithrombotic or Thrombolytic Therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Fourth Edition) April 2018. doi: 10.1097/AAP.00000000000000763	
You can fax this form back to (905) 237-5098	

For further inquiries, please contact (905) 237-4623