

AUGUST 2021, VOL:1, NO:2

# prithivya

AN OFFICIAL NEWSLETTER OF  
WCB RESEARCH FOUNDATION &  
WCB RESEARCH LAB



© Sidharth Amin

## Read about

Leucism in Indian Flying Fox  
reported from Gujarat






## Editorial Board

<b>Editor in Chief</b>	: <b>Dr. Sagarika Damle</b> <i>Head, Department of Life Sciences, KC college, HNSC University, Mumbai</i>
<b>Associate Editor</b>	: <b>Dr. Nishith Dharaiya</b> <i>Associate Professor, Department of Life Sciences, HNG University, Patan</i>
	: <b>Dr. Jagruti Rathod</b> <i>Director, Center for Nature &amp; Ecosystem Wellbeing (CNEW), Vadodara</i>
<b>Editors</b>	: <b>Prof. V. C. Soni</b> <i>Retd. Professor, Department of Biosciences, Saurashtra University, Rajkot</i>
	: <b>Dr. B.K. Jain</b> <i>Rtd. Principal, MG Science Institute, Ahmedabad</i>
	: <b>Prof. Hitesh Solanki</b> <i>Professor, Department of Botany, Gujarat University, Ahmedabad</i>
	: <b>Dr. Nuria Selva</b> <i>Associate Professor, Institute of Nature and Science Polish Academy of Science, Krakow, Poland</i>
	: <b>Mr. Thomas Sharp</b> <i>Director of Conservation and Research, Wildlife SOS, USA</i>
	: <b>Dr. Krzysztof Klimaszewski</b> <i>Assistant Professor, Department of Zoology, Warsaw University of Life Sciences, Warsaw, Poland</i>
	: <b>Dr. Zornitza Aguilar</b> <i>Scientist, SNAP Ecuador</i>
	: <b>Dr. Darshan Marjadi</b> <i>Assistant Professor, Department of Biotechnology, Sarvajanik University, Surat</i>
	: <b>Dr. Rajesh Senma</b> <i>Assistant Professor, Government Science College, Vankal</i>
	: <b>Dr. Archana Gajjar</b> <i>Assistant Professor, Bahauddin Science College, Junagadh</i>
	: <b>Dr. Nisha Singh</b> <i>Scientist, Bombay Natural History Society, Mumbai</i>
	: <b>Dr. Sumesh Dudhani</b> <i>Project Consultant, Natural heritage, INTACH, Delhi</i>
<b>Editorial Assistant</b>	: <b>Ms. Jahnvi Patel   Mr. Pratik Desai   Ms. Rinku Bhati</b> <i>WCB Research Foundation, Patan</i>
<b>Language Editor</b>	: <b>Dr. Ramaa Gaury Mandava</b> <i>WCB Research Foundation, Patan</i>

Contact details: +918320519905 | Email: [prithivya.wcb@gmail.com](mailto:prithivya.wcb@gmail.com)



# TABLE OF CONTENTS

## Prithivya

A Quarterly Newsletter of  
WCB Research Foundation and

Volume 1 | Issue 2  
(August 2021)



**Chief Editor:**  
Dr. Sagarika Damle

**Associate Editors:**  
Dr. Nishith Dharaiya  
Dr. Jagruti Rathod

**Editorial Assistant:**  
Ms. Rinku Bhati  
Jahnvi Patel  
Pratik Desai

**Design & Creation:**  
Mr. Sachin and Dr. Nishith Dharaiya

**Cover page:**  
Indian Flying Fox  
(*Pteropus giganteus*)  
By Sidharth Amin

© Printed & Published by  
WCB Research Foundation, India  
[www.wcbresearch.in](http://www.wcbresearch.in)

### Field observations

- 1-3 *Range extinction of Asian pied starling (*Gracupica contra*) in Gujarat state, Western India*  
**Chaudhary and Desai**
- 4-11 *Observation on nesting pattern of different bird species around university road, Rajkot*  
**Dave D**
- 12-17 *Record of leucism in Indian flying fox (*Pteropus giganteus*) from South Gujarat, India*  
**Amin et al.**
- 18-21 *Notes on behavioural observation of some water birds during the annual solar eclipse*  
**Thakkar and Gohil**

### Short Articles

- 22-23 *Restoration of Earth with bio based polymer*  
**Marjadi D**
- 24-27 *Single-used surgical mask: human life saver or biodiversity destroyer*  
**Patel and Patel**
- 28-36 *Does the contemplation of forest field staff about wildlife differ than a common man?*  
**Pithva and Dharaiya**


### Research Papers

- 37-43 *Qualitative phytochemical analysis of some medicinal plants (leaves) from the surrounding areas of Talod taluka, North Gujarat, India*  
**Panchal and Jha**

### Assorted news

- Final nominees of M I Patel Award 2021*
- Announcements*
- Winners of Mangrove day poster competition*

# EDITORIAL NOTE



पर्यावरणनाशेन विस्मेत् विस्तो भवेत्, मानवो मानवो भूत्वा कुर्यात् प्रकृति रक्षणम् ।

The word Sustainable livelihood explains the essence of maintaining the values of the assets and enhancing the capabilities of the system, for the current as well as future scenario, while conserving and respecting the natural resources. The second issue of the newsletter 'Prithivya' attempts to draw the attention of the readers to various aspects of sustainability through traditional conservation to high-tech environmental monitoring practices. It offers a wide range of articles ranging from Tribal artifacts and life style to Earth restoration, contributed by young researchers. From endemic species to sacred sites, all natural habitats inherit conservation values. Those biological, ecological, social or cultural values of outstanding significance are known as 'High Conservation Values' or HCVs and an observer's role is pivotal in keeping the meticulous records that help in assigning these HVC's. The Stewardship of Biodiversity Conservation has been ably carried forward by the young brigade of learners, who belong to diverse disciplines but converge towards their common goal of protecting the Mother earth. The recent disasters of untimely, uncontrollable rains and floods, landslides and raging forest fires at global levels, point towards an undesirable climatic shift in earth's environment and the human species is racing against time to be disaster ready. The 'Code Red for Humanity' flashed in the latest U.S.Climate Change report demands for doubling our efforts in ecological restoration regimes. It would happen only and only when the old conservation practices and new technological insights work in tandem to bring about the synergistic effect. 'Prithivya' is the right platform for the caregivers of environment to showcase their ideas and outcomes. We thank all the contributing authors of this 2nd issue and appeal to each reader of this newsletter to participate in some activity that will make a positive difference in sustainable livelihood of the species around us. Let's come together to build a greener tomorrow...




**Dr. Sagarika Damle**

*Chief Editor*

*Prithivya*

*sagarika.damle@kccollege.edu.in*



## DIRECTOR'S COLUMN

Wildlife conservation has captivated great attention from all the sectors of the community. Stakeholders including scientists, policymakers, local and global organizations and citizens have been putting relentless efforts in conserving the wildlife and the wilderness on the earth. This emphasizes on need for adopting foresighted approach and using scientific measures for producing desired results. However, all we need is to adopt a scientific approach in such conservation activities so that it retains for longer period of time and give the desired results. The key to wildlife conservation is target oriented awareness and appreciation of the relationship between wildlife conservation policies and practices. In order to build such relationship, it is crucial that the correct scientific information about the wildlife reaches ground level. Wildlife researchers should be recognised as an important disseminator of science-based information, with education in schools and universities. Along with that, the media should bolster the social responsibility for communicating accurate and reliable conservation information.

At WCB Research Foundation, we strive to bring different stakeholders and media on a single platform to spread the science-based information about wildlife and encourage the community to appreciate wildlife. Our community based conservation approach has now become our mission of “Adding science to the Conservation”.

The foundation will now complete one year! And during this first year of establishment, we initiated several activities through which the scientific information can reach to every single or all sectors of the society. We have started the sloth bear conservation education program in the central Gujarat, where people and sloth bear co-exists and the territorial forest patches connects two important protected areas. We have transformed the scientific information into the fun learning materials such as animation film, documentary videos, stories, comics and activity book. We distribute these products among the school children and the local community and we expect that this will enhance the tolerance and appreciation of locals towards the sloth bears. In this difficult time of the pandemic, we have organized several webinars, online workshops and spread the message on wildlife conservation through social media. In order to provide the platform to wildlife researchers and students, we initiated WCBinars allowing students to present their research and the same we broadcast on our YouTube channel.

WCB is now connected with several national and international organizations through signing MoU and I am sure this will enhance our capacity and network to reach out more people and help wildlife conservation. We are always open to collaborate with organizations and institutions working in the area of mutual interest and I welcome you all to join us, work with us and collaborate with us to serve for Mother Nature and conserving her wilderness.



**Nishith Dharaiya, PhD**

*Director of Research*

*WCB Research Foundation*

*nadharaiya@gmail.com*



## Range Extension of Asian Pied Starling (*Gracupica contra*) in Gujarat state, Western India

Chaudhary Sarthak V.<sup>1</sup> and Desai Pratik G.<sup>2\*</sup>

<sup>1</sup>M. N. College, Visnagar (Gujarat) India.

<sup>2</sup>WCB Research Lab, HNG University, Patan (Gujarat) India.

\*E-mail: [pratikdesai825@gmail.com](mailto:pratikdesai825@gmail.com)

India has more than 1300 bird species, which is about 12.5% of world's total avifauna (Grimmett et al. 2011). Among them, a total of 576 bird species have been reported from Gujarat state (Ganpule, 2016). Asian pied starling (*Gracupica contra*) commonly found in foothills of south Asia until elevation of 700m. It is also commonly found near water bodies of human dominated areas, grasslands and farmland (Ali & Ripley, 1983). Ali & Ripley (1983) recorded three subspecies of Pied starling in India, whereas Grimmett et al. (2005) mentioned only two subspecies of Asian pied starling in India; viz. *G. superciliaris*, observed around Manipur only while, *G. contra* is widely observed in North, central and Eastern parts of India (Grimmett et al. 2005). Monga & Naoroji (1983) mentioned sighting of pied starling in Jamnagar and south Gujarat as rare/vagrant. Presence and breeding of *G. Contra* was also reported from Rajkot and Dahod districts of Gujarat consequently (Rasmussen & Anderton, 2005; Ganpule, 2016) this species was introduced in Mumbai of Maharashtra state, India (Rasmussen & Anderton, 2005).



*Asian pied starling (G. contra) in Vadnagar town of northern part of Gujarat (N 23.7896, E 72.6291)*

During field visit of Vadnagar, Mehsana district in northern part of Gujarat (February 28, 2021), we have observed bird with white on the black wings with white ventral mark and orange bill at 07:30 hrs. After validation through reference book (Grewal, 2016) we have identified it as Asian pied starling (*Gracupica contra*). We have found this individual near the water body of human habitation (N 23.7896, E 72.6291). The individual was



observed for 10 minutes and its behaviour was recorded. Bird was feeding on some unknown excreta material while foraging with other bird species viz., Glossy ibis (*Plegadis falcinellus*), Streak-throated swallow (*Petrochelidon luvicola*). Another individual of the same species pair was observed on April 5, 2021, 16:13 hrs at Kheralu town, approximately 11 km far from previous observation (N 23.8831 E 72.6109) where the bird was sitting on electric wire above the agriculture farm.



*Asian pied starling (G. contra) perching on electric wire above the agriculture farm (N 23.8831 E 72.6109)*



*Asian pied starling (G. contra) feeding on some unidentified excreta*

Asian pied startling was recorded for the first time the semi-arid region of Gujarat, as we did not found any record of this species in this area. This observation reflects the possible range expansion of species. Both of our sightings were around agriculture and human dominated areas indicating the generalist niche and adaptability of this species in human dominated landscape.



## References

- Ali, S., & Ripley, S.D., 1986. Handbook of the bird of India and Pakistan (Vol. 5- Lark to Grey Hypocolius). Oxford University Press.
- Grimmett, R., Inskipp, C., & Inskipp, T., (2005). Birds of Indian Subcontinent. Oxford university press. Pp 402-403.
- Grewal, B., 2016. A Pictorial Field Guide to Birds of India: Pakistan, Nepal, Bhutan, Sri Lanka and Bangladesh. India: Om Books International.
- Monga, S. G., & Naoroji, R. K., 1983. Birds of the Rajpipla forests – SouthGujarat. With notes on nests found and breeding recorded and some new observations. *JBNHS* 80 (3): 575-612
- Rasmussen, P. C., & Anderton, J. C., 2005. Birds of South Asia: The Ripley guide. Washington DC & Barcelona: Smithsonian Institute & Lynx Edition.
- Ganpule, P., 2016. The birds of Gujarat: Status and distribution. *Flamingo*, 8(3), 2\_40.

---

### **Suggested citation:**

Chaudhary SV and Desai PG (2021). Range Extension of Asian Pied Starling (*Gracupica contra*) in Gujarat state, Western India. *Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab. Vol 1(2) 1-3.*







## Observation on nesting pattern of different bird species around University road, Rajkot

**Dave Deepali**

“Anjaney”, Sanjay vatika 8, Neel city club campus, Near Saurashtra University, Rajkot-3640005 (Gujarat), India.

\*E-mail: [deepalidave955@gmail.com](mailto:deepalidave955@gmail.com)

The dense vegetation near Saurashtra university circle provided a preferable avifaunal habitat (Trivedi & Vaghela, 2020). This helped in the study of Avifaunal nesting sites in the urban area of Rajkot (22°17'7.37"N, 70°44'53.88"E, 22°17'10.26"N 70°44'44.03"E, 22°17'2.38"N 70°44'45.14"E, 22°16'58.27"N 70°44'52.31"E), Gujarat. A 2 hours study was conducted while exploring the area 2km away from my residence in the morning for a week in May 2021. Up to, 33 Avifaunal species and seven nesting sites were reported. The nesting sites reported during the survey were of Black Drongo (*Dicrurus macrocercus*), Eurasian collared Dove (*Streptopelia decaocto*), Indian Thick knee (*Burhinus indicus*), Shikra (*Accipiter badius*), Rufous Treepie (*Dendrocitta vagabunda*), Red wattled Lapwing (*Vanellus indicus*), and Cattle egret (*Bubulcus ibis*).



**Figure 1. Detailed Map of Study area**



Point 1 from the above figure indicates my residence, where fieldwork was started. Based on vegetation and human disturbance, the study area was divided into two parts: Sub-urban Area and Rural Area. Point 7 and 8 indicates plantations done by the Saurashtra University. The line transect method was used for the field survey. Images were captured with the help of Canon 1200D, 55-250 mm lens.



Figure 2(a)



Figure 2(b)



Figure 2(c)



Figure 2(d)

**Figure 2: Different photographs of saurashtra university plantation area (Study Area)**

### **Black Drongo (*Dicrurus macrocercus*):**

The nesting season for Black Drongo is April to August, with 3-4 eggs as clutch size and 14-15 days incubation period. The nest of Black drongo was observed at Point 8 (22°17'7.37"N 70°44'53.88"E) in the plantation site of Saurashtra University. It is about 9 to 10 feet high from ground level and built with twigs, rootlets and threads. Drongo is known to be a protective parent and its mobbing behaviour towards Greater Coucal (*Centropus sinensis*), Rose-ringed Parakeet (*Alexandrinus krameri*) and Asian Koel (*Eudynamis scolopacea*) were



observed during the survey (Nijman, 2004). Interspecies interaction between Eurasian Collared-Dove and Black Drongo was observed during the survey and in previous studies (Jahan et al., 2018) as well, where two Eurasian collared Dove made nest near the nest of Black Drongo. Usually, birds like Dove do not cause any harm to Drongo. Feeding gills for both species is different as well (Ali, 2017, Grimmett et al., 2011). Therefore, to protect their eggs from other predators, Dove makes their nest in the vicinity of Drongo's nest.



Figure 3 (a)



Figure 3 (b)

**Figure 3: Photographs presenting nest of Black Drongo (*Dicrurus macrocercus*)**

#### **Eurasian collared Dove (*Streptopelia decaocto*):**

The nesting season for Eurasian collared Dove is throughout the year. The nest was



**Figure 4: A photograph presenting the nest of Eurasian collared Dove (*Streptopelia decaocto*) Detailed Map of Study area**

observed at Point 8 in the plantation site of Saurashtra University (22°17'7.37"N 70°44'53.88"E). It was made with twigs and rootlets at 8-9feet height. The clutch size is 1-2 (Robertson, 1990) and incubation period of eggs is 15-19 days. Within a distance of 3 meters, two nests of Eurasian collared Dove were found. Both nests were in the vicinity of Black Drongo's nest. It's a great example of species-species interaction. By making nests nearby aggressive Drongo's nests,



Dove could gain protection from other predator species (Jahan et al., 2018).

**Indian Thick-knee (*Burhinus indicus*):**

Indian Thick-knee is also known as Indian Stone curlew. Its nest was found at Point 8 in the plantation site of Saurashtra University (22°17'7.37"N 70°44'53.88"E). The breeding season is from March to April. The nest was made on the ground surrounded by dead leaves and few twigs. The clutch size for stone curlew is two (Sharma & Sharma, 2015) with incubation period around 24 days. While exploring the study area, one egg and one hatchling were observed, which were wonderfully camouflaged with the colour of the ground. The egg was off-white coloured with random brown spots on it. At the end of May 2021, another egg was found broken. Both, male and female are involved in caring and protecting the hatchlings against predators (Sharma & Sharma, 2015).



Figure 5 (a) Nest with hatchling and egg



Figure 5 (b) Egg



Figure 5 (c) Hatchling



Figure 5 (d) Adult Indian Thick-knee

**Figure 5: Different photographs of Indian Thick-knee (*Burhinus indicus*) presenting developmental stages**



### **Shikra (*Accipiter badius*):**

The nesting season for Shikra is March to June. Shikra's nest was found at Point 8 (22°17'10.26"N 70°44'44.03"E) in the plantation site of the Saurashtra University. The nest built with twigs and rootlets at 20feet height from ground level was observed on the tallest tree. The clutch size for the Shikra is 3 to 4 eggs with 18-21days incubation period. Usually, females hatch the eggs but in rare cases males



hatch eggs as observed by Suryawanshi (2021). During the survey female Shikra was observed chasing Large-grey Babbler (*Argyamalcolmi*).

**Figure 6: A photograph of Adult female Shikra (*Accipiter badius*) and her nest**

### **Rufous Treepie (*Dendrocittavagabunda*):**

The breeding season for Rufous Treepie is April to June and with clutch size 3-6eggs.



**Figure 7: Adult Rufous treepie (*Dendrocitta vagabunda*) and her nest**

At Point 8 of Saurashtra University plantation site (22°17'7.37"N 70°44'53.88"E) two to three individuals of Rufous Treepie were seen.



One of them was seen carrying food material while the other was collecting the nesting materials. A single nest of Rufous Treepie was observed and was built with twigs, dead leaves, threads, old clothing material, rootlets, and cotton. The nest was found at a height of approximately 20 feet. On May 16<sup>th</sup>, 2021 Rufous Treepie was seen feeding juvenile Koel. After a few days, another Rufous Treepie was found hatching the eggs.

### **Red-wattled Lapwing (*Vanellus indicus*):**

Breeding season for Red-wattled Lapwing is April to June. The egg or nest was not part of the observation, however, a juvenile was observed at Point 3 of the Sub-urban area (22°17'2.38"N 70°44'45.14"E). A survey was conducted in May and the eggs might be laid during April. The clutch size is 3 to 4 with an incubation period of 28-30 days. The nesting of this bird has no protection strategies other than camouflaging the eggs with the surroundings to protect those (Balkhande et al., 2017).



*Figure 8 (a) Juvenile Red-wattled Lapwing      Figure 8 (b) Adult Red-wattled Lapwing*

***Figure 8: Photographic presentation of juvenile and adult Red-wattled Lapwing (*Vanellus indicus*)***

### **Cattle Egret (*Bubulcus ibis*):**

The breeding season for Cattle Egret is June to August. Between Point 5 and 6 (22°16'58.27"N 70°44'52.31"E), ten to twelve Cattle Egret nests were found and Egrets were observed in their breeding plumage. During their breeding season, adults develop orange buff plumage on the back, breast and crown area. Cattle Egrets from their nest in



colonies. Rootlets and twigs are used to build nests. Clutch size for Cattle Egret is 2-4 and the incubation period is between 22-28 days (Patankar et al., 2007).



Fig. 9 (a)



Fig. 9 (b)

**Figure 9: Cattle egrets (*Bubulcus ibis*) and their colony**

#### **Acknowledgements:**

I would like to express my gratitude to Dr. Jagruti Rathod for guiding and inspiring me throughout my study. I would also like to thank all the members of WCB Research Foundation for organizing this internship program and giving me a chance to expand my knowledge.

#### **References:**

- Ali (2017), *The Book of Indian Birds*. Bombay Natural History Society. ISBN 8173714851.
- Ali, Hadjela & Hassan, Syed & Rana, Shahnaz & Beg, Mirza & Mahmood-ul-Hassan, Muhammad. (2007). Brood parasitism of Asian koel (*Eudynamys scolopacea*) on the house crow (*Corvus splendens*) in Pothwar region of Pakistan. *Pakistan Journal of Agricultural Sciences*. 44. 627.
- Balkhande, Dr. Jayvardhan. (2017). Study of Nesting pattern and breeding Biology of Red-Wattled Lapwing *Vanellus Indicus* in agricultural field near to Asna River bridge, Nanded (Maharashtra) *Research maGma An International Multidisciplinary Journal* ISSN NO- 2456-7078 VOL-1, ISSUE-6, AUG-2017.
- Grimmett *et al* (2011), *Birds of Indian Subcontinent*. Helm field guides. ISBN: 9781408127636



- Jahan, I., Begum, S., Feeroz, M. M., Das, D. K., & Datta, A. K. (2018). Nesting pattern of birds in Jahangirnagar University Campus, Bangladesh. *Journal of Threatened Taxa*, 10(5), 11618–11635. <https://doi.org/10.11609/jot.2799.10.5.11618-11635>
- Nijman, V. (2004). Seasonal variation in naturally occurring mobbing behaviour of drongos (Dicuridae) towards two avian predators. *Ethology Ecology and Evolution*, 16(1), 25–32. <https://doi.org/10.1080/08927014.2004.9522651>
- Patankar, Pratyush & Desai, Isha & Shinde, Kavita & Balakrishnan, Suresh. (2007). Ecology and breeding biology of the Cattle Egret *Bubulcus ibis* in an industrial area at Vadodara, Gujarat. *Zoos' Print Journal*. 22. 10.11609/JoTT.ZPJ.1566.2885-8.
- Robertson, H. A. (1990). Breeding of collared doves *Streptopelia decaocto* in rural Oxfordshire, England. *Bird Study*, 37(2), 73–83. <https://doi.org/10.1080/00063659009477043>
- Sharma, M., & Sharma, R. K. (2015). Ecology and breeding biology of Indian stone curlew (*Burhinus indicus*). *Nature Environment and Pollution Technology*, 14(2), 423–426.
- Suryawanshi, K., 2021. Nesting behaviour and diet of Shikra *Accipiter badius* in Ajanta, Maharashtra. *Indian BIRDS* 17 (2): XX–XX
- Trivedi, V., & Vaghela, S. (2020). Avifauna of Saurashtra University campus, Rajkot, Gujarat, India. *Journal of Threatened Taxa*, 12(13), 16764–16774. <https://doi.org/10.11609/jott.5113.12.13.16764-16774>

---

**Suggested citation:**

Dave D. (2021). Observations on nesting pattern of different bird species around university road, Rajkot. *Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab*. Vol 1(2) 4-11.







## Record of Leucism in Indian Flying Fox *Pteropus giganteus* (Mammalia: Chiroptera: Pteropodidae) from South Gujarat, India

Amin Siddharth M<sup>1</sup>, Patel Hiren J.<sup>2\*</sup> and Vaghela Prashant P.<sup>3</sup>

<sup>1</sup> C/129 Bhagyoday Society, Opposite Sahyog Police chowki, Gorwa Refinery Road, Vadodara - 390024(Gujarat), India.

<sup>2\*</sup> Sardar Patel Zoological Park, Kevadiya – 393151(Gujarat), India.

<sup>3</sup> C-2/7 New Narmada Colony, Chhani Jakatnaka, Vadodara - 390024(Gujarat), India.

\*E-mail: [hirenp9408@gmail.com](mailto:hirenp9408@gmail.com)

### Abstract:

Leucism in fruit bat species *Pteropus giganteus* (Brunnich, 1782) is recorded for the first time from India based on photographic evidence collected from the Anaval Village located in Surat District of South Gujarat, India. A total lack of pigmentation in the whole body due to the failure of melanocytes to migrate to the skin, and hair follicles, resulting in white or whitish hairs, pale skin, but normal coloured eyes are the sign of leucism. The present record can serve as baseline data and will also help understand colour aberration in bats.

**Key Words:** Colour aberration, Chiropteran, Pigmentation, Mutation

### Introduction:

The Indian Flying Fox is one of the most common and widely distributed chiropterans found in India, Pakistan, Sri Lanka, Bhutan, Nepal, Bangladesh, Maldives, Myanmar and China (Tsang 2020; Johnsingh & Manjrekar 2013; Srinivasulu & Srinivasulu 2012). Flying Fox is the largest fruit bat known to occur on mainland India (Bates et al. 1994) out of 13 Indian fruit bat species (Bates & Harrison 1997; Srinivasulu et al. 2010). It has a reddish-brown



**Figure 1:** Leucistic individual roosting along with the normal individuals

head with a darker, sometimes blackish, snout; pale brownish-yellow shoulders and hindneck, yellowish-brown ventrally and black wings (Prater 1971). It is a colonial species and roosts during the day, often in the midst of busy towns, villages, on large trees along the roadside, near cropland and water bodies (Bates et al. 1994; Johnsingh & Manjrekar 2013; Menon 2014). Their roosting



colonies are generally found on large trees such as *Ficus benghalensis*, *F. religiosa*, *Tamarindus indica*, *Mangifera indica*, *Dalbergia sissoo* and *Eucalyptus sp* (Vendan 2003) and the colony size can vary from hundreds to several thousands of individuals (Bastawade & Mahabal 1976). Although fruit bats are responsible for many ecological services (Saikia 2018), yet Indian Flying Fox is one of the most persecuted fruit bats in South Asia and is listed as vermin under Schedule V of the Indian Wildlife (Protection) Act and IUCN Red List Data included it as a Least Concern species (Tsang, 2020).

In mammals, a wide variety of colours and forms are found as a result of the presence of certain pigments, mostly melanin which provides colour to the skin, hairs and eyes (Nordlund et al. 1998; Uieda 2000; Lucati & López Baucells 2016). Often this pigment is affected as a result of mutations giving rise to diverse colour aberrations. The effect of such abnormal colouration is generally the pigmentation anomalies of chromatic disorders (Rook et al. 1998). Similar to other mammals, bats are also vulnerable to genetic disorders that affect pigmentation. Globally, such chromatic disorders are being reported at an increasing rate in bats (Lucati & López Baucells 2016) and are caused by either an increase or a decrease in the production of melanin in some regions or over the entire body (Hofreiter & Schoneberg 2010; Abreu et al. 2013). These include albinism, leucism, piebaldism, hypomelanism and melanism (Lucati & López Baucells 2016). Among these conditions, leucism stands out as a rare phenomenon in nature.

Leucism is a total lack of pigmentation in the whole body due to the failure of melanocytes to migrate to the skin, and hair follicles, resulting in white or whitish hairs, pale skin, but normal coloured eyes. It occurs regardless of the normal production of the enzyme tyrosinase and melanin. Moreover, it can be caused by one of the several mutations that will give rise to apparently similar phenotypes (van Grouw 2006, 2013; Abreu et al. 2013; Lucati & López- Baucells 2016). The occurrence of colour aberration in 55 Indian mammalian species has been compiled by Mahabal et al. (2019). Although cases of albinism and piebaldism have been reported in the family Pteropodidae (Neal 1971; Karim 1983; Anonymous 2012; Anonymous 2013), however, there is no report on colour aberration in Indian Flying Fox (*Pteropus giganteus*). Here we report the first case of leucism in Indian Flying Fox from Surat district of South Gujarat, India.

**Observations:**

On 18 June 2017, we were on the way to Vansda National Park and at 1230 h we stopped as we saw a group of Indian Flying Fox roosting on *Ficus religiosa*, near Anaval Village (20.838°N, 73.263°E) in Surat District, Gujarat. Suddenly all the fruit bats flew in the sky, as a local was hunting them with the help of a catapult maybe for eating purposes. In the middle of all these, at 1249 h a white bat flying in the sky grabbed our attention as it flew further to roost on one of the branches with other Indian Flying Fox. We took some photographs (Figure 1) and observed with the help of binoculars to note down its characteristics. Its whole body i.e. both fur and skin were white or whitish including patagium and ears but it had normal black coloured eyes. With the help of the classification proposed by Lucati & López Baucells (2016), we identified it as a case of leucism and not albinism.

The roosting colony had around 250 individuals and was situated near a village pond among human habitation on the roadside. We observed the leucistic individual at least for an hour and also alerted the locals to cease hunting of bats. The leucistic individual was probably a female and showed similar activity as that of normal individuals such as opening and closing of wings, folding the wings around belly and grooming.

**Discussion:**

Although physiological causes of colouration, including melanism, are evident (Caro 2005), the genetic processes responsible for these colour variations remain unknown in most mammals (Lucati & López Baucells 2016). Moreover, chromatic disorders in bats can affect both fur and skin, including patagium, ears, and muzzle, and their impact on the survival rate is vague (Lucati & López Baucells 2016). However, bats are most likely less vulnerable to predation than the majority of diurnal animals as they are active at night and prefer dark roosting sites (Buys et al. 2002; Rocha et al. 2013). Lucati & López Baucells (2016) states that aberrant individuals roosting in sheltered places such as caves, mines, and buildings may have greater survival rates than the one roosting in the open leaves. Here, we found the leucistic Indian Flying Fox roosting in the open leaves and is likely to be vulnerable to predation.

Albinism is one of the most commonly occurring colour aberrations in bats (Lucati & López Baucells 2016). In the family Pteropodidae, albinism has been reported in *Cynopterus sphinx*



from Nepal (Anonymous 2013), *Lissonycteris angolensis* from Uganda (Neal 1971) and *Rousettus leschenaultia* from India (Karim 1983), while piebaldism is reported in *Rousettus amplexicaudatus* from Philippines (Anonymous 2012). However, another such colour aberration leucism is not yet reported from the family Pteropodidae. In this regard, our report of leucism in *Pteropus giganteus* can be the first of its kind in the entire range of its distribution.

Today, bats in India face the harsh reality of declining habitats and resources, ultimately making them disappear altogether (Graham 2001; Tsang 2020) and the people's perception regarding the presence of bats in their surroundings for different myths. These facts raise an urgent need to protect bats throughout the world by studying more about their behaviour and ecology. Further, studies on *Pteropus giganteus* from Gujarat are mainly sighting reports (Mahato et al. 2012) and on roosting sites (Vyas & Upadhyay 2014). Due to the lack of information on colour aberration in the family Pteropodidae, this report can add importance to bat biologists.

#### **Acknowledgement:**

We are thankful to Dhaivat Patel and Vivek Patel for assisting us in fieldwork.

#### **References:**

- Abreu, M. S. L., Machado, R., Barbieri, F., Freitas, N. S., & Oliveira, L. R., 2013. Anomalous colour in Neotropical mammals: a review with new records for *Didelphis* sp. (Didelphidae, Didelphimorphia) and *Arctocephalus australis* (Otariidae, Carnivora). *Brazilian Journal of Biology* 73: 185–194.
- Anonymous., 2012. Monfort Bat Sanctuary. Blanca Bella. Website: <http://www.monfortbatsanctuary.org/#!blanca-bella/c12m>. [Accessed 20 October 2019.]
- Anonymous., 2013. SMCRF (Small Mammals Conservation & Research Foundation). Website: <http://www.smcrf.org/index.php>. [Accessed 20 October 2019.]
- Bastawade, D. B., & Mahabal, A., 1976. Some behavioural aspects of the Indian flying fox *Pteropus giganteus giganteus*. *Bio-Vigyanam* 2: 209–212.
- Bates, P. J. J., & Harrison, D. L., 1997. *Bats of the Indian Subcontinent*. Harrison Zoological Museum Publication, UK. 258pp.
- Bates, P. J. J., Harrison, D. L., & Muni, M., 1994. The bats of western India. Part 1. *JBNHS*



91 (2): 1–15.

Buys, J., Heijligers, H., & Dorenbosch, M., 2002. First record of an albino long-eared bat *Plecotus auritus* in The Netherlands. *Lutra* 45: 49–52.

Caro, T., 2005. The adaptive significance of coloration in mammals. *BioScience* 55: 125–136.

Graham, G. L., 2001. *Bats of the World*. St. Martin's Press, New York, New York.

160pp. Hofreiter, M., & Schoneberg, T., 2010.

The genetic and evolutionary basis of colour variation in vertebrates. *Cellular and Molecular Life Science* 67: 2591–2603.

Johnsingh, A. J. T., & Manjrekar, N. (eds.) 2013. *Mammals of South Asia*, Volume 2. Universities Press (India) Private Limited, Telangana, India, pp. 545–547.

Karim, K. B., 1983. A note on an albino *Rousettus leschenaulti*. *Bat Research News* 24: 39–40.

Lucati, F., & López Baucells, A., 2016. Chromatic disorders in bats: a review of pigmentation anomalies and the misuse of terms to describe them. *Mammal review*, 47(2), 112–123.

Mahabal, A., Sharma, R. M., Patil, R. N., & Jadhav, S., 2019. Colour aberration in Indian mammals: a review from 1886 to 2017. *Journal of Threatened Taxa*, 11(6), 13690–13719. Mahato, A. K. R., Kumar, V. V., & Patel, N., 2012. Indian flying fox in Hamirsir Lake, Bhuj city needs conservation attention. *Current Science*, 103(4), 354–355.

Menon, V., 2014. *Indian mammals: a field guide*. Hachette, India, pp. 386. Tsang, S.M., 2020. *Pteropus giganteus* (errata version published in 2021). *The IUCN Red List of Threatened Species* 2020: e.T18725A194134899. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T18725A194134899.en>.

Downloaded on 10 April 2021

Neal, E., 1971. *Uganda Quest*. Collins, London, UK.

Nordlund, J. J., Boissy, R. E., Hearing, V. J., King, R. A., Oetting, W. S., & Ortonne, J. P. (Eds.) 1998. *The pigmentary system: physiology and pathophysiology*. Oxford University Press, New York.

Prater, S. H., 1971. *The Book of Indian Animals (Revised)*. Bombay Natural History Society, Bombay, pp. 200–201.

Rocha, P. A., Feijo, J. A., Donato, C. R., & Ferrari, S. F., 2013. Leucism in Seba's short-tailed bat, *Carollia perspicillata* (Linnaeus, 1758), from a rock shelter in northeastern



- Brazil. *Chiroptera Neotropical* 19: 1151–1153.
- Rook, A., Wilkinson, D. S., Ebling, F. J. B., Champion, R. H., & Burton, J. L., 1998. *Textbook of Dermatology*, 6th edn. Blackwell Science, Boston, Massachusetts, USA.
- Saikia, U., 2018. A review of chiropterological studies and a distributional list of the bat fauna of India. *Records of the Zoological Survey of India*, 118(3), 242–280.
- Srinivasulu, C., & Srinivasulu, B., 2012. *South Asian mammals: their diversity, distribution, and status*. Springer Science & Business Media.
- Srinivasulu, C., Racey, P. A., & Mistry, S., 2010. A key to the bats (Mammalia: Chiroptera) of South Asia. *Journal of Threatened Taxa* 2(7): 1001–1076.
- Uieda, W., 2000. A review of complete albinism in bats with five new cases from Brazil. *Acta Chiropterologica* 2: 97–105.
- Van Grouw, H., 2006. Not every white bird is an albino: sense and nonsense about colour aberrations in birds. *Dutch Birding*, 28(2), 79–89.
- Van Grouw, H., 2013. What colour is that bird? *British birds*, 106, 17–29.
- Vendan, S. E., 2003. Roost and diet selection in the Indian Flying Fox *Pteropus giganteus* (Mega chiroptera). M.Sc. Thesis, Madurai Kamaraj University, India. 85pp.
- Vyas, R., & Upadhyay, K., 2014. Study of the Indian Flying Fox (*Pteropus giganteus*) Colonies of Jambughoda Wildlife Sanctuary, Gujarat, India: Record of the largest roosting congregation at Targol. *Small Mammal Mail*, 6(1), 2–8.

---

**Suggested citation:**

Amin, S; Patel, H; and Vaghela P (2021). Record of Leucism in Indian Flying Fox *Pteropus giganteus* (Mammalia: Chiroptera: Pteropodidae) from South Gujarat, India. *Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab*. Vol 1(2) 12-17.





## Notes on behavioural observation of some water birds during the Annular Solar Eclipse

**Thakkar Karan\* and Gohil Bharatsinh**

*Department of Life Sciences, Maharaja Krishnakumarsinhji Bhavnagar University, Bhavnagar-364001 Gujarat, India.*

\*E-mail: [ksonpal07@gmail.com](mailto:ksonpal07@gmail.com)

---

### **Abstract:**

*In this communication, we report the behaviour of various water birds observed during the annular solar eclipse which happened on 26<sup>th</sup> November 2019. We observed the behaviour of eight species of birds viz. Greater Flamingoes *Phoenicopterus roseus*, the Indian Cormorant *Phalacrocorax fuscicollis*, Pied Avocet *Recurvirostra avosetta*, Black-winged stilt *Himantopus himantopus*, Eurasian spoonbill *Platalea leucorodia*, Northern shoveler *Spatula clypeata*, Little grebe *Tachybaptus ruficollis* and Dalmatian pelican *Pelecanus crispus* throughout the eclipse period i.e. from 8:05 AM to 10:51 AM. Various behavioural changes were observed in all the targeted species except the Indian cormorant. It rested on a pole for the whole eclipse period. Detailed observations are given in the note. Reduction in light levels due to solar eclipse does affect the activity of birds.*

**Key Words:** Solar Eclipse, Behaviour, Birds, Bhavnagar

An annular solar eclipse was visible from 8:05 to 10:51 AM on 26 December 2019 in India. Solar eclipse, a rare and uncommon environmental event, can influence and modulate the behaviour of animals (Murdin 2001; Gil-burmann & Beltrami 2003). Following observations were made during the eclipse at airport Road Lake (21.762837° N, 72.183624° E) in Bhavnagar city using Nikon binoculars (10\*50).

Greater flamingos (n=24) on the day of the eclipse were seen foraging constantly from 8:05-9:20 AM. All of them started calling together from 9:21 to 9:35 AM. After the peak of the eclipse (i.e. after 9:21 AM), many flamingo individuals started preening. Activities of Indian cormorants varied throughout the eclipse period. Pied avocets (n=13) were observed feeding through the initial phase of the eclipse i.e. from 8:05 to 8:50 AM. But pied avocets went onto the resting phase from 8:51 to 9:35 AM. Eight of them continued resting even after 9:35 AM while the other five started feeding again. Black-winged stilts (n=50) were seen resting in a flock throughout the study period. But as the eclipse increased, flocks of black-winged stilts from outside the lake started arriving and joined the original flock of 50. A group of five black-winged stilts arrived at 8:52 followed by a flock of 20 at 9:06 AM. Three other groups of 30, 50, and 10 black-winged stilts arrived from 9:21 to 9:35 AM. Eurasian spoonbills (n=30) were seen resting in water during the initial stage of eclipse i.e. from 8:08 to 9:20 AM. From 9:21 to 9:35 AM five spoonbills started feeding, 21 continued resting while 12



started vocalizing and four flew away from the lake. After 9:35 AM, 12 individuals continued calling, 29 rested in water whereas eight individuals arrived from outside the lake. Activities of northern shoveler (n=80) varied throughout the eclipse. Little cormorant (n=1) rested on a pole for most of the observation. It started feeding at 9:21 to 9:35 AM along with one little cormorant which arrived from somewhere outside the lake. After 9:35, both perched at the wood in the lake. Little grebe (n=1) spent most of the time feeding. Between 9:35 and 9:50 AM, it was joined by a flock of eight grebes that arrived from outside the lake. Pelicans (n=2) rested on a small patch of land in the lake throughout the study period.

**Table 1: Ethogram**

Bird	Time intervals						
	08:05-08:20	08:21-08:35	08:36-08:50	08:51-09:05	09:06-09:20	09:21-09:35	09:36-09:50
Greater flamingo	FD(n=24)	FD(n=24)	FD(n=24)	FD(n=24)	FD(n=24)	C(n=24)	PR(n=4), FD(n=17), C(n=3)
Indian Cormorant	Rp(n=2)	Rp(n=2)	Fi(n=2), FD(n=4)	Rp(n=2), Fb(n=2)	Rp(n=1)	Rp(n=1), Fb(n=1)	_
Pied Avocet	FD(n=13)	FD(n=13)	FD(n=13)	Rw(n=13)	Rw(n=13)	Rw(n=13)	FD(n=5), Rw(n=8)
Black winged stilt	Rw(n=50)	Rw(n=50)	Rw(n=50)	Rw(n=50), Fi(n=5)	Rw(n=55), Fi(n=20)	Rw(n=75), Fi(n=30), Fi(n=50), Fi(n=10)	Rw(n=165)
Eurasian Spoonbill	Rw(n=30)	Rw(n=30)	Rw(n=30), FD(n=4)	Rw(n=34)	Rw(n=34)	FD(n=5), C(n=12), Fo(n=4), Rw(n=21)	Fi(n=8), C(n=13), Rw(n=29)
Northern Shoveler	PR(n=10), RI(n=50), Rw(n=20)	RI(n=50), Rw(n=20)	PR(n=4), Rw(n=50), RI(n=10)	Rw(n=54), RI(n=10)	FD(n=5), Rw(n=54), RI(n=5)	Rw(n=54), RI(n=10)	Rw(n=54), RI(n=10)
Little Cormorant	Rp(n=1)	Rp(n=1)	Rp(n=1)	Rp(n=1)	Rp(n=1)	FD(n=1), Fi(n=1)	Rp(n=2)
Little grebe	Rw(n=1)	FD(n=1)	FD(n=1)	FD(n=1)	FD(n=2)	Fi(n=8), FD(n=10)	FD(n=18)
Dalmatian pelican	RI(n=2)	RI(n=2)	RI(n=2)	RI(n=2)	RI(n=2)	RI(n=2)	RI(n=2)



**Table 2: Different type of behaviours observed in bird species**

	<b>BEHAVIOURS</b>
FD	Feeding
R	Resting
Rp	resting on an object; prolonged perching on a pole or a stick in water
RI	resting on land; simply sitting or standing with no noticeable movement
Rw	resting in water; no efforts for movement and neck resting on body(in case of ducks)
F	Flight
Fb	taking off from place 'A' within a lake and settling at place 'B' within a lake
Fa	taking off from place 'A' within a lake and settling at same place
Fo	taking off from within lake and flying outside the lake
Fi	arrived at lake from outside the lake
PR	Preening
C	Calling; making some kind of voice, could either be begging call, roosting call, or some other
B	Bathing; Rapid action of submerging in water and coming out

Kumar S. (1981) observed the changes in the activity of cattle egrets because of the solar eclipse. Trigunayat (1997) observed that roosting Black-crowned Night-herons altered a variety of behaviours, including increased calling, preening, and perch shifting. Nanikov et al. (2001) also mention behavioural changes in several birds due to solar eclipse. A similar trend was observed in this study as well. From these limited field observations, we can conclude that reduced light level due to annular solar eclipse does interrupt normal bird behaviour patterns.

### **Acknowledgements:**

The first author is very thankful to Mr. Dhavalkumar Vargiya for his kind support and encouragement in the preparation for this article.

### **References:**

- Altmann, J., 1974. Observational Study of Behavior: Sampling Methods. *Behaviour*, 49(3/4), 227-267. Retrieved on March 23, 2020, from [www.jstor.org/stable/4533591](http://www.jstor.org/stable/4533591) • Gil-Burmann, C., Beltrami, M., 2003. Effect of the solar eclipse on the behavior of a captive group of hamadryas baboons (*Papio hamadryas*). *Zoo Biology*, 22: 299–303.
- Kumar S., 1981. Solar eclipse – notes on behavior of egrets. *J. Bombay natural history society*. 78, 3: 133-136
- Murdin, P., 2001. Effects of the 2001 total solar eclipse on African wildlife. *Astronomy*



*Geophysics*, 42, 40–42.

Nankinov, D., Dalakthieva, S., Nikolov, B., Shurulinkov, P., Popov, K., Kirilov, S., Zareva, K., Borisov, B., Stojanov, G., Nankinov, N., Minchev, N., Kjachukov, D., Nikolov, I., Stanchev, R., Ganchev, E., 2001. Studying the behaviour of birds during the solar eclipse on August 11, 1999. — *Russian Ornithological Journal*, 142: 355-369.

Trigunayat, M. M., 1997. Some behavioral observations on Night Heron *Nycticorax nycticorax* (Linnaeus) during the total solar eclipse in Keoladeo National Park, Bharatpur, India. *Pavo* 35:61–65.

Tramer, E. J., 2000. Bird Behavior During a Total Solar Eclipse. *The Wilson Bulletin*, 112(3): 431–432.

---

**Suggested citation:**

Thakkar K and Gohil B (2021). Notes on behavioural observation of some water birds during the Annular Solar Eclipse. *Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab. Vol 1(2) 18-21.*





## Restoration of Earth with Biobased Polymer

**Marjadi Darshan**

*Shree Ramkrishna Institute of Computer Education & Applied Sciences, MTB Campus,  
Athwalines, Surat-395001*

*E-mail: [darshan.marjadi@srki.ac.in](mailto:darshan.marjadi@srki.ac.in)*

---

Humanity is facing increasingly numerous challenges in life despite the great advancements in science and technology over the last century. All the needs of modern society, e.g. food, fuel, energy, and materials, are highly dependent on diminishing fossil resources. The demand for creating and using plastic throughout the world, driven by the increasing usage of plastic-based materials has been increasing and adding stress to the current waste management infrastructures. There is significant interest to decrease the reliance on petroleum-based plastic products, which causes global environmental pollution. The advancement in science and technology has enabled mankind to live longer and consume more of the world's resources. More than eight million tons of plastic waste leak into oceans every year, which can be eased through innovative redesigns of packaging materials.

Issues such as global climate change and depletion of fossil fuels, both resulting from the increased usage of energy, have triggered alarms among scientists and politicians worldwide. Therefore, governments, industries, and academia are inputting much effort towards finding a sustainable solution for the increasing energy crisis. However, a similar concerted effort is still lacking to bring about the creation and usage of sustainable materials in the modern lifestyle. In addition to the increasing demand for energy, the usage of materials in the modern lifestyle has also been increasing rapidly. Considering the widespread usage of plastics in food packaging, clothes, shelter, communication, transportation, construction, health care, and leisure industries; plastics are the most dependable material among the other materials that mankind is currently dependent upon. Currently, most of the plastics that are widely used in various sectors are produced from petrochemical products. The plastic industry has been among the most profitable businesses and is expected to grow further, especially because of their increased demands from rapidly developing countries like India and other parts of South-East Asia. While plastics are superior materials in terms of their production costs and diverse properties, the sustainability of this synthetic material is undoubtedly an issue that needs to be addressed. Biobased and biodegradable polymeric materials are considered amongst the most suitable alternatives for some applications. The



excessive usage of petroleum also contributes to the increased emission of CO<sub>2</sub> into the atmosphere, which is thought to be among the principal reasons for global warming and climate change. These are some of the strong and valid reasons leading towards the development of technologies to produce biobased and biodegradable plastics.

Science and technology have created the modern lifestyle that mankind is living in which efficiency is the keyword. Drawing an analogy from a bacterial culture in a shaken flask, the depletion of nutrients and the simultaneous accumulation of toxic wastes are the main reasons for the onset of the stationary growth phase and the subsequent death phase. However, unlike bacteria, mankind is aware of the consequences of such activities. As a result, the world population is now concerned about sustainable development.

Biobased and biodegradable plastics most likely to remain less superior in terms of production cost and material properties compared to petrochemical plastics. Petroleum-based plastics have serious environmental and social impacts due to their non-degradable nature and the leaching of carcinogenic substances like dioxin when exposed to scratch or heat. Compared to the ease at which fossil resources are utilized; renewable biobased materials will need more effort and efficiency to be used in daily life. However, biobased materials are considered superior in terms of sustainability and environmental friendly provided that they are used judiciously and in a contented manner. Hence, it is in our hands to utilize our knowledge of the environment to work towards the restoration of the earth and make it free from single-use plastics.

---

***Suggested citation:***

*Marjadi D (2021). Restoration of Earth with Biobased Polymer . Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab. Vol 1(2) 22-23.*





## Single-use surgical mask: Human lifesaver or Biodiversity destroyer


Patel Jeel<sup>1</sup> and Patel Rushika<sup>2,3\*</sup>

<sup>1</sup>GMERS, Sola Civil Medical College, Ahmedabad-380008 (Gujarat), India.

<sup>2</sup>Assistant Professor, Rai University, Dholaka, Ahmedabad-380008 (Gujarat), India;

<sup>3</sup>Fellow, Wildlife and Conservation Biology Research Foundation, Pata-384265 (Gujarat), India.

\*E-mail: [rush2907@gmail.com](mailto:rush2907@gmail.com)



The green lines represent sustainable and biodiversity conservation ways. The dark brown lines represent possible threats of single-use surgical mask to the biodiversity.

**Key Words:** Disposable mask; Plastic and micro plastic pollution; Conservation; Threats

The COVID-19 is transmitted by respiratory droplets or aerosols and can survive in the air for more than 24 hrs, which was detected and momentarily created a pandemic situation, globally. The only way to stop the transmission is to wear face mask, washing hands frequently and to maintain social distance. Wearing a mask is declared as one of the important precautionary measures by the World Health Organization (WHO) and policymakers in every country, because masks help in the significant reduction of infection caused by touching the face/mouth/nose with contaminated hands (Schmutz et al. 2020). The virus, COVID19 has increased the demand for the surgical single-used mask in the healthcare professionals as well as among the community. Such masks are made up of polypropylene or



high-density polyethylene (HDPE), a fossil-derived plastic which has a very high half-life (Dharmaraj et al. 2021). The mask production to the disposal statistics is presented in Table 1. Based on the reusability, the mask is divided into two major groups' disposable (made up of polymers) and reusable mask (made up of cloth material) in terms of production. In present study, the details of only disposable mask is presented.

**Table 1: Global mask production, usage and disposable statistics**

No	Statistics	Reference
1	Estimated worldwide usage of disposable mask: 129 billion/month	Prata et al. 2020
2	Production in China as on Feb. 2020: 116 million/day	Adyel 2020
3	Order of mask in Japan as of April 2020: over 600 million/month	Fadare and Okoffo 2020
4	Mask disposal in the UK: 53 million/day	Website URL 1
5	Gloves and mask entered into the environment: 200 million/month	Website URL 2
6	Mask reached to oceans: 1.5 billion Marine plastic pollution increased in last year: 6200 tonnes	Website URL 3
7	Mask waste generation in India: Minimum: 928/week; Maximum: 4640/week	Selvaranjan et al 2021
8	*Plastic footprint in the environment due to mask and gloves: $4.5 \times 10^5$ Kg/Month	Present Study

\*Calculated based on the global entries of masks and gloves mentioned (200 million/month). The average of one mask and single glove weight is  $2.25 \pm 0.01$ .

#### **Mask: a threat to the ecosystem:**

In current situation, recycling of any household medical waste is not a feasible and advisable solution, as there is a chance that it may be exposed to infectious particles. There are no particular laws or guidelines for separation and segregation, which might involve several issues. Globally, the dumping of masks was observed in areas such as, roadside, river, oceans and several open areas. Usually, such masks sustain in the environment for a longer period of time (more than 400 years)(Hasan et al., 2021). The misconception of masks to various birds, animals and marine creatures, considering as a food and leads to trapping and entangling in the strips of masks. The single-use surgical mask not only adds litter to the environment, but



also hurting and killing a wide range of terrestrial and marine living organisms. Moreover, the production process of such masks also releases greenhouse gases (Prata et al. 2021). The disposal of mask also reported as a novel source of micro plastic pollution in the environment and entry point of the food chain (Hasan et al. 2021). The single-use surgical masks are also noted as a major causative agent of secondary disease outbreaks in all life forms, including humans (Fadare and Okoffo 2020). India does not have any sufficient resources for the household medical waste management (Prata et al. 2021). Every minute 10 kg of disposed masks and gloves entered into the environment (Table 1). Hence, it is very important to deal with the mentioned issues and look into more sustainable sources at individual level. Herewith, we suggest some measures by which we can protect the environment without compromising human health in current pandemic situation.

- Snip the strips of the masks before the disposal to avoid the entangling and entrapping issues.
- Dispose masks as a separate medical waste and submit it to plastic recycling companies.
- Moving towards the development of biopolymer mask (Torres and Torre 2021). Biopolymer mask were made up of gluten and nanofibers having 95% efficiency.
- A reusable mask could be applied in routine life for the general community. Regular cloth masks having multi-layer of high thread-count and textiles are moderately effective and reported to block 50-70 % fine droplets and particles. Reusable masks help in the reduction of waste production (85 %). Moreover, economically viable option in a country like India as it is 3.7 times cheaper than other masks (Prata et al. 2021).

### References:

- Adyel, T. M., 2020. Accumulation of plastic waste during COVID-19. *Science*, 396(6509): 1314–1315. <https://doi.org/10.1056/nejmms2004740>.
- Dharmaraj, S., Ashokkumar, V., Hariharan, S., Manibharathi, A., Show, P. L., Chong, C. T., & Ngamcharussrivichai, C., 2021. The COVID-19 pandemic face mask waste: A blooming threat to the marine environment. *Chemosphere*. Elsevier Ltd. <https://doi.org/10.1016/j.chemosphere.2021.129601>.
- Fadare, O. O., & Okoffo, E. D., 2020. Covid-19 face masks: A potential source of microplastic fibers in the environment. *Science of The Total Environment*, 737: 140279. <https://doi.org/10.1016/j.scitotenv.2020.140279>.



- Hasan, N. A., Heal, R. D., Bashar, A., & Haque, M. M., 2021. Face masks: protecting the wearer but neglecting the aquatic environment? - A perspective from Bangladesh. *Environmental Challenges*, 4: 100126. <https://doi.org/10.1016/j.envc.2021.100126>
- Prata, J. C., Silva, A. L. P., Duarte, A. C., & Rocha-Santos, T., 2021. Disposable over Reusable Face Masks: Public Safety or Environmental Disaster? *Environments*, 8(4), 31. <https://doi.org/10.3390/environments8040031>.
- Schmutz, M., Hischier, R., Batt, T., Wick, P., Nowack, B., Wäger, P., & Som, C., 2020. Cotton and surgical masks—what ecological factors are relevant for their sustainability? *Sustainability (Switzerland)*, 12(24): 1–13. <https://doi.org/10.3390/su122410245>.
- Selvaranjan, K., Navaratnam, S., Rajeev, P., & Ravintherakumaran, N., 2021. Environmental challenges induced by extensive use of face masks during COVID-19: A review and potential solutions. *Environmental Challenges*, 3: 100039. <https://doi.org/10.1016/j.envc.2021.100039>.
- Torres, F. G., & De-la-Torre, G. E., 2021. Face mask waste generation and management during the COVID-19 pandemic: An overview and the Peruvian case. *Science of the Total Environment*, 786: 147628. <https://doi.org/10.1016/j.scitotenv.2021.147628>.
- Website URL 1: <https://www.greenpeace.org.uk/news/single-use-face-masks-hurting-wildlife-what-you-can-do/> [Accessed on 23<sup>rd</sup> June 2021].
- Website URL 2: <https://www.greenmatters.com/p/face-masks-wildlife> [Accessed on 23<sup>rd</sup> June 2021].
- Website URL 3: <https://economictimes.indiatimes.com/news/environment/ecosystem-under-threat-as-birds-and-marine-life-ensnared-in-discarded-face-masks/widespread-contamination/slideshow/80230026.cms> [Accessed on 23<sup>rd</sup> June 2021].

---

**Suggested citation:**

Patel J and Patel R (2021). *Single-use surgical mask: Human lifesaver or Biodiversity destroyer* Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab. Vol, 1 (2) 24-27.







## Does the contemplation of forest field staff about wildlife differ than a common man?

Pithva Krishna<sup>1,2</sup> and Dharaiya Nishith<sup>1</sup>

<sup>1</sup> Wildlife and Conservation Biology Research Lab, Hemchandracharya North Gujarat University, Patan (Gujarat) India

<sup>2</sup>Institute of Environment Education and Research, Bharati Vidyapeeth Deemed University, Pune, India

\*E-mail: [krishnapithva17@gmail.com](mailto:krishnapithva17@gmail.com)

### Introduction:

A field-based training was organized by Vadodara wildlife division, Gujarat forest department and WCB Research Lab of Hemchandracharya North Gujarat University from September 17<sup>th</sup> to 19<sup>th</sup>, 2020 with the aim to enhance the capacity of forest field staff for monitoring and rescue of sloth bear and associated fauna. The training was organized at Jambughoda wildlife sanctuary, Central Gujarat. Along with the awareness about sloth bear, a small survey was carried out to understand how the field staff envisages the wildlife that is found in their work place. Total 18 frontline forest staff of different cadre such as beat guards and round foresters from different forest ranges of central Gujarat has participated in this survey.



This study provides an insight on how forest staff's perception differs in different animals. As Q method provides qualitative and quantitative data which helps identifying people's perception in detail. As participants have to provide justification of their answers, it reveals some underlying conflicts or reasons. These justifications of participants can help identify the gaps and can be better used for conservation planning as well as capacity building of forest field staff.

*Figure 1: Forest frontline staff organizing photos on Q Board © Nishith Dharaiya*




**Methodology:**

We used Q methodology for this study; this method is designed and developed by William Stephenson in 1930s (McKeown & Thomas, 1988) and allows to disclose underlying reasons. This method is widely used in social sciences studies. In this method, photos were used instead of statements to know respondent’s perception allowing them to justify their answer in detail without any restriction. This method allows both qualitative and quantitative data on perception of the person being interviewed.



**Figure 1(a):** Representative sort of organized Q-board by a participant



**Figure 1(b):** A Q-board on which the respondents organize the photos

We categorised the Q Method into two parts, first, organising 16 photos of locally found animals on a Q-board (figure 1(b)) followed by explanation for each photo which are placed on Q-board by the respondent. As shown in the figure 1(a), the participants were asked to organise photos in Q-board as per their liking and disliking towards the animals and to provide reasons which were recorded in mobile phone device. A list of all the wildlife photos used for this survey is provided in table 1.



**Table 1: Photos of wildlife used for this study**

Photo ID	Common name	Scientific name
1	Wild boar	<i>Sus scrofa</i>
2	Hanuman langur	<i>Semnopithecus entellus</i>
3	Indian cobra	<i>Najana naja</i>
4	Indian giant flying squirrel	<i>Petaurista philippensis</i>
5	Small Indian civet	<i>Viverricula indica</i>
6	Sloth bear	<i>Melursus ursinus</i>
7	Indian hare	<i>Lepus nigricollis</i>
8	Indian python	<i>Python molurus</i>
9	Barn owl	<i>Tyto alba</i>
10	Grey francolin	<i>Francolinus pondicerianus</i>
11	Red-wattled lapwing	<i>Vanellus indicus</i>
12	Rhesus macaque	<i>Macaca mulatta</i>
13	Black kite	<i>Milvus migrans</i>
14	Common krait	<i>Bungarus caeruleus</i>
15	Indian crested porcupine	<i>Hystrix indica</i>
16	Blue bull	<i>Boselaphus tragocamelus</i>

### Q sort analysis:

In this study, three factors were derived based on participant's justification which is explained in detail in the result section. A factor in this study is a category representing the group of people who have similar perspective (Brown, 1980). The higher the factor loading, the more highly that sorts are correlated with that factor (Ramlo, 2008, Ramlo & Newman, 2011). The sorts refer to the photos assembled by participants on Q-board (figure 1(b)). In order to analyse the data, a software, PQMethod (<http://schmolck.org/qmethod/>) was used which is specially designed for Q-analysis. Q sorts are the number of participants that took part in the survey. All the data were entered manually in this software and correlation was calculated among each sort. The correlation matrix for the extracted factor was analysed through a principal component factor analysis with varimax rotation for which options are provided in the software (figure 2).



```

C:\PQMethod\PQMethod.exe
(Mar 2014)
-----
by Peter Schmolck
Adapted from Mainframe-Program QMethod
by John Atkinson at KSU
-----
The QMethod Page:
http://schmolck.org/qmethod/
-----

Enter [Path and] Project Name:
ForestQ

Current Project is ... C:\PQMethod\ForestQ
Choose the number of the routine you want to run and enter it.

1 - STATES - Enter (or edit) the file of statements
2 - QENTER - Enter q sorts (new or continued)
3 - QCENT - Perform a Centroid factor analysis
4 - QPCA - Perform a Principal Components factor analysis
5 - QROTATE - Perform a manual rotation of the factors
6 - QVARIMAX - Perform a varimax rotation of the factors
7 - QANALYZE - Perform the final Q analysis of the rotated factors
8 - VIEWLIST - View output file ForestQ.lis
X - Exit from PQMethod

Last Routine Run Successfully - (Initial)

```

**Figure 2: Screenshot of analytical option available in PQM method Software**

Participants having similar perceptions are put together in their respective factors and marked in bold and have “X” next to their score. Out of 18 sorts, 17 sorts were found complete by the software and were further analysed. Each of three factors represents a different perspective towards the photos provided to them. Participants were named as PART001, PART002 and so on to keep their identity unrevealed.

In table 2, there are 8 participants belonging to factor 1 with 29% of the variance explained followed by 7 participants in factor 2 having 26% explained variance and 2 participants belonging to factor 3 with 14% explained variance. Once factor score calculated by the software, distinguishing tables were developed in this analysis for each factor which differ from each other that is explained further in this section. The distinguishing tables for each factor explains differences between factors (Brown, 1971, Ramlo & Newman, 2011, Brown, 1993). In order to determine distinguishing statements, average Z-score of respondent’s factor score was calculated by the software.

## Results and Discussion:

The three factors that emerged are shown in Table 2 with automatic pre-flagging. Participants having similar perceptions are put together in their respective factors and marked in bold and have “X” next to their score. Out of 18 sorts, 17 sorts were found complete by the software and were further analysed. Each of



*Table 2: Extracted factor score from Q-sorts*

Q-Sorts	ID	Factor 1	Factor 2	Factor 3
1	PART001	<b>0.8530X</b>	0.1248	-0.1223
2	PART002	0.0414	<b>0.8084X</b>	0.1708
3	PART003	<b>0.6945X</b>	0.0471	0.3983
4	PART004	<b>0.7683X</b>	0.1839	-0.0156
5	PART005	<b>0.5450X</b>	0.0166	0.2332
6	PART006	0.4208	<b>0.7241X</b>	0.4006
7	PART007	0.1728	0.1377	<b>0.9013X</b>
8	PART008	<b>0.6930X</b>	0.5214	0.3237
9	PART009	<b>0.6516X</b>	0.4289	0.4265
10	PART010	0.5119	<b>0.6142X</b>	0.0921
11	PART011	<b>0.8475X</b>	-0.0072	-0.2626
12	PART012	0.3742	-0.5333	0.4456
13	PART013	0.0481	<b>0.8292X</b>	0.3240
14	PART014	-0.0002	<b>0.6395X</b>	-0.3574
15	PART015	<b>0.6577X</b>	0.2179	0.1940
16	PART016	-0.0614	0.1991	<b>0.6372X</b>
17	PART017	0.4469	<b>0.7611X</b>	0.1835
18	PART018	0.5679	<b>0.7564X</b>	0.1465
<b>%</b>	<b>Explained variance</b>	<b>29</b>	<b>26</b>	<b>14</b>

*Note: number in bold shows respondents belong to those respective factors.*

Table 3 shows different factor score for each animal photo and depending on statistical significance the photo load to a specific factor. The above table contains 16 photos and their grid position for all three factors (perception, table 2). For example, Indian python was disliked by participants therefore it is scored at -2 for respondents grouped under factor 1, +1 for factor 2 and for factor 3 it was scored at +2. For factor 1 the most liked species by participants is Indian hare and most disliked species is wild boar. Most liked species for factor 2 is Indian leopard and most disliked animal is fruit bat. For factor 3, most liked species is Indian hare scored at +3 and most disliked species is wild boar scored at -3 by the participants.



**Table 3: Aggregate factor values of each 16 photos**

No.	Photos	Aggregate values		
		Factor 1	Factor 2	Factor 3
1	Indian python	-2	1	2
2	Hanuman langur	0	0	2
3	Monitor lizard	-1	-2	-1
4	Red-wattled lapwing	1	-1	1
5	Sloth bear	2	2	0
6	Indian hare	3	1	3
7	Leopard	2	3	0
8	Grey francolin	1	-1	-1
9	Indian giant flying squirrel	1	1	-2
10	Blue bull	0	0	0
11	Small Indian civet	0	0	-1
12	Fruit bat (Flying fox)	-1	-3	-2
13	Indian cobra	-2	2	1
14	Barn owl	0	0	0
15	Indian chameleon	-1	-1	0
16	Wild boar	-3	-2	-3

#### Factor 1: Economic impact

This factor was described by 8 participants mainly concerning economic impact. Animals that cause harm economically by destroying crops and threat to human life and livestock. Participants of factor 1 thinks fruit bat, Indian python and Indian cobra causes high level of economic harm. Grey francolin is scored +1 due to its contribution to reduce impact by eating pest insects from agricultural field. Though the most disliked animal is wild boar for factor 1 as shown in table 3.

**Table 4: Distinguishing photos for factor 1**

Photo ID	Photos	Grid position for Factor 1	Z-score of Factor 1	Grid position for Factor 2	Z-score of Factor 2	Grid position for Factor 3	Z-score of Factor 3
8	Grey francolin	1	0.95	-1	-0.46	-1	-0.67
12	Fruit bat (Flying	-1	-0.2	-3	-1.82	-2	-1.1
1	Indian python	-2	-1.03	1	0.34	2	1.22
13	Indian cobra	-2	-1.19	2	1.27	1	0.37

Factor 2: Aesthetic, spiritual values and conservation aspects

This factor explains aesthetic, spiritual values and conservation aspects which includes appearance, tourist attraction, rehabilitation and religious belief. There are 7 responses in this factor (table 2). Respondents are observed to have negative opinions towards animals which play major role in economic loss but positive towards animal's beauty and its natural charisma. Table 5 shows how score of animals for factor 2 is different from the score of factor 1 and factor 3. For example, Indian cobra is scored at +2 in factor 2 as it is attractive to participants but it is scored -2 for factor 1 as it threatens human life.

**Table5: Distinguishing photos for factor 2**

Photo ID	Photos	Grid position for Factor 1	Z-score of Factor 1	Grid position for Factor 2	Z-score of Factor 2	Grid position for Factor 3	Z-score of Factor 3
13	Indian cobra	-2	-1.19	2	1.27	1	0.37
6	Indian hare	3	1.48	1	0.67	3	1.89
1	Indian python	-2	-1.03	1	0.34	2	1.22
4	Red-wattled lapwing	1	0.14	-1	-0.65	1	0.79
16	Wild boar	-3	-1.89	-2	-1.25	-3	-2.01



### Factor 3: Lack of awareness

This factor describes lack of awareness which includes lack of interest compared to other animals and species. This factor mainly concerns individuals who lacks knowledge, proper information and are not aware about the species. For example, in table 6, the score of Indian giant flying squirrel differs from factor 3 to factor 1 and 2 as it is scored at -2, +1 and +1 respectively. The justification for negative score given by the participants is that they are not aware about the presence of the species therefore they do not have any information on the species.

**Table 6: Distinguishing photos for factor 3**

Photo ID	Photos	Grid position for Factor 1	Z-score of Factor 1	Grid position for Factor 2	Z-score of Factor 2	Grid position for Factor 3	Z-score of Factor 3
1	Indian python	-2	-1.03	1	0.34	2	1.22
2	Hanuman langur	0	-0.2	0	-0.45	2	1.10
13	Indian cobra	-2	-1.19	2	1.27	1	0.37
5	Sloth bear	2	1.38	2	1.35	0	-0.00
7	Leopard	2	1.33	3	1.81	0	-0.00
9	Indian giant flying squirrel	1	0.96	1	0.62	-2	-1.22

### Conclusion:

The perception of forest field staff was better understood using Q method analysis considering various aspects. It is seen that participants tend to like animals such as Indian leopard, sloth bear, Hanuman langur and Indian hare as they are scored positive or neutral. The animals that are not given negative score in any factor due to their appearance, are seen frequently and does not lack awareness. It is observed from the data that frontline forest staff lacks knowledge about animals that are not seen or present in their forest range/beat. As frontline forest staff, they should be aware about existence of animal species that are found in Gujarat state along with their ecological importance, threats and conservation values. Some participants were not even aware about the existence of the species in the wild such as Indian giant flying squirrel. Although, being frontline forest staff, their perception is biased towards animals such as wild boar, monitor lizard and flying fox bat due to the economic loss they





may cause and weird appearance. Their perception becomes the same as a common man when every animal should be equal to frontline forest staff as it is their job to protect forest and its animals. Some participants believe in superstitions of bad luck of barn owl and monitor lizard which shows their common man perception and lack of right information.

### **Recommendations:**

A separate training or workshops should be organised by forest department focused on animal species found in Gujarat. This will help them enhancing their existing knowledge and it will provide them with some scientific insights. Having basic scientific knowledge will help remove some barriers like superstitions and other beliefs. Field trips should be organised in different part of state to gain practical experience. Team building activities should be conducted between staff of different forest divisions.

### **Acknowledgements:**

We are thankful to all the forest field staff who participated in this survey. We also thank to the Deputy Conservator of Forests, Vadodara Wildlife Division of Gujarat Forest Department and sincerely acknowledge the help from our research team members.

### **References:**

Brown, S. R. (1980). *Political subjectivity: applications of Q methodology in political science*.

E. Ramlo, S., and Newman, I. (2011). Q Methodology and Its Position in the Mixed-Methods Continuum. *Operant Subjectivity*, 34(3), 172–191.

McKeown, B. and Thomas, D. B. (1988). *Q Methodology*. SAGE Publications

R. Brown, S. (1993). A Primer on Q Methodology. *Operant Subjectivity*, 16(3/4), 91–138.

Ramlo, S. E. (2008). Determining the various perspectives and consensus within a classroom using Q methodology. *AIP Conference Proceedings*, 1064, 179–182.

<https://doi.org/10.1063/1.3021248>

---

### **Suggested citation:**

Pithva K and Dharaiya N (2021). Does the contemplation of forest field staff about wildlife differ than a common man? *Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab*. Vol 1(2) 28-36.





## Qualitative phytochemical analysis of some medicinal plants (leaves) from the surrounding area of Talod Taluka, North Gujarat, India.

**Panchal Mital\* and Jha Charumati**

Department of Botany, Smt. Panchal Science college, Talod –383215, North Gujarat, India.

\*E-mail: [mitalpanchal0@gmail.com](mailto:mitalpanchal0@gmail.com)

### **Abstract:**

The tremendous biological variety of North-Eastern India has long been recognised. Five medicinal plants were chosen for this study: *Manilkara zapota* (L.) van Royen, *Manilkara hexandra* (Roxb.) Dub., *Mangifera indica* L., *Madhuca indica* J.F. Gmel., and *Annona squamosa* L. The purpose of this study was to look into the existence of Phytochemicals such as Alkaloid, Amino acid, Carbohydrate, Protein, Glycoside, Tannin, Terpenoid, Saponin, Flavonoid, and Steroids in the medicinal plants that were chosen. For the organic solvent extraction, the Soxhlet apparatus was utilized. Water, acetone, and chloroform were utilised as solvents. Our findings show that the examined plants' crude aqueous and organic solvent extracts contain medicinally relevant bioactive components, justifying their usage in traditional medicines for the treatment of various ailments.

**Key Words:** Qualitative phytochemical analysis, *Manilkara zapota*, *M. hexandra*, *Madhuca indica*, *Mangifera indica*, *Annona squamosa*.

### **Introduction:**

Phytochemicals are non-nutritive substances. Chemical compounds are produced naturally on plants during metabolic processes and contain a wide range of proactive and disease-preventive qualities. These compounds are known to be produced by plants in order to defend themselves. Recent study has shown that they can also play a vital role in illness prevention in humans. Even some of these herbs have been used for millennia as traditional medicine. Most phytochemicals, such as flavonoids, carotenoids, and polyphenols, have antibacterial action and can be used to treat infections. The following plant species were studied: *Manilkara zapota* (L.) van Royen, *M. hexandra* (Roxb.) Dub., *Mangifera indica* L., *Madhuca indica* J.F. Gmel., and *Annona squamosa* L.

Plants play a universal function in the treatment of sickness, as seen by their use in every major system of medicine, regardless of philosophical foundation. Plants are vital to the pharmaceutical industry because they are a rich supply of pharmaceuticals and a vast reservoir of chemical variety for drug development screening procedures. The majority of the medications mentioned in the Indian medicinal system are derived from plants.



Medicinal plants have been used to treat disease all across the world for thousands of years. It is generally understood that some plants with active chemicals can limit microbial development. Plants' ability to generate chemicals via secondary metabolism determines their antimicrobial potency. Secondary metabolites emerged as the most important class of chemicals, with a wide spectrum of antibacterial and antifungal properties. When compared to conventional fungicides used to inhibit microbial growth and survival, these plant chemicals have different structures and activities.

Plants are a major source of Traditional medicine and can be used to cure a variety of conditions with little adverse effects. Traditional medical practises are not only beneficial in the treatment of diseases, but they also aid in the discovery of pharmaceutically active compounds in plants, which can aid in the commercial manufacture of pharmaceuticals. From ancient times, the importance of plant diversification in health care has been extensively recognised. According to a literature review, more than 50,000 plant species have been successfully employed for medical purposes globally, with flowering plants accounting for over 13% of these.

### **Material and methods:**

#### ***Plant collection and Identification***

*M. zapota* (L.) van Royen, *M. hexandra* (Roxb.) Dub., *M. indica* L., *M. indica* J.F. Gmel., and *A. squamosa* L. were among the plant species studied. The plants were harvested from the land in the Talod taluka in North Gujarat, India. The washed plant leaves were kept in for drying after being cleaned with tap water around 2-3 times to evaporate the water content. With the use of a mechanical blender, the sample was ground into a fine powder after drying. The powder is then kept in an airtight plastic container for future usage with adequate labelling.

#### ***Extraction technique***

Extraction is the separation of inert plant tissue constituents from medicinally active plant tissue constituents using a conventional extraction process. Menstrum is a selective solvent that is used to eliminate inert material and to obtain the curative part of the procedure through therapy.

***Method and Plant extraction:******Solvent extraction***

The crude plant extract was made using the Soxhlet extraction method. 10 gm of powdered plant material was placed into a thimble, and 300 ml of solvents were extracted separately. Acetone and chloroform were utilized as solvents. In a syphon tube of an extractor, the extraction process continued for 24 hours until the solvent became colourless. The extract was then placed in a beaker. The extract was then retained and cooked on a hot plate at 30- 40 °C until all of the solvent had evaporated. The dried extract was stored at 4°C in a refrigerator for future phytochemical investigation.

***Methods of phytochemical analysis:******Alkaloid***

**Wagner's test:** A few drops of Wagner's reagent were applied to 2mg of extract that had been acidified with 1.5 percent v/v hydrochloric acid. The presence of alkaloids is indicated by a yellow or brown ppt.

***Carbohydrates***

**Molisch's test:** 2mg of ethanolic extract was mixed with 10ml water, filtered, and concentrated. 2ml of conc. sulphuric acid was added to these 2 drops of freshly prepared 20% alcoholic alpha-naphthol solution, forming a layer below the mixture red-violet ring, showing the existence of carbohydrates that disappears when sufficient alkali is added.

***Amino acid***

**Ninhydrin test:** Boil for a few minutes 2ml Ninhydrin reagent + 2ml extract. The production of blue colour indicates the presence of amino acids.

***Steroids***

**Salkowski reaction:** 2mg of dry extract was combined with chloroform, to the chloroform layer sulphuric acid was gently introduced by the sides of the test tube. The emergence of a red colour indicated the presence of steroids.

**Tannin:** A few drops of a 5 percent w/v FeCl<sub>3</sub> solution were added to 1-2 ml of the ethanolic extract. Gallo tannins are shown by a green colour, whereas pseudo tannins are indicated by a brown colour.

**Flavonoids:** After mixing 2ml of each extract with a few drops of 20% sodium hydroxide, a bright yellow colour was seen. A few drops of 70% dilute hydrochloric acid were added to



this, and the yellow coloration disappeared. The presence of flavonoids in the sample extract is shown by the formation and disappearance of yellow colour.

**Saponins:** 6ml distilled water was added to 2 ml of each extract and rapidly shaken; the presence of saponin is indicated by the production of bubbles or persistent foam.

**Proteins:** adding 1 ml of 40 percent sodium hydroxide and a few drops of 1 percent copper sulphate to 2ml of each extract. The production of violet colour shows the presence of peptide linkage molecules in the sample extract.

**Glycosides:** 0.5ml of glacial acetic acid and 3 drops of 1% aqueous ferric chloride solution added in to 1ml of each extract. The creation of a brown ring at the interface shows the presence of glycosides in the sample extract.

**Terpenoids:** 1ml of each solvent is mixed with 0.5ml chloroform and a few drops of strong sulphuric acid to produce a reddish-brown precipitate that confirms the presence of Terpenoid in the extract.

**Table 1: Preliminary Phytochemical analysis**

Class of compounds	<i>Manilkara zapota</i>			<i>Manilkara hexandra</i>			<i>Mangifera indica</i>			<i>Madhuka indica</i>			<i>Annona squamosa</i>		
	AE	CE	WE	AE	CE	WE	AE	CE	WE	AE	CE	WE	AE	CE	WE
<b>Alkaloid</b>	+	+	-	+	-	-	+	+	-	+	-	-	+	-	-
<b>Amino acid</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Carbohydrate</b>	+	+	-	+	-	+	+	-	-	-	-	-	-	-	-
<b>Protein</b>	-	-	+	+	-	+	+	-	+	-	-	-	-	-	+
<b>Glycoside</b>	+	-	-	+	-	+	-	-	-	-	-	+	-	-	-
<b>Tannin</b>	-	-	+	-	-	-	+	+	+	+	-	+	+	+	+
<b>Terpenoid</b>	-	+	-	-	-	-	-	-	-	-	-	+	-	-	-
<b>Saponin</b>	-	+	+	+	-	+	+	-	+	+	-	+	+	-	-
<b>Flavonoid</b>	-	+	+	-	-	+	-	-	+	-	-	+	+	-	-
<b>Steroid</b>	+	-	+	-	-	+	-	-	+	-	-	-	-	-	-

Where, '+' = positive and '-' = negative

AE= Acetone extract, CE= Chloroform extract, WE= Water extract

Alkaloid, carbohydrate, glycoside, and steroid are all found in Acetone extract of *M. zapota* leaves. Chloroform extract of *M. zapota* leaves contains alkaloid, carbohydrate, terpenoid,



saponin, and flavonoid. Protein, Tannin, Saponin, Flavonoid, and Steroid can be found in the aqueous extract of *M. zapota* leaves., alkaloid, carbohydrate, protein, glycoside, and saponin are all found in Acetone extract of *M. hexandra* leaves. All phytochemicals are missing in the chloroform extract of *M. hexandra* leaves. Aqueous extract of *M. hexandra* leaves contains carbohydrate, protein, glycoside, saponin, flavonoid, and steroid. Alkaloid, Carbohydrate, Protein, Tannin, and Saponin are all found in Acetone extract of *M. indica* leaves. Alkaloid and Tannin are found in Chloroform extract of *M. indica* leaves. Protein, Tannin, Saponin, Flavonoid, and Steroid are found in aqueous extract of *M. indica* leaves. Alkaloid and Tannin are found in Acetone extract of *M. indica* leaves. Chloroform extract of *M. indica* leaves is devoid of all phytochemicals. Aqueous extract of *M. indica* leaves contains Glycoside, Tannin, Terpenoid, Saponin, and Flavonoid. Alkaloid, Tannin, Saponin, and Flavonoid are all found in Acetone extract of *A. squamosa* leaves. Aqueous extract of *A. squamosa* leaves contains protein and tannin.

### Discussion:

Plant samples were subjected to phytochemical analysis, which confirmed the existence of elements with medicinal and physiological properties. Phytochemicals such as carbohydrate, amino acid, protein, tannin, flavonoid, saponin, glycoside, steroid, terpenoid, and alkaloids were found in the plant extracts.

Tannins attach to proline-rich proteins, preventing them from being synthesized. Flavonoids are hydroxylated phenolic compounds that plants produce in response to microbial infection and have been discovered to have antibacterial properties in vitro against a wide range of pathogens. Their capacity to combine with extracellular and soluble proteins, as well as the bacterial cell wall, is most likely the reason for their activity. They are also powerful antioxidants with anticancer properties.

Saponins, which are known to have an anti-inflammatory impact, were also discovered in the plant extracts. Saponins have the ability to coagulate and precipitate red blood cells. Saponins are known for their ability to produce foams in aqueous solutions, as well as their hemolytic activity, cholesterol binding capabilities, and bitterness. Steroids have been shown to have antimicrobial characteristics, and they are extremely essential molecules, particularly in regard to sex hormones. For millennia, alkaloids have been associated with medical purposes, and cytotoxicity is one of their most prevalent biological features. Alkaloids have been shown



to have analgesic, antispasmodic, and antibacterial effects by several researchers. Glycosides have been shown to reduce blood pressure in numerous studies. The findings of this study indicate that the detected phytochemical substances may be bioactive constituents, and that these plants are proving to be a valuable reservoir of bioactive compounds with significant medical value.

### **Conclusion:**

The findings demonstrated that the plants investigated contained medicinally essential components. Many previous investigations had accumulated evidence that the discovered phytochemical was bioactive. Several studies have demonstrated that these phytochemicals contribute pharmacological and physiological qualities to the plants researched in the treatment of various illnesses. As a result, extracts from these plants could be considered a valuable source of drugs. Further research should be carried out to separate, purify, and characterize the active ingredients responsible for the activity of these plants, as well as further work to isolate, purify, and characterize the active constituents responsible for the action of these Plants. In addition, more research into the putative mechanisms of action of these extracts is urged.

### **Acknowledgement:**

Authors are thankful to Disha life sciences Pvt. Ltd. for practical work during this study.

### **References:**

- Ashwin Godghate, Rajaram Sawant, Ashok Sutar (2012) Phytochemical Analysis of Ethanolic extract of Roots of *Carrisa carandus* Linn., *Rasayan.J.Chem.* 5(4):456-459.
- K. Girija, P. Udaya Chandrika, K. Lakshman, N. Pruthvi (2010) Evaluation of Antimicrobial activity of various Bark extracts of *Bombax malabaricum.*, *Int.J.Res.Pharm.Sci.*, 1(2):199-204.
- Khalid S, Shahzad A, Bashrat N, Abubakar M, Anwar P (2018) Phytochemical screening and Analysis of selected Medicinal Plants in Gujarat., *J. Phytochemistry Biochem.* 2:108.
- Kodangala C, Saha S, Kodangala P (2010) Phytochemical studies of Aerial parts of the Plant *Leucas lavandulaefolia.*, *Der Pharma Chemica*, 2(5):434-437.



- Manash PS et al. (2016) Phytochemical Analysis of Traditional Medicinal Plants and their Antimicrobial activity: An Experience from North East India. J Pharm Res, 1(1):000104.
- Prabhavathi R. M, Prasad M. P and Jayaramu M (2016) Studies on Qualitative and Quantitative Phytochemical Analysis of *Cissus quadrangularis*., Advances in Applied Science Research. 7(4):11-17.
- Pushendra K Patel, Narendra K Prajapati and B.K. Dubey (2012) *Madhuka indica*: A Review of its Medicinal property, IJPSR, 3(5): 942-950.
- RNS Yadav and Munia Agrawala (2011) Phytochemical analysis of some medicinal plants, Journal of Phytology, 3(1):10-14.
- S. Hemadri Reddy, Aisha Khalid AL-Hinai, Hajer Hafed AL-Yaqoobi and Fatima Jafar AL-Ajmi (2016) Phytochemical analysis, antimicrobial screening and Hypoglycemic effect of some selected Medicinal plant extract from Oman., Journal of Experimental Biology and Agricultural Sciences, 4(2):218-224.

---

**Suggested citation:**

Panchal M and Jha C (2021). *Qualitative phytochemical analysis of some medicinal plants (leaves) from the surrounding area of Talod Taluka, North Gujarat, India. Prithivya, An Official Newsletter of WCB Research Foundation and WCB Research Lab. Vol 1(2) 37-43.*





## M. I. Patel Award for Young Wildlife & Conservation Biologist

### Nominations for 2021 Award

#### Ms. Shivangi Mishra



*Shivangi is a Senior Research Fellow (Basic Scientific Research Fellow, UGC) at University of Lucknow and is currently working on Egyptian Vultures. She has also worked on Butterflies of Uttar Pradesh from 2014- 2016. She has presented her research on Egyptian Vultures in Conference of Raptor Research Foundation in 2018 (South Africa) 2019 (USA) and awarded with William C Andersen Award and James Koplun Award respectively. In future also she wants to contribute her research for the conservation of Egyptian Vultures in India.*

#### Dr. Tarang Kumar Shah



*Tarang is an Assistant Professor in Fisheries polytechnic at Dau Shri Vasudev Chandrakar Kamdhenu Vishwa Vidyalaya, Durg. He studied the Effect of Thymus linearis plant Extract on Growth Performance and Non-Specific Immune Responses of endangered Golden Mahseer (Tor putitora) against the disease caused by Oomycetes Saprolegnia parasitica. He found that the leaf extract of Thymus plant has the antifungal properties and a small quantity of which can control the population against saprolegniosis.*

#### Dr. Sujit Kolangath



*Sujit is a subject specialist in animal biotechnology at Wildlife Research & Training Center, Gorewada, Nagpur. He studied the qualitative and quantitative aspect of hepatozoon felis epidemiology in a tiger cub. He reported that the infection is transmitted by ingestion of infected ticks, infected prey and carrion and concurrent infection with bacterial or viral infections may be fatal to the infected animal.*

### **Ms. Jis Sebastian**



*Jis recently completed her doctoral research in Botany from Madurai Kamraj University, Madurai- Tamil Nadu. She is studying the pattern of diversity, spatial distribution and community structure of epiphytic orchids of southern Western Ghats responds to availability of microenvironments (Microhabitat supported by Microclimates) in the structurally diverse trees. There for Conservation of epiphytic orchids would ensure conservation of old aged, natural, structurally complex trees in the forest ecosystem.*

### **Dr. Harshil Patel**




*Harshil is a naturalist working mainly on systematics and taxonomy of reptiles and amphibians. He studied the amphibians & reptiles of northern Western Ghats of Gujarat. He found a new, Miocene divergent, endemic lineage of a racer snake; evolutionary history of an endemic snake resulted in the resurrection of a genus; reinvestigation of a gecko species complex that resulted into the discovery of two new species.*

*The final round including the presentation by the nominees followed by the award ceremony will be conducted on September 27, 2021 at M N College, Visnagar (Gujarat). WCB and the M I Patel Award committee congratulate all the nominees and wishing all the best for their presentation.*

*Applications and nomination for the 2022 awards are now open through our website ([www.wcbresearch.in](http://www.wcbresearch.in)) till June 30, 2022.*

## PROF. M. I. PATEL AWARD



Award for Young Wildlife & Conservation Biologist

The Award will be conferred to any young scientist / student/ researcher below the age of 35 years working in the field of wildlife and conservation biology (Zoology/ Botany/ Environmental Science/ or allied subject) and carried out significant research in the respective field.

How to apply:

Register online with an extended research abstract in 1000 words and the outstanding contribution to the wildlife conservation along with your one page CV and one letter of nomination/ recommendation. The registration fees is Rs. 200 (non refundable)

Selection criteria:

A selection committee will review your application and invited you for the online presentation before the Jury Panel and the best five researchers will be selected for grand finale.

### *Announcement*

As United nation declared this decade (2021-2030) an Ecosystem Restoration Decade; We, WCB Research Foundation with CNEW are organizing a field based workshop and hands-on training on

### **Ecological Restoration: Principles and Field applications**

A four days training course in **October 2021** with very limited seats. A renowned, nationally and internationally recognized eco-restoration experts will be providing the hands-on training and sharing their experiences.

An ecology student, researcher, faculty member, policy makers, industrialist, architecture, civil engineers should participate in this field based workshop.

**Keep looking at our website for date and registration details**

[www.wcbresearch.in](http://www.wcbresearch.in)

**Write us for more details:**

[wcbresearch@gmail.com](mailto:wcbresearch@gmail.com) | [cnew.1099@gmail.com](mailto:cnew.1099@gmail.com)

## Internship Opportunities @ WCB Research Foundation

We, at WCB Invites applications for internship, training and dissertations in hybrid mode (Online and Offline) under our team of experts in the following areas



*Adding Science to Conservation*

For more details contact to

Director of Research:  
8329519903

Admin Officer:  
9712313698

Email:  
[wcbresearch@gmail.com](mailto:wcbresearch@gmail.com)  
Apply online at  
[www.wcbresearch.in](http://www.wcbresearch.in)

- **Research methodology**

- **Wildlife Biology**

- Mammalian ecology
- Human-wildlife interaction
- Avifaunal diversity
- Herpetology
- Marine biology
- Invertebrate diversity

- **Environment & Ecology**

- Biodiversity & Ecosystem Assessment
- Eco-restoration
- Urban Environment
- Wetland Conservation
- Solid-waste Management

- **Forestry & Plant biology**

- Plant diversity
- Ethnobotany
- Habitat Assessment

- **Geo-Spatial Analysis**

- **Biotechnology & Conservation Genetics**

- Bio and Eco Informatics
- Environmental Biotechnology

WCB Research Foundation with K J Somaiya College, Mumbai have organized Digital and Handmade Posters Competition on the occasion of World Mangrove Day July 26, 2021. Out of entries of thirty participants three best posters were selected by the committee and awarded.

## Posters of Winners

### MANGROVES

a tree with roots which are above the ground and that grows along coasts of large rivers



# 30-50%

of mangrove forest have been lost due to shrimp farming, intrusive coastal development, tourism and other factors.

#### Threats of mangroves



DEFORESTATION



POLLUTION



CLIMATE CHANGE

#### Benefits of Mangroves



protection from coastal erosion



carbon sequestration




controls flooding



water quality improvements



biodiversity



mangroves are home of many Endangered species

#### • Why Should We Protect Mangroves?

They perform important ecological functions like nutrient cycle, hydrological regime, coastal protection, fish-fauna productions, etc. They reduce high tides and waves help prevent soil erosion.

more than 50 species of mangroves found in the world

*1<sup>st</sup> Prize Digital Poster Ms. Yukta Sawant*

## MANGROVE CONSERVATION / IMPORTANCE OF MANGROVE

A tidal habitat comprised of salt-tolerant trees & shrubs.

Prop roots, Pneumatophores which helps them to impede water flow & thereby enhance the deposition of sediments in areas

### IMPORTANCE

- Moderate Monsoonal floods & reduce inundation of inland low lands.
- Protect coastal areas from Tsunami, hurricanes & floods.
- Enhance the natural recycling of nutrients.
- Support & flourish various & wild life.
- Provide woods, firewood, medicinal plants, shrimps, lobsters, fish, shell, honey to local communities.
- Mangroves store more oil than most of the other forests.

### REASON OF DECLINE

- Over-exploitation of Timber, fish, shellfish, crustaceans, Aquaculture, Saltpans.
- **URBANIZATION**
- Deforestation
- To construct continuous landmass
- Blue carbon emissions are being artificially augmented through mangrove deforestation.

### SIGNIFICANCE OF FUNCTION

- Important nursery grounds & breeding sites for birds, fish, crustaceans, shellfish, reptiles & mammals.
- Renewable source of wood.
- Accumulation sites for sediments, contaminants, carbon & nutrients.
- Offer protection against coastal erosion.

### FUTURE OF MANGROVE / STRATEGIES FOR CONSERVATION

- Depends on
- Technological and ecological advances
- Multipurpose silviculture
- Genetics
- Forestry modelling
- Rehabilitation

### HEAVY METAL DISTRIBUTION AND ACCUMULATION FROM HISTORICAL AND ANTHROPOGENIC SOURCES IN TROPICAL MANGROVES OF INDIA AND BANGLADESH

Location	Lead	Cadmium	Copper	Chromium	Zinc	Iron	Manganese	Nickel	Vanadium
Chennai	0.0015	0.0001	0.0005	0.0002	0.0010	0.0020	0.0005	0.0001	0.0001
Coimbatore	0.0012	0.0001	0.0004	0.0002	0.0008	0.0015	0.0004	0.0001	0.0001
Hyderabad	0.0018	0.0001	0.0006	0.0003	0.0012	0.0025	0.0006	0.0001	0.0001
Kolkata	0.0020	0.0001	0.0007	0.0004	0.0015	0.0030	0.0008	0.0001	0.0001
Madurai	0.0010	0.0001	0.0003	0.0001	0.0005	0.0010	0.0003	0.0001	0.0001
Thiruvananthapuram	0.0014	0.0001	0.0005	0.0002	0.0009	0.0018	0.0005	0.0001	0.0001
Visakhapatnam	0.0016	0.0001	0.0006	0.0003	0.0011	0.0022	0.0006	0.0001	0.0001

### TYPES OF MANGROVE

Red Mangrove (*Rhizophora* sp.)  
 Black Mangrove (*Avicennia* sp.)  
 White Mangrove (*Laguncularia* sp.)

Mangrove provides: Ecological niches & Interconnectivity of the marine & land ecosystems

**SOMAIYA VIDYAPEETH**

TANYA SAWARDEKAR | ZOOLOGY | Msc - Part 1 | K. J. Somaiya College

1<sup>st</sup> Prize Digital Poster Ms. Yukta Sawant

## MANGROVES: Living On The Edge ...

**WATER PURIFICATION**

**CARBON STORAGE**

**COASTAL PROTECTION**

**PROTECTS FROM CLIMATE CHANGE**

1<sup>st</sup> Prize Handmade Poster Ms. Tanya Sawardekar



## Governing Council of WCB Research Foundation

<b>Executive Director</b>	<b>Dr. Hardik Patel</b>	Assistant Professor Parul University, Vadodara <a href="mailto:p.hardik80@gmail.com">p.hardik80@gmail.com</a>
<b>Director of Research</b>	<b>Dr. Nishith Dharaiya</b>	Associate Professor HNG University, Patan <a href="mailto:nadharaiya@ngu.ac.in">nadharaiya@ngu.ac.in</a>
<b>Member</b>	<b>Dr. Sagarika Damle</b>	Associate Professor & Head Department of Life Sciences, KC College, Mumbai <a href="mailto:sagarika.damle@kccollege.edu.in">sagarika.damle@kccollege.edu.in</a>
<b>Member</b>	<b>Dr. Nuria Selva</b>	Associate Professor Institute of Nature Conservation Polish Academy of Science, Krakow, Poland <a href="mailto:nuriaselva@gmail.com">nuriaselva@gmail.com</a>
<b>Member</b>	<b>Mr. Pratik Desai</b>	Jr. Research Fellow WCB Research Lab, Patan <a href="mailto:pratikdesai825@gmail.com">pratikdesai825@gmail.com</a>
<b>Member</b>	<b>Mr. M. S. Patel</b>	Associate Professor M N College, Visnagar <a href="mailto:martinleems@gmail.com">martinleems@gmail.com</a>
<b>Member</b>	<b>Dr. Haresh Gondaliya</b>	Assistant Professor Shri G D Modi Science College, Palanpur <a href="mailto:haresh.gondliya55@gmail.com">haresh.gondliya55@gmail.com</a>
<b>Member</b>	<b>Mr. Prasad Patil</b>	Range Forest Officer Gujarat Forest Department <a href="mailto:jalu1890@gmail.com">jalu1890@gmail.com</a>
<b>Member</b>	<b>Ms. Madhuri Padaya</b>	Assistant Professor K J Somaiya College of Science, Mumbai <a href="mailto:padayamadhuri@yahoo.com">padayamadhuri@yahoo.com</a>
<b>Member</b>	<b>Dr. Jagruti Rathod</b>	Director, Center for Nature & Ecosystem Wellbeing, Vadodara <a href="mailto:rathod21@gmail.com">rathod21@gmail.com</a>
<b>Member Secretary</b>	<b>Ms. Jahnvi Patel</b>	Admin Officer, WCB Research Foundation, Patan <a href="mailto:admin@wcbresearch.in">admin@wcbresearch.in</a>



## **Instructions for authors**

**Scope:** Pruthivya is a tri-annual newsletter of WCB Research Foundation and will publish manuscripts on wildlife, conservation biology and its allied areas.

## **Guidelines to contributors of Prithiviya**

- *Manuscript Types:* Prithiviya publishes, original research articles, short notes, field notes, reviews and comments. Submissions should be in English (UK) only.
- All manuscripts should be easy to read and comprehend. Manuscripts may be edited for length, content and style, and be sent to referees when found to be appropriate. The editor will discuss contributions with authors and advise to modified whenever required.
- Submission is considered on the condition that papers are previously unpublished, have not been offered simultaneously elsewhere, and that all contributors have read and approved the contents.
- The editor will acknowledge manuscript contributions and assign them a unique registration number, which latter must be quoted in correspondence. Papers are peer-reviewed, although the editor reserves the right to reject one without a review.

## **Preparation and submission of manuscripts**

1. Title page:
  - Add full title,
  - Author/s name and affiliation with email of all the Authors
  - Abstract (250 Words) and Key words (4-5)
2. Submitted manuscripts should generally follow the normal convention of scientific manuscripts, keywords, generally not appearing in the title, Introduction, Materials and Methods, Results and Discussion, Acknowledgments, Authors contribution (optional), Appendices (optional) and References) with the replacement or addition of sections allowed in between the Introduction and the Acknowledgments.
3. The manuscript should be typed in MS Word with left-aligned typesetting and only true type fonts (Times new roman preferable) should be used. The line spacing must be 1.5 and the font size of 12. Headings should be bold and sub-headings should be made italicized.





4. **Title:** A concise, informative and attractive and short title with 5-20 words will be encouraged
5. **Abstract:** Word limit for the abstract is 250 words and should contain a summary of all major findings or conclusions of the study; however abstract is not required in case of short notes, field notes and other short communication.
6. **Introduction:** should be clearly written and must covering the objectives and significance of the study.
7. **Methodology:** A detailed methodology is always encouraged. The methodology should contain the study area (if applicable), data collection, analysis and interpretation. Authors should also mention the instruments, software used for the experiments and analysis.
8. **Result and discussions:** which contain main result, with proper references and authentic justifications include, graph, table, photographs (No checklist, it directly goes into appendix or supplementary data).
9. **Conclusions:** A short and comprehensive conclusion with significant /novel findings and depicting your research in a nut-shell.
10. **Acknowledgments:** which contain all funds support, persons who help during project or research or findings, name of university other than they work etc.
11. Checklist-based papers, like trip reports, should contain dates of the trip, total number of birders or participants, Mammals, reptiles, amphibians, invertebrates etc.
12. Manuscripts should be sent electronically as an email attachment to the Editorial Assistant with a cover letter to .....
13. The text, tables, figure/photograph captions (which must be self-explanatory), and appendices should be combined in one Microsoft Word® (“.doc,” OR “.docx”) file format. The preferred font is ‘Times New Roman’ in 12 point.
14. Photographs, artwork, maps (the outlines of the Republic of India should conform to Government of India regulations), diagrams, *etc.*, should be digitized and sent separately along with the manuscript, or if they are large



files, in separate, trailing e-mails. These should be in TIFF and at least 8×11 inches in 300 dpi resolutions. JPEG files must be ‘maximum’ quality, i.e., at their minimum compression. Maps should be marked with a scale and North arrow, and should carry a clear caption.

15. Authors, whose work involves the handling of any necessary permissions from the requisite government agencies, to do so. The editor reserves the right to view such permissions at her/his discretion.
16. While photographing nesting/courtship display or any behavioural activity, including those of nests, eggs, courtship *etc.*, authors/photographers must ensure the safety of the animals involved, and also ensure that they do not disturb in any way whatsoever, in the course of their photography. The editor reserves the right to view permissions for such activities within protected areas. The editor/referee reserves the right to query an author regarding the ethics of photography, and the methodology followed while photographing including that of rejecting a manuscript if unsatisfied on these counts.

### **Types of Manuscript**

1. Short Research articles: 2000 to 2500 words
2. Short notes: Up to 1500 words
3. Important sightings: 250-350 words with proper GPS location and photographic evidence
4. Important findings: 250-500 words with proper GPS location with image
5. Full-length Research paper: Up to 10000 words with appendix excluding photographs only.

### **Other important Guidelines**

- Scientific binomials should comprise the generic and specific names, but should not include the name of the author and year of citation. Scientific trinomials should be mentioned only if the nature of the manuscript requires them.
- For example, both the English and scientific name must be given, thus, “House Crow *Corvus splendens*”, thereafter, only the English name, “House Crow” will suffice. English and scientific names should follow the proper book, that should be mention in reference and title of the table



- Metric units and their international symbols must be used, e.g., “sq. km”, and “1,000 m asl.”; dates and times should be of the form of MMDDYYYY (January 01, 2021”, and “1345 h” respectively.
- Tables should be within gridlines (which may or may not be printed). Tables should never be formatted with wrap-around text.
- Latitude and longitude coordinates should follow the format, “27.807°N, 74.094°E”.
- Enzyme Nomenclature: The trivial names recommended by the IUPAC-IUB Commission should be used. When the enzyme is the main subject of a paper, its code number and systematic name should be stated in its first citation in the paper.
- Symbols and Abbreviations: Use only standard abbreviations. The use of non-standard abbreviations can be extremely confusing to readers. Avoid abbreviations in the title. The full term for which an abbreviation stands should precede its first use in the text unless it is a standard unit of measurement.
- Page number (page 1 Of 10 formats) must be on top right corner of the word documents, font size 10 and times new roman word type.

### **Citation and References**

Pruthivya encourages the authors to use Mendely, EndNote or other reference management tools for proper citations. We follow APA referencing style.

- Do not abbreviate references; pagination should be given in the expanded form, e.g., “i-vi, 1-250”, not “vi, 250”; the page numbers should be separated by an ‘En’ dash.
- Italics should be used for book titles but not volume numbers; when citing periodicals use italics for the journal name, not the article title.
- References should be arranged alphabetically by the first author (and in the case of more than one author, by second and subsequent authors).
- Citations to references in the text should follow the style, “(Santharam 1978)”, “(Barua & Sharma 1999)”, “(Pande et al. 2002)”, “(Lainer 1999, 2004)”, “(Lainer 1999a,b,c,d; Lainer 1999a, 2000a)” [for several papers by an author/s from



a/multiple calendar year/s], and “(Neelakantan et al. 1993; Sashikumar et al. 2011)”.

- A string of citations in the text should follow a chronological, not an alphabetical order, thus: “(Ali & Ripley **1981**; Kazmierczak **2000**; Grimmett et al. **2011**)”.
- When citing unpublished sources, please use the following formats: if the source is written material, “(Harkirat Sangha, *in litt.*, e-mail/letter dated 02 January 2013)”; if oral, “(Rajah Jayapal, *verbally*, dated 15 December 2013)”.
- When citing from e-Bird or any other magazine checklist, please enter the author of the e-Bird checklist’s name or others name, and the year the checklist was uploaded, in the text, “(Raman 1994)”. If a checklist that’s shared amongst multiple observers is being referred to, only the relevant checklist(s) should be used for citation.

See examples below for detailed formatting:

#### Journal articles

- Pittie, A., 2011. The dates of seven new taxa described by W. E. Brooks (*Certhia Hodgsoni*, *Sitta Cashmirensis*, *Dumeticola major*, *Horites [sic] Pallidus*, *Phylloscopus Tytleri*, *Motacilla Cashmirensis*, and *Alauda Guttata*). *Pruthivya*7 (2): 54–55.
- Naoroji, R., & Sangha, H. S., 2011. Threats to habitat and wildlife in Changthang and Rupshu areas of Ladakh: a case study at Hanle. *Pruthivya*7 (1): 2–6.
- Choudhary, D. N., Mandal, J. N., Mishra, A., & Ghosh, T. K., 2010. First ever breeding record of Black-necked Stork *Ephippiorhynchus asiaticus* from Bihar. *Pruthivya*6 (3): 80–82.

#### Books

- Futehally, Z. (ed.) 2006. *India through its birds*. 1st ed. Bangalore, India: Dronequill Publishers Pvt. Ltd. Pp. i–ii, 1–214.
- Pittie, A., 2010. *Birds in books: three hundred years of South Asian ornithology—a bibliography*. 1st ed. Ranikhet: Permanent Black. Pp. i–xxi, 1–845.
- Sashikumar, C., Praveen J., Palot, M. J., & Nameer, P. O., 2011. *Birds of Kerala: status and distribution*. 1st ed. Kottayam, Kerala: DC Books. Pp. 1–835.

#### Book chapter

- Pittie, A., 2011. Stray Feathers (1872–1899) (p. 247). In: *Priority! The dating of scientific names in ornithology: a directory to the literature and its*



reviewers. Dickinson, E. C., Overstreet, L. K., Dowsett, R. J., & Bruce, M. D. (eds.). Northampton, UK: Aves Press Limited.

#### Website

- 2013. Kadalundi makes history with new gull species. *The Hindu* (Thiruvananthapuram ed.) dated February 7, 2013. Website: <http://www.thehindu.com/todays-paper/tp-national/tp-kerala/kadalundi-makes-history-with-new-gull-species/article4388171.ece>. [Accessed on 21 July 2014.]

#### Any magazine checklist

- Raman, T. R. S., 1994. Website URL: <http://ebird.org/ebird/view/checklist?subID=S21694571>. [Accessed on 15 December 2015.]

#### **ETHICAL MATTERS**

Authors involving in the usage of experimental animals and human subjects in their research work should seek approval from the appropriate Institutional Animal Ethics Committee in accordance with "Principles of Laboratory Animal Care". The material and methods section of the manuscript should include a statement to prove that the investigation was approved and that informed consent was obtained.

#### **Disclaimer**

- While every care is taken of manuscripts and their accompanying material, *Pruthiviya* cannot be held responsible for accidental loss or damage of any material sent for publication or for their return, whether they are accepted for publication or not.
- Material published in *Pruthiviya* reflects the views of the authors and not necessarily those of the publishers, editors, referees, etc.
- The editor reserves the rights to make necessary changes in manuscripts in consultation with the author.

All submissions are evaluated under the assumption that the conditions listed here have been understood and accepted by the author(s).

**Submission fees:** There will be no submission fees for e-publication; however, we may produce the reprints on request and on payment basis.



**Special announcements**

We also accept announcement of workshop, webinar, conference, symposium, job opportunities, academic courses, etc. with nominal charges and guarantee for its wider circulation.

**Contacts**

**Editor in Chief**

: Dr. Sagarika Damle  
Email: [visaanika@gmail.com](mailto:visaanika@gmail.com)

**Associate Editors**

: Dr. Nishith Dharaiya  
Email: [nadharaiya@gmail.com](mailto:nadharaiya@gmail.com)

: Dr. Jagruti Rathod  
Email: [rathod21@gmail.com](mailto:rathod21@gmail.com)

**Editorial Assistant**

: Ms. Rinku Bhati  
Email: [bhatirinku178@gmail.com](mailto:bhatirinku178@gmail.com)



**WCB Research Foundation**

*Adding Science to Conservation*

आत्मवत् सर्व भूतेषु

