

Psychiatric Rehabilitation Journal

Text Message Exchanges Between Older Adults With Serious Mental Illness and Older Certified Peer Specialists in a Smartphone-Supported Self-Management Intervention

Karen L. Fortuna, John A. Naslund, Kelly A. Aschbrenner, Matthew C. Lohman, Marianne Storm, John A. Batsis, and Stephen J. Bartels

Online First Publication, July 16, 2018. <http://dx.doi.org/10.1037/prj0000305>

CITATION

Fortuna, K. L., Naslund, J. A., Aschbrenner, K. A., Lohman, M. C., Storm, M., Batsis, J. A., & Bartels, S. J. (2018, July 16). Text Message Exchanges Between Older Adults With Serious Mental Illness and Older Certified Peer Specialists in a Smartphone-Supported Self-Management Intervention. *Psychiatric Rehabilitation Journal*. Advance online publication. <http://dx.doi.org/10.1037/prj0000305>

Text Message Exchanges Between Older Adults With Serious Mental Illness and Older Certified Peer Specialists in a Smartphone-Supported Self-Management Intervention

Karen L. Fortuna

Dartmouth Centers for Health and Aging, Lebanon, New Hampshire, and CDC Health Promotion Research Center at Dartmouth, Lebanon, New Hampshire

John A. Naslund

Harvard Medical School

Kelly A. Aschbrenner

Dartmouth Centers for Health and Aging, Lebanon, New Hampshire, and CDC Health Promotion Research Center at Dartmouth, Lebanon, New Hampshire

Matthew C. Lohman

University of South Carolina

Marianne Storm

The Dartmouth Institute, Lebanon, New Hampshire, and University of Stavanger

John A. Batsis and Stephen J. Bartels

Dartmouth Centers for Health and Aging, Lebanon, New Hampshire, and CDC Health Promotion Research Center at Dartmouth, Lebanon, New Hampshire

Objective: To identify the strategies peer specialists use to provide illness self-management support for older adults with serious mental illness (SMI) through text messaging. **Method:** Transcripts of text message exchanges between 8 older adult participants with SMI who completed the PeerTECH intervention and 3 older adult certified peer specialists who delivered the 12-week program were analyzed. Text message analyses explored themes relevant to peer support and health behavior change. Quantitative data comprised frequency of text messages by either the peer or consumer. **Results:** Consumers ($N = 8$) had a mean age of 68.8 years ($SD = 4.9$) and were mainly women (88%), White (100%), and married (75%). Certified peer specialists ($N = 3$) were all 55 or older; 100% were female, 66% identified as White, and 33% identified as African American. Overall, peers sent 215 text messages whereas consumers sent 141 text messages. In the peer specialist–consumer text message exchanges, we identified 4 themes on different aspects of illness self-management, including health behavior change, self-management therapeutic techniques, engagement in health technology, and peer support. **Conclusions and Implications for Practice:** This exploratory qualitative study offers preliminary support that peers are able to use text messages to support the delivery of a peer-delivered home-based medical and psychiatric self-management intervention. Certified peer specialists can potentially provide a range of illness self-management support to older adults with SMI via text messaging. These findings will inform the

Karen L. Fortuna, Dartmouth Centers for Health and Aging, Lebanon, New Hampshire, and CDC Health Promotion Research Center at Dartmouth, Lebanon, New Hampshire; John A. Naslund, Department of Global Health and Social Medicine, Harvard Medical School; Kelly A. Aschbrenner, Dartmouth Centers for Health and Aging and CDC Health Promotion Research Center at Dartmouth; Matthew C. Lohman, Department of Epidemiology and Biostatistics, University of South Carolina; Marianne Storm, The Dartmouth Institute, Lebanon, New Hampshire, and Faculty of Health Sciences, University of Stavanger; John A. Batsis and Stephen J. Bartels, Dartmouth Centers for Health and Aging and CDC Health Promotion Research Center at Dartmouth.

Matthew C. Lohman is now at The Dartmouth Institute and Faculty of Health Sciences, University of Stavanger.

Dr. Batsis' research in this publication was supported in part by National Institute on Aging of the National Institute of Health under Award K23AG051681. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of

Health. Research reported in this publication was supported by the Dartmouth Clinical and Translational Science Institute, under award number UL1TR001086 from the National Center for Advancing Translational Sciences (NCTAS) of the National Institute of Health (NIH). The content is solely the responsibility of the author(s) and does not necessarily represent the official views of the NIH.

This study was funded by the Health Promotion Research Center at Dartmouth through a Centers for Disease Control and Prevention grant (Cooperative Agreement U48 DP005018). Additional support was provided by National Institute of Mental Health Grant T32 MH073553-11. Funding for the fifth author was received from the Norwegian Research Council–Harkness Fellowship in Health Care Policy and Practice, 2017–2018 (Project 276638). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the article.

Correspondence concerning this article should be addressed to Karen L. Fortuna, Dartmouth Centers for Health and Aging, 46 Centerra Parkway, Suite 200, Lebanon, NH 03766. E-mail: karen.l.fortuna@dartmouth.edu

development of standardized peer text-messaging services to augment evidence-based illness self-management interventions for older adults with SMI.

Impact and Implications

This study offers preliminary support that certified peer specialists' services can potentially be advanced with the use of text messaging to provide illness self-management support to older adults with a serious mental illness.

Keywords: serious mental illness, mHealth, peer support, illness self-management, text messaging

Since the deinstitutionalization of mental hospitals in the 1960s and the successive consumer movement led by former patients in the 1970s, peer support has been recognized as a valuable component of consumer recovery (Stroul, 1989). Peer support is not a form of therapy (Zinman, Harp, & Budd, 1987); rather, it is a nonmanualized form of social support that is provided by a person living with a mental health condition to others sharing a similar mental health condition to achieve individually identified goals (Solomon, 2004). Peer support interventions delivered in inpatient and outpatient settings have been shown to be feasible and acceptable, with high potential for clinical effectiveness in multiple domains, including hope, sense of personal control, ability to make positive changes, and decreased psychiatric symptoms (Chinman et al., 2014). Peer support services are typically provided through face-to-face interactions in clinical settings, the community, or the home (Chinman et al., 2014). However, the proliferation of mobile health (mHealth) in mental health services has created an opportunity to advance delivery of peer support services beyond traditional approaches. Researchers have used mHealth with peers to promote intervention fidelity and delivery of evidence-based practices and to increase intervention dose without in-person sessions (Fortuna et al., 2018). As researchers increasingly use mHealth to enhance peer services delivery, it will be important to understand how peers interact with technology to optimize peer-delivered interventions. In this study, we examined peers' text-messaging strategies to support a home-based integrated medical and psychiatric self-management intervention.

It is estimated that 72% to 93% of people with serious mental illness (SMI; people with schizophrenia spectrum disorders, bipolar disorder, or persistent major depressive disorder) own smartphones (Ben-Zeev, Davis, Kaiser, Krzysos, & Drake, 2013; Naslund, Aschbrenner, & Bartels, 2016). Mobile health interventions using smartphone apps afford opportunities to promote physical activity (Naslund, Aschbrenner, Barre, & Bartels, 2015) and assist people with SMI with psychiatric symptom monitoring (Naslund, Marsch, McHugo, & Bartels, 2015). Peers' role in mHealth services delivery is a largely unexplored service delivery approach. Our prior research has shown peer-delivered interventions enhanced with an app appear feasible and acceptable and are potentially associated with improvements in psychiatric self-management, self-efficacy for managing chronic health conditions, hope, quality of life, medical self-management skills, and empowerment among adults with SMI and chronic health conditions (Fortuna et al., 2018).

Approximately 78% of all mobile phone owners with SMI use text messaging (Naslund et al., 2016). Initial studies using

text messaging for people with SMI have focused on remote symptom monitoring (Španiel et al., 2008), improving functioning (Pijnenborg et al., 2010), delivering appointment reminders (Maiga, 2011), and supporting medication and treatment adherence (Beebe, Smith, & Phillips, 2014; Montes, Medina, Gomez-Beneyto, & Maurino, 2012; Sims et al., 2012). However, there has been little research on the practice of peer-to-consumer text messaging to support medical and psychiatric self-management among adults with SMI. mHealth could potentially provide support outside of structured clinical environments including real or automated social interactions that are tailored to an individual's goals and delivered through video and audio modalities (Whiteman, Lohman, & Bartels, 2017; Whiteman, Lohman, Gill, Bruce, & Bartels, 2017).

Despite the rapid progress of research in the emerging field of mHealth with traditional providers, few studies have incorporated mHealth in peer support services (Fortuna et al., 2018). Peers are viewed as role models who have a shared intimate "lived experience," and they offer access to embedded support networks in the community. Certified peer specialists have professional practice standards, including (a) sharing their personal stories of recovery, (b) facilitating learning among those they support, (c) learning from those they support, and (d) focusing on those they support and encouraging them to achieve what they want in life (International Association of Peer Support Providers, 2011). Unlike the case with traditional providers, these professional practice standards allow for flexibility in the content of intervention delivery (e.g., text message exchanges) within previously described parameters. Identifying and understanding the potential impact of strategies used by peer specialists to support home-based integrated medical and psychiatric self-management using text messaging can inform the development of peer-delivered mHealth interventions.

We explored the content of smartphone text message interactions between consumers with SMI and certified peer specialists who were engaged in a 12-week in-person peer-delivered intervention augmented by a smartphone app that incorporated a text-messaging feature that was compliant with the Health Insurance Portability and Accountability Act of 1996. The intervention targeted training adults with SMI to self-manage their medical and psychiatric illnesses through providing knowledge and skill development in psychoeducation, behavioral tailoring, coping skills training, and motivational interviewing while also allowing consumers to determine and work toward their own self-management goals at their own pace. Findings pertaining to the feasibility, acceptability, and preliminary effectiveness of this peer-delivered

mHealth self-management intervention have been reported elsewhere (Fortuna et al., 2018). The present study seeks to expand on this prior work by identifying strategies that certified peer specialists use to support a home-based integrated medical and psychiatric self-management intervention for older adults with SMI.

Method

Study Design and Participants

The study design and recruitment procedures for PeerTECH have been described in a previously published article (Fortuna et al., 2018). Briefly, 10 adults with SMI and medical comorbidity (i.e., cardiovascular disease, obesity, diabetes, chronic obstructive pulmonary disease, hypertension, and/or high cholesterol) ages 60 years and older received the PeerTECH intervention in their homes. Eight of these adults completed the intervention and were included in the current analysis. PeerTECH was delivered by three certified peer specialists augmented with a smartphone application over a period of 12 weeks (Whiteman, Lohman, & Bartels, 2017; Whiteman, Lohman, Gill, et al., 2017).

All procedures were conducted in accordance with the ethical standards of the (World Medical Association, 2013) Institutional Review Board Dartmouth College and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. Written informed consent was obtained from all participants.

Peer Training

Training of peers was provided over 4 days by the first author and included (a) information about physical and mental health, (b) techniques used in PeerTECH, (c) setting goals and actions steps to achieve goal, (d) delivering PeerTECH sessions using eModules on the tablet, (e) providing the structure for the weekly sessions, (f) using role-play in teaching self-management skills, and (g) presenting an orientation to the smartphone application. Each peer had a two- to three-person caseload and worked a total of 10 hr per week, including providing direct care, text-messaging consumers, and receiving supervision. Peers were supervised by a trained peer supervisor for 1 hr each week. A verbal informed consent to participate as an interventionist in the PeerTECH study was obtained from peers during the peer training.

Research staff, with a certified peer specialist, met with potential participants in their home and provided an overview of the study and discussed the voluntary nature of study participation and confidentiality issues. The potential participant could ask any questions as needed and, if interested, complete the consent form. For those who agreed to participate, Peers met with them in participants' homes for 1 hr each week over a 12-week period.

Intervention

We adapted an evidence-based medical and psychiatric self-management intervention for older adults with SMI—Integrated Illness Management and Recovery (Bartels et al., 2014)—for in-person delivery by a certified peer specialist with the use of guided eModules and a smartphone application designed to complement in-person eModule sessions. The eModules are designed to be reviewed on a tablet side-by-side with a peer and consumer during 1-hr, weekly, in-person sessions in a community setting. Each eModule includes videos and experiential learning tasks on psychoeducation and coping skills training (see Table 1).

The smartphone intervention portion was designed to reinforce in-person sessions and to provide support to people in real-world environments. Certified peer specialists were instructed to message consumers a minimum of three times a week. There was no maximum number of text messages required. Text message exchanges were to focus on providing peer support and following up on goals and discussions during in-person sessions (e.g., “hope you are doing well on your goals—journaling and walking”). Peers were permitted to text message at any time throughout the day. All text message content was logged, and time–date was recorded. Requirements for text messaging were purposely left unstructured in an effort to examine naturalistic interaction between certified peer specialists and consumers with SMI.

Smartphones and data plans were provided free of charge or participants could choose to use their own smartphone. Participants were not provided an incentive to send text messages; however, they were provided \$20 compensation to complete baseline, 1-month, and 3-month assessments (total of \$60 over the entire study duration). These assessments and related findings were described in detail in a previously published report (Fortuna et al., 2018). Briefly, PeerTECH baseline, 1-month, and 3-month assessments were associated with statistically significant improvements in psychiatric self-management, and improvements were

Table 1
PeerTECH eModule Sessions

<i>Session 1 & 2 Identifying Your Recovery and Wellness Goals:</i> Setting recovery and health goals and strategies to achieve goals and orientation to the smartphone application;
<i>Session 3 & 4 Psychoeducation:</i> Psychoeducation on SMI and medical illness;
<i>Session 5 Stress Vulnerability and Illness:</i> Causes of mental illness and factors that influence its course;
<i>Session 6 Building Social Supports and Recovery and Wellness:</i> How to build social supports;
<i>Session 7 Medication Adherence Strategies:</i> Behavioral tailoring and motivational techniques for medication adherence;
<i>Session 8 Psychiatric and Medical Relapse Prevention:</i> Identify warning signs and develop a relapse prevention plan;
<i>Session 9 Coping with Psychiatric Symptoms and Health-related Stress and Solving Problems:</i> Establish a method managing symptoms;
<i>Session 10 Coping with Stress, Chronic Pain and Medical Symptoms:</i> Identifying stressors that exacerbate symptoms and strategies to cope with stress;
<i>Session 11 Medication Misuse:</i> Addressing medication misuse and the effects on symptoms and functioning; and
<i>Session 12 A Guide to Navigating the Mental Health and Medical Healthcare System:</i> Accessing mental health and medical health services and making informed decisions.

Note. SMI = serious mental illness.

found in self-efficacy for managing chronic health conditions, hope, quality of life, medical self-management skills, and empowerment. For this study, incoming and outgoing text messages were securely stored within the smartphone app database. Data were extracted into an Excel worksheet.

Data Analysis

Text message transcripts were analyzed for eight participants and three certified peer specialists who delivered the 12-week intervention. Our analysis of text message data was informed using the grounded-theory approach (Martin & Turner, 1986). The codebook consisted of a priori peer- and researcher-driven codes, derived from interviews and inductively derived codes from qualitative data (Martin & Turner, 1986). The first and second authors read data and incorporated new codes and operational definitions from transcript coding, a validated approach that allows for multiple perspectives (Martin & Turner, 1986). Codes were assigned to text and then grouped and checked for themes. Thematic analysis was used to summarize themes identified in the text message data (Braun & Clarke, 2006). Analyses assessed within-group consensus or disagreement. Member checking was employed to validate qualitative results and resolve any incongruent findings. As such, the lead author contacted the participating certified peer support specialists to discuss the key themes that emerged from the data. This approach helped ensure that these findings are consistent with how the certified peer specialists intended to use text messaging to provide peer support for health behavior change and self-management among consumers. Quantitative data comprised frequency of text messages by either the peer or consumer participant. Frequency data were captured directly from the app. Frequency data were integrated at the conclusion of the study.

Results

Study Sample

Our sample consisted of eight consumer participants and three certified peer specialists. Consumer participants had a mean age of 68.8 years ($SD = 4.9$; range = 62–77) and were primarily women (88%), White (100%), and married (75%). The sample included people diagnosed with major depressive disorder (63%), schizophrenia spectrum disorders (25%), and bipolar disorder (13%).

Certified peer specialists were all 55 or older; 100% were female, 66% identified as White, and 33% identified as African American. All peers completed the certified peer specialist training (Kaufman, Kuhn, & Stevens Manser, 2016) and the certified older adult peer specialist training.

Text Message Exchanges

Peers sent text messages to participants' smartphones based on consumers' preference, ranging from 8 a.m. to 10 p.m. on weekdays and weekends. Over the course of the 12-week intervention, a total of 356 text messages were sent. Peers sent 215 text messages, whereas consumers sent 141 text messages (see Figure 1). Consumer participants responded to 51% of the initial text messages from peers (as measured by a returned text message within 24 hr).




Figure 1. Aggregated frequency of consumer participant and peer text messages over the 12-week intervention period.

We identified a final set of 68 codes relating to four themes in the certified peer specialist–consumer participant text message exchanges. Themes covered different aspects of illness self-management for adults with both mental health and physical health needs. The four themes were *health behavior change*, *self-management therapeutic techniques*, *engagement in health technology*, and *peer support*.

Health Behavior Change

The first theme, health behavior change, included these subcategories: responding to help seeking, reinforcement, promoting self-care, and self-efficacy. Most consumer participants referred to a barrier to receiving support (e.g., a consumer texted “for support if feeling down and holidays can be difficult don’t feel there is much support just additional stress”). Certified peer specialists or consumer participants regularly reinforced self-management practices discussed during the in-person session (e.g., a certified peer specialist texted “how did the Managing Stress module help you and what can you use in the future to reduce or to manage your stress?”). Peer specialists frequently recommended self-care, encouraging consumer participants to take care of their mental and/or physical health through participating in meaningful activities (e.g., a certified peer specialist texted “did you think of something special to do for yourself yet?”). Certified peer specialists or consumer participants referred to their self-efficacy in their ability to self-manage (e.g., a consumer participant wrote “also on a personal note, the types of breathing in the video do not work well for me. But I got that covered”).

Self-Management Therapeutic Techniques

The second theme included text messages that were characterized as self-management therapeutic techniques. Peers used empowering text messages to encourage participants to pursue their personalized health and recovery goals that were identified during in-person sessions. This included goal setting; discussions of the interaction between mental health and physical health; and reminders to engage in daily self-management tasks such as watching mindfulness or breathing technique videos, building social supports, and advocating for themselves. Peers were able to monitor

completion of daily self-management tasks and physical activity through pedometer readings (e.g., a peer wrote “great to see you in action”). As such, peers used text messaging to follow up on consumer participants’ engagement with the smartphone application (e.g., a peer wrote “glad to see you active on the checklist”). Peers used motivational interviewing techniques via text messaging, including responding to change talk; asking open-ended questions; and providing affirmations, reflections, and summaries (e.g., a peer provided these affirmations: “as I always have said: You are doing the work. I’m just along for this part of your journey. And It is a pure joy!! To see you doing by leaps and bounds. I’m proud of you”).

Engagement in Health Technology

The third theme represented peer specialists’ impact on consumer participants’ engagement in the intervention and included these subcategories: accountability and scheduling appointments and appointment reminders. Accountability included efforts by the peer to encourage participants to follow up on in-person topics related to self-management. This included questions about what activities worked or did not work. Additionally, peers taught consumers how to engage in the text-messaging portion of the intervention (e.g., a consumer wrote “I did my first ‘SOLO’ tonight, THANKS to ‘YOU’!!! My first Text Ever!!!”). Peers and consumers frequently used text messaging for scheduling and to coordinate and/or confirm in-person sessions.

Peer Support

The fourth theme was peer support, which included these subcategories: self-determination (i.e., right to determine their own goals), shared knowledge, and shared experience (i.e., experiential learning—experiential teaching, including psychoeducational resources such as peers’ sharing their personal experiences with managing diseases; e.g., a peer wrote “somebody once told me ‘Do you want to be right or do you want to be loved’. . . . I really had to struggle with my un-forgiveness and resentment . . . I was right but what did that get me . . .”). Another example reflected reciprocal—mutual support and empowerment (a peer texted “You have the POWER to move forward. It’s up to guess who? YOU! You decided who has the power. Are you gonna give to someone else? Or are you in Control of [participant’s name]? You can’t blame anyone. YOU have always and will always had the power to keep or give it up!”). Shared values and bonding reflecting an attachment or relationship between a participant and a peer was also observed. Bonding included efforts by peer specialists to encourage participants and vice versa that demonstrated an attachment or a relationship (e.g., a peer wrote “great to spend time with you. happy birthday. I know you enjoyed yesterday, I know you will enjoy today. And you are working on enjoying the rest of your life. you are doing all this for YOU, remember that!!!”).

Finally, social support included a subset of categories that demonstrated emotional support (i.e., showing empathy, compassion, and concern for another person), informational support (i.e., providing advice, guidance, and feedback), tangible support (i.e., transportation), and companionship (e.g., sense of belonging). For example, text messages reflected peer support when consumers highlighted the value of their relationship with their peer specialist

and the importance of receiving support (e.g., a consumer participant wrote to a peer “you helped me get complete after 42 years. Thank you so much. You are such a caring person and I am thankful we are matched together. I so enjoy being with you”).

Discussion

We identified four broad text-messaging strategies that certified peer specialists use to support a home-based integrated medical and psychiatric self-management intervention. The four strategies included health behavior change, self-management therapeutic techniques, engagement in health technology, and peer support. This exploratory qualitative study offers preliminary evidence that peers are able to use text messages to support the delivery of a peer-delivered home-based medical and psychiatric self-management intervention (PeerTECH).

Peers provided reinforcement to complete self-management tasks and promoted self-care and self-efficacy through text messaging. These strategies align with accepted theories of human behavior and health behavioral change (Naslund, Aschbrenner, & Bartels, 2016). There has been little research on the use of peer-delivered text messaging to support medical and psychiatric self-management health behavior change with older adults with SMI. Older adults with SMI are a rapidly growing population with high rates of chronic illness that is associated with impaired functioning and early mortality (De Hert et al., 2011; Colton & Manderscheid, 2006; Walker, McGee, & Druss, 2015), primarily due to chronic health conditions (De Hert et al., 2011). Self-management skills such as medication management and stress reduction can help to improve physical and mental health outcomes (Whiteman, Naslund, DiNapoli, Bruce, & Bartels, 2016). Peers could potentially use text messages to support health behavior change among older adults with SMI by offering support in completing self-management tasks.

Self-management therapeutic techniques were a common topic in peer and consumer text message exchanges. Both peers and older adults with SMI were willing to share their difficulties with completing daily self-management tasks, try new coping strategies (e.g., mindfulness, guided breathing techniques), build social supports, and advocate for themselves. Self-management skills developed during in-person sessions (i.e., in a controlled environment) appear to have been independently practiced when encouraged by peers’ text messages. Although older adults have sensitivity to potential age-related changes with respect to memory and attention (Glisky, 2007), mastering new skills can be difficult. Peer text messaging around self-management therapeutic techniques appears to have acted as a reinforcement to independently practice self-management skills.

Certified peer specialists’ used text messaging to promote participant engagement in self-management activities and behavioral change. Despite the proliferation of mHealth interventions for people with SMI, mHealth interventions are often impacted by low engagement among people with SMI (Berrouiguet, Baca-García, Brandt, Walter, & Courtet, 2016; Eysenbach, 2005). The addition of a support person in the delivery of mHealth interventions can potentially increase initial and sustained engagement in health behavioral change activities (Mohr, Cuijpers, & Lehman, 2011); however, it is not known who support persons should be or what role they should have in the intervention. Initial studies using text

messaging for people with SMI have focused on clinician-delivered remote symptom monitoring (Španiel et al., 2008), support functioning (Pijnenborg et al., 2010), and delivering appointment reminders (Maiga, 2011), as well as medication and treatment adherence (Beebe, Smith, & Phillips, 2014; Montes, Medina, Gomez-Beneyto, & Maurino, 2012; Sims et al., 2012). With further development and validation, peers in combination with mHealth appears promising for promoting participant engagement in self-management activities and behavioral change.

Peers demonstrated a unique relationship with consumer participants who were grounded in principles of peer support, including mutual support, mutual respect, understanding, affection, trust, and companionship. This type of interpersonal relationship is more commonly experienced between friends or family members and not typically between a service provider and a consumer. For example, relationships documented between consumers and clinicians through text messaging demonstrated consumers were interested in discussing techniques for coping with mental health symptoms and health behavior change (Aschbrenner, Naslund, Gill, Bartels, & Ben-Zeev, 2016)—not relationship building. The impact of social ties between peers and consumers on health outcomes is well known. Examining the mechanistic effect of social ties may elucidate social networks that can potentially support the objectives of peer-delivered self-management interventions.

In interpreting these findings, it is important to consider the limitations of this study. First, it is not known whether we met saturation. Customarily, qualitative interviews are conducted until there is saturation of data (i.e., saturation means that researchers reach a point in their analysis where sampling more data will not lead to more information related to their research questions; Seale, Gobo, Gubrium, & Silverman, 2004). By drawing from grounded-theory design (Martin & Turner, 1986), saturation would generally occur with 20–30 participants total (Creswell, 2007); however, the sample size was small because the primary study was conducted to assess feasibility. It is important to note that findings cannot be generalized; however, the themes identified can be used to guide the development of peer support text-messaging services as an adjunct to evidence-based interventions. Further, we were unable to stratify our data by demographic characteristics due to the sample size. For example, one peer had a master's degree in social work, and this advanced educational background likely influenced the person's delivery of services. Second, peers met in person with consumer participants over a 12-week time frame; thus, in-person follow-up discussions from text messages are not reported. Third, the sample included a heterogeneous group of people with psychotic disorders and mood disorders that predominately was composed of people with major depressive disorder. Fourth, the consumer participants in this study were all receiving mental health services, and therefore our findings cannot generalize to individuals with serious mental illness not enrolled in care or without access to mental health services. Peer-delivered services using widely available mobile technology may hold promise for reaching and supporting individuals with serious mental illness who are reluctant or unable to seek traditional services. This is also an important area for future inquiry. Finally, the results elucidate common text message themes between certified peer specialists and consumers with serious mental illness; however, it is not known whether the peer-to-consumer text message exchanges can improve self-management and other clinical outcomes. A future

study with adequate power and a suitable comparison condition is necessary to isolate the impact of text message support on health outcomes.

Despite these limitations, this is the first study to our knowledge that has explored the content of text message exchanges between peers and consumers with serious mental illness in the context of an integrated medical and psychiatric self-management intervention. Our findings demonstrate that certified peer specialists appear to use text messages to promote health behavior change and engagement in health technology and to provide peer support and self-management therapeutic techniques. Peer-supported text messaging could potentially impact effectiveness of self-management interventions, augment the capacity of peer-supported services, and increase certified peer specialists' reach beyond traditional service delivery mechanisms.

References

- Aschbrenner, K. A., Naslund, J. A., Gill, L. E., Bartels, S. J., & Ben-Zeev, D. (2016). A qualitative study of client-clinician text exchanges in a mobile health intervention for individuals with psychotic disorders and substance use. *Journal of Dual Diagnosis, 12*, 63–71. <http://dx.doi.org/10.1080/15504263.2016.1145312>
- Bartels, S. J., Pratt, S. I., Mueser, K. T., Naslund, J. A., Wolfe, R. S., Santos, M., . . . Riera, E. G. (2014). Integrated IMR for psychiatric and general medical illness for adults aged 50 or older with serious mental illness. *Psychiatric Services, 65*, 330–337. <http://dx.doi.org/10.1176/appi.ps.201300023>
- Beebe, L., Smith, K. D., & Phillips, C. (2014). A comparison of telephone and texting interventions for persons with schizophrenia spectrum disorders. *Issues in Mental Health Nursing, 35*, 323–329. <http://dx.doi.org/10.3109/01612840.2013.863412>
- Ben-Zeev, D., Davis, K. E., Kaiser, S., Krzosos, I., & Drake, R. E. (2013). Mobile technologies among people with serious mental illness: Opportunities for future services. *Administration and Policy in Mental Health and Mental Health Services Research, 40*, 340–343. <http://dx.doi.org/10.1007/s10488-012-0424-x>
- Berrouguet, S., Baca-García, E., Brandt, S., Walter, M., & Courtet, P. (2016). Fundamentals for future mobile-health (mHealth): A systematic review of mobile phone and web-based text messaging in mental health. *Journal of Medical and Internet Research, 18*(6), e135.
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology, 3*, 77–101. <http://dx.doi.org/10.1191/1478088706qp0630a>
- Chinman, M., George, P., Dougherty, R. H., Daniels, A. S., Ghose, S. S., Swift, A., & Delphin-Rittmon, M. E. (2014). Peer support services for individuals with serious mental illnesses: Assessing the evidence. *Psychiatric Services, 65*, 429–441. <http://dx.doi.org/10.1176/appi.ps.201300244>
- Colton, C. W., & Manderscheid, R. W. (2006). Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Preventing Chronic Disease: Public Health Research, Practice, and Policy, 3*(2), A42.
- Creswell, J. (2007). *Qualitative inquiry and research design: Choosing among five approaches* (2nd ed.). Thousand Oaks, CA: Sage.
- De Hert, M., Correll, C. U., Bobes, J., Cetkovich-Bakmas, M., Cohen, D., Asai, I., . . . Leucht, S. (2011). Physical illness in patients with severe mental disorders: I. Prevalence, impact of medications and disparities in health care. *World Psychiatry, 10*, 52–77. <http://dx.doi.org/10.1002/j.2051-5545.2011.tb00014.x>
- Eysenbach, G. (2005). The law of attrition. *Journal of Medical and Internet Research, 7*(1), e11.

- Fortuna, K. L., DiMilia, P. R., Lohman, M. C., Bruce, M. L., Zubritsky, C. D., Halaby, M. R., . . . Bartels, S. J. (2018). Feasibility, acceptability, and preliminary effectiveness of a peer-delivered and technology supported self-management intervention for older adults with serious mental illness. *Psychiatric Quarterly*, *89*, 293–305. <http://dx.doi.org/10.1007/s11126-017-9534-7>
- Glisky, E. L. (2007). Changes in cognitive function in human aging. In D. R. Riddle (Ed.), *Brain aging: Models, methods, and mechanisms* (pp. 3–20). Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK3885/>
- Health Insurance Portability and Accountability Act of 1996. Pub. L. No. 104-191, § 264, 110 Stat.1936.
- International Association of Peer Support Providers. (2011). *National practice guidelines for peer supporters*. Retrieved from <https://na4ps.files.wordpress.com/2012/09/nationalguidelines1.pdf>
- Kaufman, L., Kuhn, W., & Stevens Manser, S. (2016). *Peer specialist training and certification programs: A national overview 2016*. Retrieved from <http://sites.utexas.edu/mental-health-institute/files/2017/01/Peer-Specialist-Training-and-Certification-Programs-A-National-Overview-2016-Update-1.5.17.pdf>
- Maiga, D. (2011). Intérêt de l'utilisation du téléphone mobile dans la réponse aux rendez-vous des patients atteints de psychoses aiguës fonctionnelles au service de psychiatrie de l'Hôpital national de Niamey [Interest of using a mobile phone in response to appointments for patients with acute functional psychoses at the psychiatric ward of Niamey National Hospital]. *L'Information Psychiatrique*, *87*, 127–132. <http://dx.doi.org/10.3917/inpsy.8702.0127>
- Martin, P., & Turner, B. (1986). Grounded theory and organizational research. *Journal of Applied Behavioural Science*, *22*, 141–157.
- Mohr, D. C., Cuijpers, P., & Lehman, K. (2011). Supportive accountability: A model for providing human support to enhance adherence to eHealth interventions. *Journal of Medical Internet Research*, *13*(1), e30. <http://dx.doi.org/10.2196/jmir.1602>
- Montes, J., Medina, E., Gomez-Beneyto, M., & Maurino, J. (2012). A short message service (SMS)-based strategy for enhancing adherence to anti-psychotic medication in schizophrenia. *Psychiatry Research*, *200*, 89–95. <http://dx.doi.org/10.1016/j.psychres.2012.07.034>
- Naslund, J. A., Aschbrenner, K. A., Barre, L. K., & Bartels, S. J. (2015). Feasibility of popular m-health technologies for activity tracking among individuals with serious mental illness. *Telemedicine and e-Health*, *21*, 213–216. <http://dx.doi.org/10.1089/tmj.2014.0105>
- Naslund, J. A., Aschbrenner, K. A., & Bartels, S. J. (2016). How people with serious mental illness use smartphones, mobile apps, and social media. *Psychiatric Rehabilitation Journal*, *39*, 364–367. <http://dx.doi.org/10.1037/prj0000207>
- Naslund, J. A., Marsch, L. A., McHugo, G. J., & Bartels, S. J. (2015). Emerging mHealth and eHealth interventions for serious mental illness: A review of the literature. *Journal of Mental Health*, *24*, 321–332. <http://dx.doi.org/10.3109/09638237.2015.1019054>
- Pijnenborg, G. H., Withaar, F. K., Brouwer, W. H., Timmerman, M. E., van den Bosch, R. J., & Evans, J. J. (2010). The efficacy of SMS text messages to compensate for the effects of cognitive impairments in schizophrenia. *British Journal of Clinical Psychology*, *49*, 259–274. <http://dx.doi.org/10.1348/014466509X467828>
- Seale, C., Gobo, G., Gubrium, J. F., & Silverman, D. (2004). Grounded theory. In C. Seale, G. Gobo, J. F. Gubrium & D. Silverman (Eds.), *Qualitative research practice* (pp. 80–94). <http://dx.doi.org/10.4135/9781848608191>
- Sims, H., Sanghara, H., Hayes, D., Wandiembe, S., Finch, M., Jakobsen, H., . . . Kravariti, E. (2012). Text message reminders of appointments: A pilot intervention at four community mental health clinics in London. *Psychiatric Services*, *63*, 161–168. <http://dx.doi.org/10.1176/appi.ps.201100211>
- Solomon, P. (2004). Peer support/peer provided services underlying processes, benefits, and critical ingredients. *Psychiatric Rehabilitation Journal*, *27*, 392–401. <http://dx.doi.org/10.2975/27.2004.392.401>
- Španiel, F., Vohlídka, P., Hrdlicka, J., Kožený, J., Novák, T., Motlová, L., . . . Höschl, C. (2008). ITAREPS: Information technology aided relapse prevention programme in schizophrenia. *Schizophrenia Research*, *98*, 312–317. <http://dx.doi.org/10.1016/j.schres.2007.09.005>
- Stroul, B. (1989). Community support systems for person with long-term mental illness: A conceptual framework. *Psychosocial Rehabilitation Journal*, *12*, 9–26.
- Walker, E. R., McGee, R. E., & Druss, B. G. (2015). Mortality in mental disorders and global disease burden implications: A systematic review and meta-analysis. *Journal of the American Medical Association Psychiatry*, *72*, 334–341. <http://dx.doi.org/10.1001/jamapsychiatry.2014.2502>
- Whiteman, K., Lohman, M., & Bartels, S. (2017). Peer and technology supported integrated medical and psychiatric self-management intervention for older adults with serious mental illness. *Psychiatric Services*, *68*(4), 420. <http://dx.doi.org/10.1176/appi.ps.68404>
- Whiteman, K. L., Lohman, M. C., Gill, L. E., Bruce, M. L., & Bartels, S. J. (2017). Adapting a psychosocial intervention for smartphone delivery to middle-aged and older adults with serious mental illness. *American Journal of Geriatric Psychiatry*, *25*, 819–828. <http://dx.doi.org/10.1016/j.jagp.2016.12.007>
- Whiteman, K. L., Naslund, J. A., DiNapoli, E. A., Bruce, M. L., & Bartels, S. J. (2016). Systematic review of general medical and psychiatric self-management interventions for adults with serious mental illness. *Psychiatric Services*, *67*, 1213–1225. <http://dx.doi.org/10.1176/appi.ps.201500521>
- World Medical Association. (2013). Declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA*, *310*, 2191–2194. <http://dx.doi.org/10.1001/jama.2013.281053>
- Zinman, S., Harp, H. T., & Budd, S. (1987). *Reaching across, mental health clients helping each other*. Riverside: California Network of Mental Health Clients.

Received January 23, 2018

Revision received March 23, 2018

Accepted April 3, 2018 ■