NEW YORK STATE OFFICE OF CHILDREN AND FAMILY SERVICES INDIVIDUAL ALLERGY AND ANAPHYLAXIS EMERGENCY PLAN

Instructions:

- This form is to be completed for any child with a known allergy.
- The child care program must work with the parent(s)/guardian(s) and the child's health care provider to develop
 written instructions outlining what the child is allergic to and the prevention strategies and steps that must be taken
 if the child is exposed to a known allergen or is showing symptoms of exposure.
- This plan must be reviewed upon admission, annually thereafter, and anytime there are staff or volunteer changes, and/or anytime information regarding the child's allergy or treatment changes. This document must be attached to the child's Individual Health Care Plan.
- Add additional sheets if additional documentation or instruction is necessary.

Allergen:	Type of Exposure: (i.e., air/skin contact/ingestion, etc.):	Symptoms include but are not limited to: (check all that apply)
		 ☐ Shortness of breath, wheezing, or coughing ☐ Pale or bluish skin, faintness, weak pulse, dizziness ☐ Tight or hoarse throat, trouble breathing or swallowing ☐ Significant swelling of the tongue or lips ☐ Many hives over the body, widespread redness ☐ Vomiting, diarrhea ☐ Behavioral changes and inconsolable crying ☐ Other (specify)
Marin and M		□ Shortness of breath, wheezing, or coughing □ Pale or bluish skin, faintness, weak pulse, dizziness □ Tight or hoarse throat, trouble breathing or swallowing □ Significant swelling of the tongue or lips □ Many hives over the body, widespread redness □ Vomiting, diarrhea □ Behavioral changes and inconsolable crying □ Other (specify)
NU ankoati	mever a tant of (mercillorianous) at	□ Shortness of breath, wheezing, or coughing □ Pale or bluish skin, faintness, weak pulse, dizziness □ Tight or hoarse throat, trouble breathing or swallowing □ Significant swelling of the tongue or lips □ Many hives over the body, widespread redness □ Vomiting, diarrhea □ Behavioral changes and inconsolable crying □ Other (specify)

OCFS-6029	(01/2021)							
Date of P		1					SQ.	
• C	ilect ebiuebli	rine immedia rescue squad	tely and note th	ne time when t	the first dos	e is given.	but not limited to ephrine when emerg	the sol
• La or	ay the person lie on their si	flat, raise legs	s, and keep warm	n. If breathing i	is difficult or the	ne child is vomi	ting, allow them to s	sit up

rand c

- If symptoms do not improve, or symptoms return, an additional dose of epinephrine can be given in consultation with 911/emergency medical technicians.
- Alert the child's parents/guardians and emergency contacts.
- After the needs of the child and all others in care have been met, immediately notify the office.

MEDICATION/DOSES

 Epinephrine brand or generi 	•	•	Epinephrine	brand	or	generic	;
---	---	---	-------------	-------	----	---------	---

Epinephrine dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

ADMINISTRATION AND SAFETY INFORMATION FOR EPINEPHRINE AUTO-INJECTORS

When administering an epinephrine auto-injector follow these guidelines:

- Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than the mid-outer thigh. If a staff member is accidentally injected, they should seek medical attention at the nearest
- If administering an auto-injector to a young child, hold their leg firmly in place before and during injection to
- Epinephrine can be injected through clothing if needed.
- Call 911 immediately after injection.

STORAGE OF EPINEPHRINE AUTO-INJECTORS

- All medication will be kept in its original labeled container.
- Medication must be kept in a clean area that is inaccessible to children.
- All staff must have an awareness of where the child's medication is stored.
- Note any medications, such as epinephrine auto-injectors, that may be stored in a different area.
- Explain here where medication will be stored:

MAT/EMAT CERTIFIED PROGRAMS ONLY

Only staff listed in the program's Health Care Plan as medication administrant(s) can administer the following medications. Staff must be at least 18 years old and have first aid and CPR certificates that cover all ages of children in

- Antihistamine brand or generic:
- Antihistamine dose:
- Other (e.g., inhaler-bronchodilator if wheezing):

*Note: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE

STORAGE OF INHALERS, ANTIHISTAMINES, BRONCHODILATOR

All medication will be kept in its original labeled container. Medication must be kept in a clean area that is inaccessible to All medication will be stored in a different and making inhalars, that may be stored in a different and making inhalars, that may be stored in a different and stored. Note any medications, such as asthma inhalers, that may be stored in a different area. Explain here:

ne following strategies will be taken by the child care program to minimize the risk of exposure to any allergens while above-named child is in care (add additional sheets if needed): Document plan here: **EMERGENCY CONTACTS - CALL 911** Ambulance: () Phone #: (Child's Health Care Provider:) Phone #: (Parent/Guardian: CHILD'S EMERGENCY CONTACTS Phone#: (Name/Relationship:) Phone#: (Name/Relationship:

Name/Relationship:			
Date:	1	1	
Date:	1	1	
Date:	1	1	
	Date:	Date: /	

)

Phone#: (