

<b>EAP</b>	_____
Phone #	_____
Website	_____
<b>Peer Support Member</b>	_____
Phone #	_____
<b>Peer Support Member</b>	_____
Phone #	_____
<b>Peer Support Member</b>	_____
Phone #	_____
<b>Peer Support Member</b>	_____
Phone #	_____
<b>Clinician</b>	_____
Phone #	_____
<b>Chaplain</b>	_____
Phone #	_____

<b><u>National Resources</u></b>	
National Suicide Hotline	988
WCPR (West Coast Post-trauma Retreat)	415-721-9789
https://www.frsn.org/contact.html	
info@frsn.org	
<b>Copline (24/7 Peer Listening Resource)</b>	1-800-267-5463
<b>Significant Others and Spouses Program (SOS)</b>	
https://www.frsn.org/significant-others-and-spouses.html	
<b>VALOR Officer Safety and Wellness Program</b>	
https://www.valorforblue.org/	
<b><u>KC Metro Area Resources:</u></b>	
<b><u>Heart Kansas Trauma Recover Network (TRN) - EMDR Assistance Program</u></b>	913-274-9944 (receive a call back in 24 hours)
<b>Monkey Brain Art (Using art to help with PTSD)</b>	
https://monkeybrainart.org/	
<b>Warriors Ascent</b>	
https://www.warriorsascent.org/	
<b>KC Metro Regional Coordinators</b>	
<b>Jon Koch</b> (Johnson County, Kansas Sheriff's Office) KC Metro Kansas Region	816-206-2410
<b>Matt Kellogg</b> (Liberty PD, MO) KC Metro Missouri Region	816-668-3180



## Critical Incident Stress Information

### Our Mission

Provide educational programs, to promote healthy development following stress and critical incidents experienced by first responders and their families including collaboration with culturally competent mental health professionals.

[frstmidwest.org](http://frstmidwest.org)

You have experienced a traumatic event or a critical incident (any event that causes unusually strong emotional reactions that have the potential to interfere with the ability to function normally). Even though the event may be over, you may now be experiencing or may experience later, some strong emotional or physical reactions. It is very common, in fact quite normal, for people to experience emotional aftershocks when they have passed through a horrible event.

Sometimes the emotional aftershocks (or stress reactions) appear immediately after the traumatic event. Sometimes they may appear a few hours or a few days later. And, in some cases, weeks or months may pass before the stress reactions appear.

The signs and symptoms of a stress reaction may last a few days, a few weeks, a few months, or longer, depending on the severity of the traumatic event. The understanding and the support of loved ones usually causes the stress reactions to pass more quickly. Occasionally, the traumatic event is so painful that professional assistance may be necessary. This does not imply craziness or weakness. It simply indicates that the particular event was just too powerful for the person to manage by himself/herself.

### **Here are some common signs and signals of a stress reaction**

**Physical:** chills, thirst, fatigue, nausea, fainting, twitches, vomiting, dizziness, weakness, chest pain, headaches, elevated bp, rapid heart rate, muscle tremors, shock symptoms, grinding of teeth, visual difficulties, profuse sweating, difficulty breathing. **These symptoms may need a medical evaluation from a physician.**

**Cognitive:** confusion, nightmares, uncertainty, hyper-vigilance, suspiciousness, intrusive images, blaming someone, poor problem solving, poor abstract thinking, poor attention/decisions, poor concentration/memory, disorientation of time, place or person, difficulty identifying objects or people, heightened or lowered alertness, increased or decreased awareness of surroundings

**Emotional:** fear, guilt, grief, panic, denial, anxiety, agitation, irritability, depression, intense anger, apprehension, emotional shock, emotional outbursts, feeling overwhelmed, loss of emotional control, inappropriate emotional response

**Behavioral:** withdrawal, antisocial acts, inability to rest, intensified pacing, erratic movements, change in social activity, change in speech patterns, loss or increase of appetite, hyper-alert to environment, increased alcohol consumption, change in usual communications

### **THINGS TO TRY:**

- WITHIN THE FIRST 24-48 HOURS – periods of appropriate physical exercise, alternated with relaxation will alleviate some of the physical reactions.
- Structure your time; keep busy. • You're normal and having normal reactions; don't label yourself crazy.
- Talk to people; talk is the most healing medicine
- Be aware of numbing the pain with overuse of drugs or alcohol, you don't need to complicate this with a substance abuse problem.
- Reach out; people do care
- Maintain as normal a schedule as possible.
- Spend time with others.
- Help your co-workers as much as possible by sharing feelings and checking out how they are doing.
- Give yourself permission to feel rotten and share your feelings with others.
- Keep a journal; write your way through those sleepless hours.
- Do things that feel good to you.
- Realize those around you are under stress.
- Don't make any big life changes.
- Do make as many daily decisions as possible that will give you a feeling of control over your life, i.e., if someone asks you what you want to eat, answer them even if you're not sure.
- Get plenty of rest.
- Don't try to fight reoccurring thoughts, dreams or flashbacks – they are normal and will decrease over time and become less painful.
- Eat well-balanced and regular meals (even if you don't feel like it).