

Forests with Exceptional Conservation Values Assessment Report



April 2025

Version 2

Corner Brook Pulp and Paper Ltd
Woodlands

EXECUTIVE SUMMARY

This report contains a detailed analysis of the identification of Forests with Exceptional Conservation Values (FECV) on the Defined Forest Area (DFA) of Corner Brook Pulp and Paper Limited (CBPPL). This report presents background information and decisions relating to the assessment for the presence of FECVs on the DFA of CBPPL.

Objective 4. Conservation of Biological Diversity, is one of fourteen principles in the SFI Forest Management Standard. Performance Measure 4.2 of this objective Indicators 1 and 2 requires a “Program to protect threatened and endangered species” and a “Program to locate and protect known sites of flora and fauna associated with viable occurrences of critically imperiled and imperiled species and communities also known as Forests with Exceptional Conservation Value.”

CBPPL’s High Conservation Value Forest (HCVF) Assessment Report (v.6, Jan 2017) was adapted to create this FECV report. To complete the initial High Conservation Value Forest Assessment Report, CBPPL formed a Committee of local experts, CBPPL staff, and secretariats to assess the candidates.

Subsequent versions of the FECV report have updated by CBPPL and the most recent assessment of all identified values determined the following FECVs on CBPPL Defined Forest Area, listed by the corresponding question:

Category 1

1. Conservation Value:

- Habitat for American Marten (Newfoundland population)
- Habitat for Little Brown Myotis and Northern Myotis
- Habitat for Woodland Caribou
- Habitat for Harlequin Duck and Barrow’s Goldeneye
- Mature Coniferous Habitat for Gray-cheeked Thrush, Northern Goshawk, Olive-sided Flycatcher, Red Crossbill, Winter Wren
- Open Forest Habitat for Northern Shrike
- Open Barrens and Grasslands Habitat for Peregrine Falcon and Short-eared Owl
- All-age Classes Forest Habitat for Sharp-shinned Hawk
- Shoreline Habitat for Piping Plover and Red Knot
- Wet Forest Habitat for Rusty Blackbird
- Riparian Habitat for Atlantic Salmon, Banded Killifish, American Eel, Freshwater Fishes, and Song Sparrow
- Habitat for Plant Species at Risk and Rare and Uncommon Vascular Plant Species (Appendix 6)
- Habitat for Blue Felt Lichen
- Habitat for Boreal Felt Lichen
- Red Pine
- White Pine
- Black Ash
- Yellow Birch

2. No Conservation Value

3. Exceptional Conservation Value: Gros Morne National Park, Cook’s Marsh, Middle Ridge Wildlife Reserve, Habitat for Woodland Caribou, Riparian Habitat for Atlantic Salmon

4. Habitat for American Marten (Newfoundland population)

5. Exceptional Conservation Value: Red Pine, White Pine, Yellow Birch, Black Ash,

6. **Exceptional Conservation Value:** Gros Morne National Park; IBP Sites: Serpentine Lakes, Crooked Bog, Brownmore Bog, Sandy Lake, Grand Lake Brook; West Brook Ecological Reserve, Little Grand Lake Provisional Ecological Reserve, Little Grand Lake Wildlife Reserve, Middle Ridge Wildlife Reserve, Flatwater Pond Provincial Park Reserve, Jonathan's Pond Provincial Park Reserve, Barachois Pond Provincial Park, Sir Richard Squires Memorial Provincial Park, T'Railway Provincial Park

Category 2

7. No Conservation Value

Category 3

8. No Conservation Value
9. **Exceptional Conservation Value:** Red Pine areas – Sandy Lake (Birchy Narrows), Howley; Grand Lake South Area (balsam fir/spruce/healthy white pine of all ages)
10. **Exceptional Conservation Value:** Hampden Downs (25,511ha); Cat Arm (12,713ha); Little Codroy (12,252ha)
11. **Exceptional Conservation Value:** Serpentine Areas – North Arm Hills, Northwest Gander river near bridge on Bay D'Espoir Hwy;
Limestone Areas – South shore of Serpentine Lake, Goose Arm area;
Riparian Ecosystems – Upper Humber River, Bottom Brook, Harry's River;
Red Pine areas – Sandy Lake (Birchy Narrows), Howley

Category 4

12. **Exceptional Conservation Value:** Thirty-six Protected Public Water Supply Areas servicing ~32 communities that fall on CBPPL's DFA
13. No Conservation Value
14. **Exceptional Conservation Value:** Slopes greater than 25° (46%)
15. **Exceptional Conservation Value:** Atlantic Salmon; Freshwater fishes

Category 5

16. **Exceptional Conservation Value:** Recreation; Cutting timber for wharves and boats; Income from working in the forest; Firewood to heat homes and businesses; Company roads for access to hunting, berry picking, and cabins; Sawlogs

Category 6

17. **Exceptional Conservation Value:** Indigenous Burial sites, Indigenous Sacred Sites, Indigenous Spirit Areas, Indigenous Medicine Plants
18. No Conservation Value

TABLE OF CONTENTS

EXECUTIVE SUMMARY	1
TABLE OF CONTENTS.....	3
1. OVERVIEW	4
2. PURPOSE AND METHODOLOGY	4
3. FOREST DESCRIPTION	6
4. ASSESSMENT FOR THE PRESENCE OF EXCEPTIONAL CONSERVATION VALUES	10
5. MANAGEMENT AND MONITORING STRATEGIES FOR ECVS AND FECVS	46
6. REFERENCES	63
7. GLOSSARY	68
APPENDIX 1 CANDIDATE ECVS ON CBPPL'S DFA	76
APPENDIX 2 INITIAL MEETINGS WITH STAKEHOLDERS/INTERESTED PARTIES	87
APPENDIX 3 CANDIDATE EXCEPTIONAL CONSERVATION VALUES SUBMITTED BY STAKEHOLDERS.....	89
APPENDIX 4 EXCEPTIONAL CONSERVATION VALUE FOREST ASSESSMENT COMMITTEE	92
APPENDIX 5 POSSIBLE EXCEPTIONAL CONSERVATION VALUES.....	93
APPENDIX 6 VASCULAR PLANTS OCCURRING ON THE DFA THAT HAVE BEEN RANKED S1-S3 AT THE SUBNATIONAL (S) LEVEL IN THE GENERAL STATUS OF WILD SPECIES.....	94
APPENDIX 7 SPECIES AT RISK ON THE ISLAND OF NEWFOUNDLAND.....	101
APPENDIX 8 MAPS OF EXCEPTIONAL CONSERVATION VALUES.....	107

1. Overview

CBPPL Woodlands first became involved in environmental and forest management standards in 2001 when it was certified to the International Standards Organization 14001 Environmental Management System Standard. This standard ensured CBPPL manages its activities through standard operating procedures that prevent or minimize damage to the environment. Most recently, the company certified to the Sustainable Forestry Initiative (SFI) Forest Management and Fibre Sourcing Standards.

The Sustainable Forestry Initiative (SFI) is an independent, non-profit organization that is dedicated to promoting sustainable forest management. SFI Inc. is governed by a three-chamber Board of Directors representing environmental, social and economic sectors equally. The SFI Forest Management Standard Principle 4 states that we must manage the forest in ways that protect and promote biological diversity, including animal and plant species, wildlife habitats, and ecological or natural community types. Objective 4, Conservation of Biological Diversity states that we must manage for Forests with Exceptional Conservation Value (FECV). The definition as per SFI suggests that there can be both critically imperiled (G1) and imperiled (G2) species and ecological communities in FECV.

These two terms are described below:

Critically imperiled: A plant or animal or community, often referred to as G1, that is globally extremely rare or, because of some factor(s), especially vulnerable to extinction. Typically five or fewer occurrences or populations remain, or very few individuals (<1000), or hectares (<809), or linear kilometers (<16) exist.

Imperiled: A plant or animal or community, often referred to as G2, which is globally rare or, because of some factor(s), is very vulnerable to extinction or elimination. Typically six to 20 occurrences, or few remaining individuals (1000-3000), or hectares (809 – 4047), or linear kilometers (16 – 80.5) exist.

Principle 4, Objective 4 of the Forest Management Standard addresses Forests with Exceptional Conservation Values (FECVs). All forests contain cultural, environmental and social values, and forest areas where these values are of outstanding significance or critical importance are called FECVs. CBPPL has a mandate to recognize non-timber forest values, as the forests of Newfoundland are used extensively by residents as a livelihood for logging, outfitting, and adventure tourism, and recreationally for activities such as hunting, fishing, cabins, berry picking, hiking, snowmobiling and bird watching. CBPPL has already identified a number of non-timber values on its limits through the 5-Year Planning Process, conducted by the provincial Fisheries and Land Resources (FLR), where Government and CBPPL present their proposed harvesting, road building, and silviculture plans for the next 5 years, and address issues raised by stakeholders.

This assessment report has been adapted and modified from CBPPL's High Conservation Value Forest Assessment Report (v. 6, Jan 2017). CBPPL maintains and updates the FECV Report.

2. PURPOSE AND METHODOLOGY

PURPOSE

The purpose of this report is to document some of the requirements of Principle 4 Forests with Exceptional Conservation Values of the SFI Forest Management Standard. The basis of Principle 4 is the identification of Forests with Exceptional Conservation Values (FECV) on the forest of the CBPPL. This complements CBPPL's commitment to "for the provision of ecosystem services...recreation and aesthetics" as stated in its Forest and Environmental Policy.

The mandate of this report is to describe the assessment process conducted for the Defined Forest Area (DFA) of CBPPL, the resultant FECVs identified, the management strategies necessary to ensure the

attributes of these values are maintained or even enhanced, and the monitoring required to assess the effectiveness of the management strategies.

ORIGINAL ASSESSMENT

CBPPL's High Conservation Value Forest Assessment Report (v. 6, Jan 2017) was adapted to create this FECV report. The High Conservation Value Forest Assessment Report was developed under contract, with data and information obtained through discussions with local experts, ENGO organizations, stakeholders, and other interested parties. The first step of the assessment process was to research related sources of information, to determine if there were any HCVF attributes on the DFA. This included a search of international, national, and provincial databases to determine the presence of HCVF attributes on the island of Newfoundland. Spatial data was collected when available and ArcMap software (ESRI Inc.) was used to analyze this data. All of the information was studied to determine, if possible, whether or not the attributes occurred on the DFA. If the attributes occurred or could possibly occur on the DFA, then they were listed in a table where additional information about the attributes was compiled: habitat, status, risk from forest operations, and current management (Table 4 and Appendix 1).

Once this background information was collected, stakeholders and other interested parties were invited to submit their candidate HCVFs (See Appendix 2 for list of groups and individuals). As a follow-up, the groups were sent a list of 19 questions and the rationale behind the questions, to help them understand the attributes that could create candidate HCVFs. The consultant, a CBPPL staff member, and one of the Assessment Committee local experts met with anyone that wanted to submit potential HCVF values, to discuss the value, their objective(s) for the value, ideas for management strategies, and to locate the value(s) spatially. These values (Appendix 3) were then included in the table of potential HCVFs, along with the information indicated above.

An assessment committee (Appendix 4) made up of local experts, CBPPL staff, a facilitator, and the consultant then reviewed the information in the tables. Using the definitive and guidance questions, the Committee determined which values listed were HCVFs, which were not HCVFs, and those for which more information was required before they could be assessed (Possible HCVF, Appendix 5). To address question 18 (values that did not meet HCVF thresholds but collectively constitute HCVFs), CBPPL and the Canadian Forest Service had initially considered using a GIS software package, developed to analyze multiple criteria in integrated land management plans. However after the assessment was completed, both agencies decided that since there was no apparent overlap of values that would lead to new HCVFs, and most of the submitted values were identified as HCVFs, further analysis was unnecessary.

For this FECV report, CBPPL has replaced the HCVF terminology with FECV.

MANAGEMENT STRATEGIES

The ECV Assessment Committee used information about the FECVs from Table 4 and Appendix 1, objectives for the values submitted by stakeholders/interested parties, and their own expert knowledge to develop management strategies for the identified FECVs (Section 5). It was impossible to develop management strategies for the maintenance of Possible FECVs as there is a knowledge gap in their occurrence, habitat preference, food availability, etc. Summarized in this plan, the management strategies are explained in greater detail in Forest Management Plan documents. For efficiency and simplicity, monitoring of the of FECV management strategies was incorporated into existing monitoring processes.

Information about the consultation, assessment, management, and monitoring of FECVs is presented in this report. The initial report was subjected to a credible, outside review and subsequently revised based on recommendations from the reviewer. It was posted for public review on CBPPL's website (cbppl.com), and through newspaper advertising CBPPL invited comments on all documents on the website

concerning forest management activities on their timber limits, including the reports developed as requirements for SFI certification.

ADAPTIVE MANAGEMENT

Conditions change and knowledge evolves, providing new information and opportunities. Consequently, an analysis of new information is essential to keeping the Forests with Exceptional Conservation Value Report current. CBPPL will regularly review and compile an updated list of Species at Risk to address additions, changes in status, and new information on existing species (particularly Possible FECVs). The Environmental Management Representative will ensure regular reviews of the report are conducted to analyze new information for Possible FECVs, assess new candidate FECVs, evaluate the monitoring results, and assess the need for new or revised management strategies. Following the principles of adaptive forest management and continual improvement, the necessary changes will be made and the report will be revised as required.

FOREST MANAGEMENT PLAN

The Forest Management Plan for CBPPL timber limits is contained in a number of documents, but mainly in the 5-Year Operating Plans for each Forest Management District and the Sustainable Forest Management Plan, all available on CBPPL's website. The 5-Year Operating Plans, developed through a public consultation process, contain a description of the forest, timber supply analysis, forest values, management objectives and strategies, and proposed activities for a 5-year period. They exhibit the impact and scale of the proposed operations in each of the applicable forest management districts. The Sustainable Forest Management Plan contains targets and management strategies to address local values identified through the Canadian Council of Forest Ministers Criteria and Elements for sustainable forest management.

3. FOREST DESCRIPTION

A part of Kruger Inc., CBPPL produces 257,000 tonnes of newsprint annually at its mill in Corner Brook, Newfoundland. CBPPL's timber limits (~ 1.35 million ha), under license from the Crown, extend from the Codroy Valley in the southwest, north to Cat Arm on the Northern Peninsula, and east to Gander, and fall into provincial Forest Management Districts 5, 6, 9, 10, 14, 15, & 16. The remainder is supplied by Crown operators and pulp chips from sawmills. A small area of CBPPL's license (and the responsibility for its management) has been transferred to or exchanged with the Crown. CBPPL maintains Sustainable Forest Management certification on the portion of its timber limits for which it has management responsibility, described as the Defined Forest Area (DFA). Operations on the DFA include harvesting, road construction, precommercial thinning and planting. No widespread land conversion is presently occurring nor has it occurred previous to this assessment. The DFA for CBPPL can be seen in Figure 1.




Figure 1. Defined Forest Area for CBPPL

The island of Newfoundland forms the northern extremity of the Appalachian geological province of Eastern North America. The island has been shaped by glaciation and is rugged as a result. The DFA includes parts of the island that have been scraped bare, and many valleys and low-lying areas with thick mantles of rocky glacial deposits. The DFA – like the island – is also characterized by rolling hills, mountainous areas, upland plateaus, and numerous bogs, barrens, and ponds. There are abundant fresh water sources in the forms of large rivers, lakes, and streams. Elevation on the island peaks at 814

metres above sea level in the Lewis Hills and can average 600-800 metres above sea level on other plateaus.

The geology of the DFA is typical of the island. The surface geology is characterized by large areas of coarse-textured material (glacial till deposits), washed sediments, peat deposits, and rock outcrops. Rock types tend to be predominantly limestone, predominantly sandstone, conglomerate and shale, or predominantly volcanic rock. Under well-drained conditions, the most common form of mineral soil in the province is podzols. These soils typically have an organic layer (duff), over a distinctive red or reddish-brown layer rich in iron. Most of these soils tend to be coarse textured and very acidic.

Damman (1983) divided the island of Newfoundland into nine ecoregion classifications (Figure 2). Ecoregions are defined as areas where comparable vegetation and soil can be found on the same parent material provided that these sites have experienced a similar history of disturbance. A significant portion of CBPPL's DFA (~81%) lies within two of these ecoregions, the Central Newfoundland Forest and the Western Newfoundland Forest. Approximately 8% falls in the Long Range Barrens, over 6% in the Maritime Barrens, and almost 5% in the Northern Peninsula Forest. The North Shore Forest comprises less than 0.2% of the DFA. The ecoregions can be described briefly as follows:

- Northern Peninsula Forest – Balsam fir is the dominant forest cover except at high elevations (300-400m) on the eastern side of the peninsula where black spruce appears to be a natural component of the stands.
- Western Newfoundland Forest – Balsam fir is the dominant forest cover, with white birch and yellow birch common in protected valleys below 200m elevations.
- Central Newfoundland Forest – Much of the original balsam fir-feathermoss forest types of this area have been converted by fire to black spruce, and some of the richer site types to hardwood forests dominated by white birch and aspen.
- Long Range Barrens – This ecoregion includes the mountainous areas above treeline. The only trees that occur are stunted and of poor form shaped by wind, salt or ice (and therefore non-productive), usually dominated by black spruce, balsam fir and eastern larch.
- North Shore Forest – The vegetation in this ecoregion is similar to that of the Central Newfoundland Ecoregion, except that white spruce is more abundant in the forests.
- Maritime Barrens – This ecoregion consists of usually stunted, almost pure stands of balsam fir, broken by extensive open heathland.

A detailed description of the ecoregions in the DFA can be found in *The Pre-Industrial Condition of the Forest Limits of Corner Brook Pulp and Paper Limited* and in the 5-Year Operating Plans.

The landscape of the island of Newfoundland is naturally fragmented, composed of productive forest, scrub (nonproductive forest), bog, barren, and water. The percentage of these land classes present on the DFA can be seen in Figure 2.




Figure 2. CBPPL Defined Forest Area by land class

The forests of Newfoundland belong to the boreal forest, characterized by periodic, catastrophic, stand-replacement, natural disturbances such as fire and insect outbreaks. The resulting forest is even-aged and comprised of few, primarily conifer species. Additional forest disturbances include harvesting, and blow down (often occurring when another disturbance, like insect damage, has weakened a stand). The dominant species on the DFA are balsam fir and black spruce, with smaller amounts of white spruce, eastern larch, trembling aspen, and white birch. Wildfires have established black spruce as a characteristic species across much of central Newfoundland while the forests of the west coast are predominantly balsam fir.

Gros Morne National Park, designated a UNESCO World Heritage Site for its geological features, borders on the DFA, and ten provincial parks and reserves (see Question 6 in Section 4) are adjacent to or lie within CBPPL's DFA. The DFA provides habitat for one species of global concern as well as a number of nationally and provincially ranked species at risk (Question 1 in Section 4). And because there are numerous communities within or adjacent to the DFA, 36 protected public water supply areas servicing 32 communities coincide with the DFA. CBPPL Woodlands employs ~ 175 employees from nearly 50 Newfoundland communities. This total includes workers for harvesting and road-building operations and seasonal forestry workers who carry out silviculture operations in the summer and fall. CBPPL employs another 300 people at the Mill in Corner Brook and 25 in the Deer Lake Power Company.

4. ASSESSMENT FOR THE PRESENCE OF SPECIES IN FORESTS WITH EXCEPTIONAL CONSERVATION VALUE

CATEGORY 1: FOREST AREAS CONTAINING GLOBALLY, NATIONALLY OR REGIONALLY SIGNIFICANT CONCENTRATIONS OF BIODIVERSITY VALUES.

1. Does the forest contain species at risk, or potential habitat of species at risk, as listed by international, national, or territorial/provincial authorities?

This indicator ensures the maintenance of vulnerable and/or irreplaceable elements of biodiversity, and allows for a single species or a concentration of species to meet FECV thresholds.

Sources of Assessment Information

Atlantic Canada Conservation Data Center (ACCDC)

Breeding Bird Surveys (up to 2021, NL) <https://wildlife-species.canada.ca/breeding-bird-survey-results/P003/A001/?lang=e>

The State of Canada's Birds 2024 <https://naturecounts.ca/nc/socb-epoc/main.jsp>

Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) <http://checklist.cites.org>

International Union for Conservation of Nature (IUCN) Red List of Threatened Species <http://discover.iucnredlist.org/>

NatureServe Explorer <https://explorer.natureserve.org/>

Committee on the Status of Endangered Wildlife in Canada (COSEWIC)

Species at Risk Act (SARA) Public Registry <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Species Status Advisory Committee (SSAC)

Interviews with local experts, NL Department of Fisheries and Land Resources ¹

Targeting conservation towards species in greatest need requires the determination of which are rare or declining. A number of factors are used to assess the conservation status of plant, animal and fungal species, leading to the designation of a conservation status rank.

In 2010 CBPPL requested that the ACCDC conduct an analysis to map the listed plants, rare plants, and rare birds in their database that were recorded as occurring on the DFA at that time. In 2019 CBPPL did further updates and sent an additional list of possible species at risk for analysis to the ACCDC. In 2024, the ACCDC provided observation data for all species of conservation concern within the extent of DFA with CBPPL.

Species of Global Significance

ACCDC Database (G1 and G2 ranked species): **Laurentian Dandelion** (*Taraxacum laurentianum*) (G1) and **Northern Long Eared Bat** (*Myotis septentrionalis*) (G2) are listed in the ACCDC database as occurring on the DFA. A ranking of G1 indicates a species that is critically imperiled across its entire range; G2 indicates the species is imperiled.

¹ Shelley Garland, Manager, Wildlife Research, Endangered Species and Biodiversity
Shelley Moores, Sr. Mgr. Endangered Species and Biodiversity
Susan Squires, Ecosystem Management Ecologist, Endangered Species

CITES: Appendix I, II, and III of CITES lists species that are protected from over-exploitation through international trade. **Northern Goshawk** (*Accipiter gentilis*), **Rough-Legged Hawk** (*Buteo lagopus*), **Northern Saw-Whet Owl** (*Aegolius acadicus*), **Boreal Owl** (*Aegolius funereus*), **American Kestrel** (*Falco sparverius*), **Red Crossbill** (*Loxia curvirostra*), **Snowy Owl** (*Nyctea scandiaca*), **Northern Hawk Owl** (*Surnia ulula*), **Lynx** (*Lynx Canadensis*), **Short-Eared Owl** (*Asio flammeus*), **Showy Ladyslipper** (*Cypripedium reginae*), **Summer Coralroot** (*Corallorhiza maculata*), **Yellow Ladyslipper** (*Cypripedium parviflorum*), **Checkered Rattlesnake Plantain** (*Goodyera tessellata*), **White Adder's Mouth** (*Malaxis monophyllos*), **Auricled Twayblade** (*Neottia auriculata*), **Northern Twayblade** (*Neottia borealis*), **Broad Leaved Twayblade** (*Neottia convallarioides*), **Greater Purple Fringed Bog Orchid** (*Platanthera grandiflora*), **Hooker's Bog Orchid** (*Platanthera hookeri*), **Ragged Fringed Orchid** (*Platanthera lacera*), and **Round Leaved Orchid** (*Platanthera orbiculata*) are the species listed by CITES for the island of Newfoundland that are also identified in the ACCDC database as occurring on the DFA. **Cooper's Hawk** (*Accipiter cooperii*), **Red-Tailed Hawk** (*Buteo jamaicensis*), **Sharp-Shinned Hawk** (*Accipiter striatus*), **Merlin** (*Falco columbarius*), **Bald Eagle** (*Haliaeetus leucocephalus*), **Black Bear** (*Ursus americanus*), and **Great Horned Owl** (*Bubo virginianus*) are also listed by CITES, and while their habitat range does include the CBPPL landbase, these species were not identified in the ACCDC database.

IUCN: **Boreal Felt Lichen** (*Erioderma pedicellatum*), **Suckley's Cuckoo Bumble Bee** (*Bombus suckleyi*), **Appalachian Matchsticks** (*Pilophorus fibula*), and **Smooth Horsehair Lichen** (*Byoria salazinicola*) are included on the Red List and are found in Newfoundland. None of the species is listed in the ACCDC database as occurring on the DFA.

NatureServe Explorer: There were 21 species ranked as G1 or G2 under NatureServe that are found in Newfoundland (Table 1).

Table 1. Natureserve species ranked as G1 or G2 in Newfoundland.

Northern Long Eared Bat (<i>Myotis septentrionalis</i>)	Small Four-toothed Moss (<i>Tetradontium repandum</i>)
Newfoundland Vallonia (<i>Vallonia terraenovae</i>)	Montane Trematodon Moss (<i>Trematodon montanus</i>)
Suckley's Cuckoo Bumble Bee (<i>Bombus suckleyi</i>)	Blunted Earwort (<i>Diplophyllum obtusatum</i>)
Baltic Saltbrush (<i>Atriplex nudicaulis</i>)	Boreal Felt Lichen (<i>Erioderma pedicellatum</i>)
Fernald's Braya (<i>Braya fernaldii</i>)	Witch's Hair Lichen (<i>Alectoria fallacina</i>)
Long's Braya (<i>Braya longii</i>)	Red Firedot Lichen (<i>Polycauliona leuteominia</i>)
Dense Draba/Whitlow-grass (<i>Draba pycnosperma</i>)	A boulder lichen (<i>Porpidia grisea</i>)
Barrens Willow (<i>Salix jejuna</i>)	A lichen (<i>Fuscidea lowensis</i>)
Porsild's Bryum (<i>Haplodontium macrocarpum</i>)	A lichen (<i>Thelocarpon epibolum</i>)
Shortleaf Bristle Moss (<i>Seligeria brevifolia</i>)	A fungus (<i>Phaeocollybia gregaria</i>)
Splachnoid Trumpet Moss (<i>Tayloria splachnoides</i>)	

Of these species, only the **Northern Long Eared Bat** is identified on the ACCDC database as occurring on the DFA. Northern Long Eared Bat is ranked globally as G2 by NatureServe and S1S3 by ACCDC.

Species of National (Canadian) Significance

In Canada, the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) assesses the national status of species that are considered to be at risk. Canada's Species at Risk Act (SARA) provides for the legal protection of wildlife species and the conservation of their biological diversity. Table 2 lists the endangered or threatened species, or species of special concern as ranked by COSEWIC that occur on the DFA (historically or currently) as identified in the ACCDC database, Breeding Bird Surveys and The State of Canada's Birds 2024.

Table 2. Species on the DFA listed in Schedule 1 of the Species at Risk Act and/or assessed as Endangered, Threatened, or Special Concern by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC).

Species	COSEWIC Designation	SARA Species at Risk
American Marten (<i>Martes americana atrata</i>)	Special Concern	Schedule 1
Woodland Caribou - NL pop'n (<i>Rangifer tarandus</i>)	Special Concern	Schedule 1
Little Brown Myotis (<i>Myotis lucifugus</i>)	Endangered	Schedule 1
Northern Myotis (<i>Myotis septentrionalis</i>)	Endangered	Schedule 1
Bank Swallow (<i>Riparia riparia</i>)	Threatened	Schedule 1
Barn Swallow (<i>Hirundo rustica</i>)	Special Concern	Schedule 1
Barrow's Goldeneye (<i>Bucephala islandica</i>)	Special Concern	Schedule 1
Harlequin Duck (<i>Histrionicus histrionicus</i>)	Special Concern	Schedule 1
Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Special Concern	Schedule 1
Piping Plover (<i>Charadrius melodus melodus</i>)	Endangered	Schedule 1
Red Crossbill (<i>Loxia curvirostra perena</i>)	Threatened	Schedule 1
Red Knot (<i>Calidris canutus rufa</i>)	Endangered	Schedule 1
Rusty Blackbird (<i>Euphagus carolinus</i>)	Special Concern	Schedule 1
Short-eared Owl (<i>Asio flammeus</i>)	Threatened	Schedule 1
Banded Killifish (<i>Fundulus diaphanous</i>)	Special Concern	Schedule 1
Blue Felt Lichen (<i>Degelia plumbea</i>)	Special Concern	Schedule 1
Boreal Felt Lichen (<i>Erioderma pedicellatum</i>)	Special Concern	Schedule 1
Wrinkled Shingle Lichen (<i>Pannaria lurida</i>)	Threatened	Schedule 1
Bobolink (<i>Dolichonyx oryzivorus</i>)	Special Concern	Schedule 1
Common Nighthawk (<i>Chordeiles minor</i>)	Special Concern	Schedule 1
Evening Grosbeak (<i>Coccothraustes vespertinus</i>)	Special Concern	Schedule 1
Mountain Holly Fern (<i>Polystichum scopulinum</i>)	Threatened	Schedule 1
Transverse Lady Beetle (<i>Coccinella transversoguttata</i>)	Special Concern	Schedule 1
Yellow-banded Bumble Bee (<i>Bombus terricola</i>)	Special Concern	Schedule 1

Species of Importance on the Island of Newfoundland

Newfoundland and Labrador's [Endangered Species Act](#) (ESA) provides special protection for plant and animal species considered to be endangered, threatened, or vulnerable (special concern) in the province. Designation under the Act follows recommendations from the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) and/or the Species Status Advisory Committee (SSAC) on the appropriate assessment of a species. SSAC is an independent, provincial committee of government and non-government scientists who determine the provincial status of species, subspecies, and populations.

Currently there are 51 species, subspecies, and populations listed under the Act. Those that ACCDC lists as occurring on the DFA are listed in Table 3. Some species have not been recorded on the DFA, but because their habitat is present on the DFA, they could possibly occur there. These species have been designated Possible ECV (Appendix 5).

Table 3. Species at Risk on the DFA listed in Newfoundland and Labrador's Endangered Species Act (NLESA).

Species	NLESA
American Marten (<i>Martes americana atrata</i>)	Vulnerable
Barrow's Goldeneye (<i>Bucephala islandica</i>)	Vulnerable
Bobolink (<i>Dolichonyx oryzivorus</i>)	Vulnerable
Common Nighthawk (<i>Chordeiles minor</i>)	Vulnerable
Harlequin Duck	Vulnerable
Newfoundland Gray-cheeked Thrush (<i>Catharus minimus minimus</i>)	Threatened
Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Vulnerable
Peregrine Falcon (<i>Falco peregrinus anatum</i>)	Vulnerable
Piping Plover (<i>Charadrius melodus melodus</i>)	Endangered
Red Crossbill (<i>Loxia curvirostra percna</i>)	Threatened
Red Knot (<i>Calidris canutus rufa</i>)	Endangered
Rusty Blackbird (<i>Euphagus carolinus</i>)	Vulnerable
Short-eared Owl (<i>Asio flammeus</i>)	Threatened
American Eel (<i>Anguilla rostrata</i>)	Vulnerable
Banded Killifish (<i>Fundulus diaphanous</i>)	Vulnerable
Black Ash (<i>Fraxinus nigra</i>)	Threatened
Crowded Wormseed Mustard (<i>Erysimum coarctatum</i>)	Endangered
Cutleaf Fleabane (<i>Erigeron compositus</i>)	Endangered
Feathery False Solomon's Seal (<i>Maianthemum racemosum</i>)	Endangered
Lindley's Aster (<i>Symphyotrichum ciliolatum</i>)	Endangered
Northern Bog Aster (<i>Symphyotrichum boreale</i>)	Endangered
Northern Twayblade (<i>Neottia borealis</i>)	Endangered
Rattlesnackroot (<i>Nabalus racemosus</i>)	Endangered
Red Pine (<i>Pinus resinosa</i>)	Endangered
Rock Dwelling Sedge (<i>Carex petricosa</i> var. <i>misandroides</i>)	Endangered
Sharpleaf Aster (<i>Oclemena acuminata</i>)	Threatened
Tradescant's Aster (<i>Symphyotrichum tradescantii</i>)	Threatened
Vreeland's Striped Coralroot (<i>Corallorhiza striata</i> var. <i>vreelandii</i>)	Endangered
Boreal Felt Lichen (<i>Erioderma pedicellatum</i>)	Vulnerable ¹
Blue Felt Lichen (<i>Degeli/Pectinia plumbea</i>)	Vulnerable

In addition to the species identified in Table 3, the ACCDC database identified a number of species occurring on the DFA that have been ranked at the subnational (S) level in the general status of wild species. Species ranked as S1 are critically imperiled, S2 imperiled, and S3 vulnerable. This S ranking serves as a guide for the Species Status Advisory Committee in the selection of species to be assessed. Upon recommendation from Environmental Nongovernmental Organizations in the province, all S1-S3 species occurring on the DFA have been considered for ECVs. The S1-S3 bird species are included in Table 4; the S1-S3 vascular plants are listed in Appendix 6.

The Atlantic Canada Conservation Data Center also lists additional tree species as rare or sensitive, and these were also been included for assessment:

- White Pine, *Betula alleghaniensis*
- Yellow Birch, *Betula alleghaniensis*

Species that are currently being assessed by the SSAC but not yet listed in the Endangered Species Act will be assessed as they are listed, at each review of this report.

All of the species at risk identified by international, national, or provincial authorities are listed in a table (Appendix 7) which includes information on their conservation status (risk designation), the potential for the presence of their habitat on the DFA, and whether they have actually been recorded on the DFA. This table helps determine which species at risk should be assessed as having an ECV.

For select species at risk occurring on the DFA as listed in Appendix 7, information on habitat, extent of occurrence on the DFA, and threats from forest operations are compiled (Table 4). The assessment decisions – ECV, Not ECV, or Possible ECV – are also included in this table.

There are no ecological or taxonomic groups of rare species that would together constitute an ECV. However, the DFA does contain critical habitat for American Marten. To date, this is the only species at risk with identified critical habitat.

Table 4. Assessment of Species at Risk occurring on CBPPL's DFA as having Exceptional Conservation Values.

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
Mammals								
American Marten – Newfoundland population <i>Martes americana atrata</i>	Yes	SARA –Special Concern; ESA – Vulnerable; Population possibly rebounding	Martens select for, or use in proportion to availability, a broad range of habitat types, including recent cuts ≤5 yr old, regenerating forest <6.5 m, precommercially thinned stands, and mature and overmature forest	Three of five subpopulations occur within or adjacent to the DFA: Georges Lake/Pinchgut Lake, Little Grand Lake, and Main River	Extensive harvesting would temporarily remove habitat; habitat modeling of harvesting plans shows a modest increase in potential marten habitat across the DFA with harvest of 100% AAC	Critical habitat identified; fully protected in ecological reserves and parks; snaring techniques for hare introduced to reduce incidental captures of marten ⁶		Habitat for American Marten ECV
Little Brown Myotis <i>Myotis lucifugus</i>	Yes	SARA – Endangered	Roost in trees in day; maternity colonies gather in barns, attics, tree cavities. ⁷ Need mature forests across the landscape.	Widespread across the DFA	Extensive harvesting would temporarily remove habitat	Bat Working Groups in Canada and the US are delineating the abundance of these species	Rapidly spreading white-nose syndrome (WNS) is ultimately expected to cause regional and likely range wide extinction in a very short ecological time frame.	Habitat for Little Brown Myotis ECV

² Confirmed by NL Wildlife Division Personnel: Shelley Moores, Emily Herdman, Susan Squires, Claudia Hanel

³ [NaturesServe Explorer](#)

⁴ Confirmed by NL Wildlife Division Personnel: Shelley Moores, Emily Herdman, Susan Squires, Claudia Hanel

⁵ Atlantic Canada Conservation Data Center (ACCDC)

⁶ [Recovery plan for the threatened Newfoundland population of American marten \(*Martes americana atrata*\)](#), 2010

⁷ [COSEWIC assessment and status report on the Little Brown Myotis *Myotis lucifugus*, Northern Myotis *Myotis septentrionalis* and Tri-colored Bat *Perimyotis subflavus* in Canada](#), 2013

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
Northern Myotis <i>Myotis septentrionalis</i>	Yes	SARA – Endangered	Found in dense forest stands and chooses maternity roosts beneath exfoliating bark and in tree cavities ⁶ . Need mature forests across the landscape	Less widespread across the DFA than Little brown Myotis	Extensive harvesting would temporarily remove habitat	Bat Working Groups in Canada and the US are delineating the abundance of these species	Rapidly spreading white-nose syndrome (WNS) is ultimately expected to cause regional and likely range wide extinction in a very short ecological time frame.	Habitat for Northern Myotis ECV
Woodland Caribou <i>Rangifer tarandus</i>	Yes	SARA – Special Concern	Mostly barren land during the summer months, moving to areas of mixed forests during the colder months	Caribou occur all across CBPPL limits, except in FMD 5; caribou occur in FMD 5 but not on CBPPL limits	Harvesting could impact calving and wintering areas	There are woodland caribou core and calving areas identified for both conservation and restoration herds with no go, restricted road building, and seasonal restricted areas. 5 Year Plans must be approved by gov't prior to harvest including in or adjacent to caribou habitat.		Habitat for Woodland Caribou ECV
Birds								

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
Bank Swallow <i>Riparia riparia</i>	No	SARA - Threatened	Aerial other than nests. Open and partly open situations, frequently near water. Nests in steep sand, dirt, or gravel banks.	Occurs on DFA	Roadbuilding could remove nesting habitat	Birds and their nests are protected under the Migratory Birds Convention Act.		Shoreline Habitat ECV
Barrow's Goldeneye <i>Bucephala islandica</i>	No	SARA –Special Concern; ESA – Vulnerable Population possibly still in decline	Occur only winter on the island of Newfoundland, in open water in bays.	One sighting at mouth of Humber River; possible on bays; Stephenville Crossing	No impact in wintering areas; required buffers on waterways protect bays from effects of harvesting.	Birds and their nests are protected under the Migratory Birds Convention Act. Need to gather knowledge of the full extent of the breeding (and other) ranges ⁸	No perceived threat from harvesting	Habitat for Barrow's Goldeneye ECV
Bobolink <i>Dolichonyx oryzivorus</i>	No	SARA –Special Concern; ESA – Vulnerable	Herbaceous wetlands	Occurs on DFA	No perceived threat from harvesting	Birds and their nests are protected under the Migratory Birds Convention Act; 50m buffer on significant wetlands		Wet Forest Habitat ECV
Common Nighthawk <i>Chordeiles minor</i>	No	SARA –Special Concern; ESA – Vulnerable	Mountains and plains in open and semi-open areas; open coniferous forests, grasslands, and fields. Nesting	Occurs on DFA	Reforestation efforts can negatively impact nesting habitat	Birds and their nests are protected under the Migratory Birds Convention Act.		Open Country Habitat ECV

⁸ [A management plan for Barrow's Goldeneye \(*Bucephala islandica*; Eastern population\) in Newfoundland and Labrador](#), 2006

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
			occurs on the ground on a bare site in an open area.					
Evening Grosbeak <i>Coccothraustes vespertinus</i>	No	SARA –Special Concern	Coniferous (primarily spruce and fir) and mixed coniferous-deciduous woodland, and second growth	Occurs on DFA	Insect control (budworm) impacts food availability	Nests are protected under NL Wild Life Regulations.		All Age-classes Forest Habitat ECV
Harlequin Duck <i>Histrionicus histrionicus</i>	No	SARA –Special Concern; ESA – Vulnerable Population in eastern North America well below historic levels; increases at four strategic local wintering areas	Breed along fast-flowing turbulent rivers; Wintering habitat consists of rocky coastline, subtidal ledges, and exposed headlands.	Breed along the Great Northern Peninsula – FMD 15 & 16 Sightings at mouth of Humber, Upper Humber.	Logging activities are known to increase stream siltation that may affect food availability.	Birds and their nests are protected under the Migratory Birds Convention Act. Monitor population levels and understand threats; 30-m buffers in waterfowl breeding, moulting, and staging areas ⁹	No nests recorded on the island of Newfoundland	Habitat for Harlequin Duck ECV
Newfoundland Gray-Cheeked Thrush <i>Catharus minimus minimus</i>	Yes	ESA - Threatened	On the breeding grounds, prefer dense low coniferous woods, including young regenerating forest, open-canopy old-	No sightings listed in the ACCDC, however reported sightings or nests in FMDs 5, 9, 14, 15, & 16	Though tolerant of local clearcutting, loss of large proportions of forested habitat on a larger scale is likely to have	Birds and their nests are protected under the Migratory Birds Convention Act. Increasing knowledge about the	At this time, buffers currently required in the <i>Environmental Protection Guidelines for Forestry Operations in</i>	Mature Coniferous Forest Habitat ECV

⁹ [Management Plan for the Harlequin Duck \(*Histrionicus histrionicus*\) Eastern Population, in Atlantic Canada and Québec, 2007](#)

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
			growth forests having a dense growth of shrubs and small conifers in the understory, and dense, stunted spruce and fir on windblown sites and near the tree line.		a negative impact.	species in Newfoundland, including population size and trends, demographics, distribution, habitat requirements and threats. ¹⁰	<i>Newfoundland and Labrador</i> [EPG] provide some protection for most if not all of the species listed	
Northern Goshawk <i>Accipiter gentilis</i>	Yes	NatureServe - S3B	Prefer mature forest	Generally distributed across the DFA; sightings listed in the ACCDC	Loss of large proportions of forested habitat on a large scale is likely to have a negative impact.	No management plans exist but they are covered by the raptor protection in the Environmental Protection Guidelines.		Mature Coniferous Forest Habitat ECV
Northern Hawk-owl <i>Surnia ulula</i>	No	NatureServe - S3	Open coniferous or mixed forest, forest edge and clearings, old deciduous forest burns, dense shrubby areas (especially tamarack), swamps, scrubby second-growth woodland and muskeg	Occurs on DFA	Though tolerant of local clearcutting, loss of large proportions of forested habitat on a larger scale is likely to have a negative impact.	No management plans exist but they are covered by the raptor protection in the Environmental Protection Guidelines.		Open Forest Habitat ECV
Northern Saw-whet Owl <i>Aegolius acadicus</i>	No	NatureServe - S3B	Dense coniferous or mixed forest, alder thickets, swamps, and tamarack bogs.	Occurs on DFA	Large clear-cuts and even-aged management remove nest trees	No management plans exist but they are covered by the raptor protection in the Environmental		Collapsed Mature Coniferous Forest Habitat

¹⁰ [The Status of Grey-cheeked Thrush \(Catharus minimus\) in Newfoundland and Labrador](#), 2010

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
						Protection Guidelines.		ECV
Northern Shrike <i>Lanius excubitor</i>	No	NatureServe - S3N	Prefer mature forest	Generally distributed across the DFA; sightings listed in the ACCDC	Loss of large proportions of forested habitat on a large scale is likely to have a negative impact.	Birds and their nests are protected under the Migratory Birds Convention Act. No management plans exist but they are covered by the raptor protection in the Environmental Protection Guidelines.		Open Forest Habitat ECV
Olive-sided Flycatcher <i>Contopus Cooperi</i>	Yes	SARA –Special Concern; ESA – Vulnerable	Mature coniferous, open areas: blowdown, insect-killed, clearcuts	No sightings listed in the ACCDC, but possible sightings in Main River	Possibly low nesting success in clearcuts versus other open areas	Birds and their nests are protected under the Migratory Birds Convention Act. A provincial management plan is in progress; need to gather more information		Collapsed Mature Coniferous Forest Habitat ECV
Peregrine Falcon ssp. Anatum <i>Falco peregrinus anatum</i>	No	SARA –Not At Risk; ESA – Vulnerable Population increasing nationally; decreases in some inland Labrador sites	Do not breed on the island of Newfoundland, but do migrate through; habitat choice during migration is quite broad; follow leading lines such as coasts, lake edges, and mountain ranges;	Near Gros Morne National Park	No perceived threat from harvesting; unlikely threat from forest access road construction	Work focuses only on breeding areas in Labrador; conduct surveys every 5 years		Open Country Habitat ECV

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
			hunt in open areas					
Piping Plover <i>Charadrius melodus melodus</i>	No	SARA – Endangered; ESA – Endangered Population increasing	Nests on sandy beaches on the southwest and west coasts of Newfoundland.	Stephenville Crossing	Nil	Birds and their nests are protected under the Migratory Birds Convention Act and the Species at Risk Act. Habitat protection is through enforcement and landowner and public education ¹¹	No perceived threat from harvesting	Shoreline Habitat ECV No Management Required
Red Crossbill <i>Loxia curvirostra percna</i>	Yes	SARA – Threatened; ESA – Threatened Population in decline	Thought to be mature pine, spruce and fir stands capable of producing abundant cones.	Highest numbers of reports occurring in older forests in western Newfoundland; Sightings in Tompkins/Doyles	Debate as to the significance of harvesting impact on food source (cones) and nesting habitat	Birds and their nests are protected under the Migratory Birds Convention Act. Emphasis on reducing knowledge gap about the species ¹²		Mature Coniferous Forest Habitat ECV
Red Knot <i>Calidris canutus rufa</i>	No	SARA – Endangered; ESA – Endangered Population in decline	Open, sandy estuaries have been identified as prime habitat, with rotting kelp deposits ranked as the second best habitat type.	Around almost the entire coast, however, the majority of sightings have been on the west coast	Nil	Birds and their nests are protected under the Migratory Birds Convention Act and the Species at Risk Act.	No perceived threat from harvesting	Shoreline Habitat ECV No Management Required

¹¹ [National Recovery Plan for the Piping Plover \(*Charadrius melodus*\), 2002](#)

¹² [Recovery Strategy for the Red Crossbill, *percna* subspecies \(*Loxia curvirostra percna*\), in Canada](#), 2006

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
				(Stephenville Crossing)		Eastern Habitat Joint Venture has a municipal stewardship agreement with Stephenville Crossing and St. Paul's ¹³		
Rough-legged Hawk <i>Buteo lagopus</i>	No	NatureServe - S2S3	Tundra, woodland – conifer, grassland/herbaceous, cliff	Occurs on DFA	Little to no impact	No management plans exist but they are covered by the raptor protection in the Environmental Protection Guidelines.		Open Country Habitat ECV
Rusty Blackbird <i>Euphagus carolinus</i>	Yes	SARA – Endangered; ESA – Vulnerable Population in decline	Boreal forest, favouring the shores of wetlands and slow-moving streams, peat bogs, marshes, swamps, beaver ponds	Sighting in Corner Brook area, Codroy Valley, Stephenville Crossing, Gander, White Bay	No impact; very rarely in interior of forest.	No specific management for the Rusty Blackbird in Newfoundland	No perceived threat from harvesting	Wet Forest Habitat ECV
Sharp-shinned Hawk <i>Accipiter striatus</i>	No	NatureServe - S3B	Open marshlands, deep grass fields, second-growth forest	Likely to occur on southwest coast from Stephenville to Codroy	Little to no impact		Sharp-shinned Hawks are very adaptable	All Age-classes Forest Habitat ECV
Short-eared Owl <i>Asio flammeus</i>	Yes	SARA – Threatened; ESA – Threatened	Relatively open habitat of marshland and deep grass fields. Not in boreal forest	Sighting in Stephenville Crossing area	No impact	Monitoring Distribution and Population Trends, Habitat Assessment ¹⁴	No perceived threat from harvesting; unlikely threat from forest	Open Country Habitat ECV

¹³ [Recovery Plan for Red Knot, rufa subspecies \(*Calidris canutus rufa*\), in Newfoundland and Labrador](#), 2009

¹⁴ [A management plan for the Short-eared owl \(*Asio flammeus flammeus*\) in Newfoundland and Labrador](#), 2005

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
		Population trend unknown; thought stable at historical levels					access road construction	
Song Sparrow <i>Melospiza melodia</i>	No	S4B	Habitat varies; marsh and shrub areas, forest openings and edge, riparian areas.	Widespread	Minimal for this species, possibly destroying some nesting and foraging habitat but creating new opportunities.	Birds and their nests are protected under the Migratory Birds Convention Act		Riparian Habitat ECV
Winter Wren <i>Troglodytes troglodytes</i>	Yes	S3B	Mature forests with complex forest structure, such as snags, downed trees, riparian areas	Widespread	Negatively affected by harvesting	Birds and their nests are protected under the Migratory Birds Convention Act		Mature Coniferous Forest Habitat ECV
Fishes								
Banded Killifish <i>Fundulus diaphanus</i>	No	SARA –Special Concern; ESA – Vulnerable Population stable; found in more areas than thought	Observed in the shallows and quiet areas of clear lakes and ponds with a muddy or sandy substrate, high detrital content and abundant submerged aquatic vegetation	Majority of populations are clustered on the south and southwest coasts, with an outlier population in the Indian Bay watershed; also near Stephenville Crossing and St. George's	Forest harvesting and road building may negatively impact banded killifish habitat if these activities result in increased sedimentation of adjacent waterbodies ¹⁵ .	Surveys and population monitoring; development of site specific best management practices to ensure habitat protection ^{16,17} In the Indian Bay watershed: 100m buffers on most ponds with 150m and 200m	No perceived threat from harvesting	Riparian Habitat ECV

¹⁵ [Forestry Guidelines for the Protection of Fish Habitat in Newfoundland and Labrador.](#)

¹⁶ [\[Draft\] Management Plan for the Banded Killifish \(Fundulus diaphanus\) in Newfoundland](#), 2006

¹⁷ [Management Plan for the Banded Killifish \(Fundulus diaphanous\), Newfoundland Population in Canada](#)

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
						on specific ponds. ¹⁸		
American Eel <i>Anguilla rostrata</i>	No	COSEWIC – Threatened; ESA – Vulnerable NL population unknown; probably due to changes in marine systems	Found in most coastal areas and adjacent accessible rivers in Newfoundland	Yes	No impact	Current management practices under DFO are related to the commercial eel fishery ¹⁹	No perceived threat from harvesting	Riparian Habitat ECV
Plants²⁰								
Laurentian Dandelion <i>Taraxacum laurentianum</i>	No	NatureServe - S1 Population trend unknown	Calcareous or basic substrates, such as limestone barrens or dolomite; “slatey” talus	One location appears to be plotted wrong in GIS to fall into DFA, but there is a possibility of existence in DFA.	Unlikely unless disturbed during road building.	Recovery plan not completed to date	No perceived threat from harvesting	Habitat for Laurentian Dandelion ECV
Rock Dwelling Sedge <i>Carex petricosa</i> var. <i>misandroides</i>	No	ESA- Endangered Population trend unknown	Limestone cliffs, barrens, tablelands, talus, and screes	William Wheeler Point, Woman’s Cove (Goose Arm), Shag Cliff (Bonne Bay)	Unlikely unless disturbed during road building.	Recovery plan not completed to date, Status Report only ²¹	No perceived threat from harvesting	Habitat for Rock Dwelling Sedge ECV
Serpentine Stitchwort <i>Minuartia marcescens</i>	No	NatureServe – G3; ACCDC – G3 Population trend unknown	Serpentine barrens	Northern and central Newfoundland	Unlikely unless disturbed during road building.	Recovery plan not completed to date	No perceived threat from harvesting	Habitat for Serpentine Stitchwort ECV

¹⁸ Memorandum of Understanding and Access Agreement between Corner Brook Pulp and Paper and Indian Bay Ecosystem Corporation.

¹⁹ [Management Plan for the American Eel \(*Anguilla rostrata*\) in Newfoundland and Labrador](#), 2010

²⁰ Habitat for plants taken from Meades et al, 2000

²¹ [The Status of Rock Dwelling Sedge \(*Carex petricosa* var. *misandroides*\) in Newfoundland and Labrador](#), 2008

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
Tradescant's Aster <i>Symphotrichum tradescantii</i>	No	ESA- Threatened Population trend unknown	Rocky backwater, sedge/rush flat, sandy/muddy steady	Southwest Brook, Bottom Brook	Unlikely unless disturbed during road building.	None to date, Status Report only ²²	No perceived threat from harvesting	Habitat for Tradescant's Aster ECV
Vreeland's Striped Coralroot		ESA - Endangered	Inhabits semi-open second-growth forest of Balsam Fir, Black Spruce, Balsam Poplar, Eastern Larch, Mountain White Birch, and Speckled Alder ²³	Occurs in Lomond Campground, near boundary of DFA.	Cutting of trees where the plants occur would likely lead to the demise of the plants	None to date, Status Report only		Habitat for Vreeland's Striped Coralroot ECV
Blue Felt Lichen <i>Degelia plumbea</i>		SARA – Special Concern; ESA - Vulnerable	Found in moist habitats or close to stream and lake margins. In Newfoundland it grows mainly on yellow birch but very occasionally occurs also on white spruce ²⁴ .	May occur on the DFA as occurs in Crabbe's River area.	The harvest of mature hardwood for firewood and browsing by the large populations of moose will limit the future availability of old yellow birch, the main host for this lichen.	Riparian buffers related to commercial forestry developments are required but are modest (~30-50m) and unlikely to conserve adequately the macro- and micro-habitat needs.		Habitat for Blue Felt Lichen ECV
Boreal Felt Lichen <i>Erioderma pedicellatum</i>	Yes	SARA – Endangered ESA – Vulnerable	On trunks and branches of coniferous trees in moist, mature forests	In FMD 6	Loss of habitat	Suitable sites for potential Boreal Felt Lichen colonization adjacent to fertile Boreal Felt Lichen thalli		Habitat for Boreal Felt Lichen ECV

²² [The Status of Tradescant's Aster \(*Symphotrichum tradescantii*\) in Newfoundland and Labrador](#), 2008

²³ [The Status of Vreeland's Striped Coralroot *Corallorhiza striata* var. *vreelandii* in Newfoundland and Labrador](#), 2009

²⁴ [COSEWIC Assessment and Status Report on the Blue Felt Lichen *Degelia plumbea* in Canada](#), 2010

Species	Focal Species	Status	Habitat ^{2,3}	DFA Occurrence ^{4,5}	Threats / Risk from Forest Operations	Current Management	Comments/ Assessment	Decision
						are left to cycle naturally ²⁵		
Red Pine <i>Pinus resinosa</i>		ESA - Endangered	Deep sands or gravel soils	Sandy Lake, Howley, Birchy Narrows, Birchy Lake, West Brook	Harvesting not permitted in red pine stands	No permits issued for red pine		ECV
White Pine <i>Pinus strobus</i>		NatureServe – S3	Moist sandy soils	Yes	Risk from scarring during road construction and harvesting operations	Harvesting of white pine not permitted unless on road or a safety risk		ECV
Black Ash <i>Fraxinus nigra</i>		ESA - Threatened	Wet sites along rivers or the margins of swamps	South of Bonne Bay; Bottom Brook	Negligible; possibly, but very rarely, during road construction	A 30m no-grub zone is required on both sides of a water crossing		ECV
Yellow Birch <i>Betula alleghaniensis</i>		NatureServe – S3	Variety of sites, often in association with Balsam fir and other hardwoods	Relatively common in District 14 (e.g. Bottom Brook, Harry's River), less so in 15 & 16.	Forest resource roads provide access to yellow birch for firewood	No management to date		ECV

²⁵ [A 5 Year \(2006 – 2011\) Management Plan For the Boreal Felt Lichen \(Erioderma pedicellatum\) In Newfoundland and Labrador, 2006](#)

Identified Exceptional Conservation Values (Species at Risk)

Habitat for American Marten (Newfoundland pop'n.)	Wet Forest Habitat for Bobolink and Rusty Blackbird
Habitat for Little Brown Myotis and Northern Myotis	Riparian Habitat for Atlantic Salmon, Banded Killifish, American Eel, Freshwater Fishes, and Song Sparrow
Habitat for Woodland Caribou	Habitat for Plant Species at Risk and Rare and Uncommon Vascular Plant Species (Appendix 6)
Habitat for Harlequin Duck and Barrow's Goldeneye	Habitat for Blue Felt Lichen
Mature Coniferous Habitat for Gray-cheeked Thrush, Northern Goshawk, Northern Saw-whet Owl, Olive-sided Flycatcher, Red Crossbill, Winter Wren	Habitat for Boreal Felt Lichen
Open Forest Habitat for Northern Hawk-owl and Northern Shrike	Red Pine
Open Country Habitat for Common Nighthawk, Peregrine Falcon, Rough-legged Hawk and Short-eared Owl	White Pine
All-age Classes Forest Habitat for Evening Grosbeak and Sharp-shinned Hawk	Black Ash
Shoreline Habitat for Bank Swallow, Piping Plover and Red Knot	Yellow Birch

Identified Exceptional Conservation Values (Focal Species)

Habitat for American Marten, Newfoundland population

Habitat for Little Brown Myotis and Northern Myotis

Habitat for Woodland Caribou

Mature Coniferous Habitat for Gray-cheeked Thrush, Northern Goshawk, Northern Saw-whet Owl, Olive-sided Flycatcher, Red Crossbill, Winter Wren

Open Barrens and Grasslands Habitat for Short-eared Owl

Wet Forest Habitat for Bobolink and Rusty Blackbird

Habitat for Blue Felt Lichen

Habitat for Boreal Felt Lichen

2. Does the forest contain a globally, nationally, or regionally significant concentration of endemic species?

Endemic means being unique to a particular geographic location. Assessing the DFA for endemic species ensures the maintenance of vulnerable and/or irreplaceable elements of biodiversity.

Sources of Assessment Information

World Wildlife Fund Ecoregion Conservation Assessment

Conservation International

Birdlife International

Globally Significant Concentrations of Endemic Species

The World Wildlife Fund initiated Global 200 to create a global strategy to protect representative examples of all the world's ecosystems, as well as those areas that contain exceptional concentrations of species and endemics (Olson and Dinerstein 2002). The prioritization identified 238 ecoregions around the globe. The Canadian Boreal Forest was selected as one of the Global 200 ecoregions; however, the defined area did not include the island of Newfoundland. Olson and Dinerstein (2002) rated the Canadian Boreal Forests as "relatively stable or intact".

Conservation International has identified biodiversity hotspots characterized both by exceptional levels of plant endemism and by serious levels of habitat loss. To qualify, a region must meet two strict criteria: it must contain at least 1,500 species of vascular plants (> 0.5 percent of the world's total) as endemics, and it has to have lost at least 70 percent of its original habitat (CI 2010). There are no "hotspots" identified in Newfoundland, or even in Canada.

Birdlife International maintains a World Bird Database to assist in the management of avian species. Endemic Bird Areas is one group of data identified. Birdlife International does not identify any Endemic Bird Areas in Canada (BI 2015).

Endemic Subspecies on the DFA

There are endemic sub-species occurring on the DFA, however, these endemic sub-species do not occur in concentrations, nor would they together constitute a globally or nationally significant concentration. American Marten and the Red Crossbill are endemic sub-species on the DFA.

American Marten have a demonstrated sensitivity to forest operations, in that intensive harvesting over an extensive area, i.e., marten home-range size areas of 10-20 km², would reduce or eliminate marten habitat on the area (Hearn et al 2010). They also found that martens used a broad range of habitat types, including recent cuts ≤5 yr old, regenerating forest <6.5 m, precommercially thinned stands, and mature and overmature (>80-yr) forest. Critical habitat for marten has been identified by the Newfoundland Wildlife Division, and some is within and adjacent to the DFA, specifically the subpopulations in Little Grand Lake, Georges Lake/Pinchgut Lake, and Main River.

The effect of forestry operations on Red Crossbills is not well known. While forest operations shorten rotation ages for conifers in Newfoundland, the overall availability of forest stands of cone-bearing age (i.e., 40+ years of age) is not a concern. Furthermore, forest management policy (Government of Newfoundland & Labrador 2014) requires at least 15% forest cover of stands 81+ years. Finally, fragmented landscapes created by forestry practices may not be a significant threat to the Red Crossbill subspecies, as the forest in Newfoundland is naturally fragmented (Env. Canada 2006).

Identified Exceptional Conservation Values (Endemic Species) No ECV

Note: Information used in the assessment of candidate ECVs for the remaining questions (3–19) can be found in Appendix 1.

²⁶ Shelley Moores, Sr. Mgr. Endangered Species and Biodiversity
Emily Herdman, Ecosystem Management Ecologist, Endangered Species
Claudia Hanel, Ecosystem Management Ecologist, Botanist

3. Does the forest include critical habitat containing globally, nationally or regionally significant seasonal concentration of species (one or several species, e.g., concentrations of wildlife in breeding sites, wintering sites, migration sites, migration routes or corridors – latitudinal as well as altitudinal)?

Habitat containing seasonal concentrations of species refers to areas such as breeding grounds, calving and wintering areas, and migration routes/corridors. This indicator addresses wildlife habitat requirements critical to maintaining population viability (regional “hotspots”).

Sources of Assessment Information

Bird Studies Canada

Conservation International

Ducks Unlimited

World Database on Protected Areas

Interviews with local experts, NL Dept. of Fisheries and Land Resources ²⁷

Areas of seasonal concentrations of birds, such as Important Bird Areas (IBA) and Ramsar Wetlands of International Importance have been designated by international agencies. The IBA Program is a conservation initiative coordinated by BirdLife International. The co-partners for the IBA Program in Canada are Bird Studies Canada and Nature Canada. Of the 25 IBAs identified on the island of Newfoundland, 4 are within 10 km of the DFA.

Middle Ridge Wildlife Reserve (adjacent to Bay Du Nord Wilderness Reserve) shares ~7 km (6%) of its northwestern border with the DFA. Together the two reserves cover 3513 km². Middle Ridge Wildlife Reserve is an upland plateau of extensive barrens, kalmia heaths, bogs, and fens which provides calving and wintering grounds for Woodland Caribou, and breeding grounds for a number of waterfowl species. The Codroy Valley (33 km²) lies within 8 km of the DFA, and is essentially a balsam fir forest mixed with yellow birch and mountain maple thickets. Two restricted range species, Red Crossbill and Ovenbird, (although not in seasonal concentrations) are found in coniferous and deciduous forest respectively, and other species uncommon or absent in the rest of Newfoundland can be seen. Gros Morne National Park (1943 km²), a federally designated conservation area, is also designated as an IBA. The DFA is adjacent to Gros Morne National Park on the park's southern and southeastern borders. The Park is a combination of mountainous barrens, forest, and diverse coastal lowlands and provides calving and wintering grounds for Woodland Caribou. Of the 207 species of birds recorded in the Park, 3 are restricted-range species: Rock Ptarmigan, Red Crossbill, and Ovenbird.

The Codroy Valley Estuary IBA (30 km²) is an important breeding and staging site for numerous waterfowl species in Newfoundland. West of the Codroy Valley IBA, the Estuary lies approximately 7 km southwest of the boundary of the DFA in FMD 14. This area is also a Ramsar Wetlands of International Importance (known as Grand Codroy Estuary), chosen for the diversity of the waterfowl present in migration rather than the numbers. The Ramsar boundaries lie within the Codroy Valley Estuary IBA.

²⁷ Shelley Moores, Sr. Mgr. Endangered Species and Biodiversity
Bruce Rodrigues, Ecosystem Management Ecologist, Biodiversity

Another IBA site, Terra Nova National Park (396 km²), is ~ 27 km from the DFA. This park is forested with black spruce, balsam fir, white birch, and tamarack, and also features numerous lakes and upland plateau fens and bogs. Red Crossbill, Ovenbird, and other forest birds are common as well as shorebirds, waterfowl and alcids.

Ducks Unlimited

Canada has a conservation agreement with CBPPL for wetland and associated upland areas in Cook's Marsh, which falls within the DFA. This area has undergone restoration activities including construction of a water control structure and fishway. Cook's Marsh is valuable for waterfowl and the area is used by many local outdoor recreation enthusiasts. Breeding species include American black duck, green winged teal, ring necked duck, common goldeneye, and common merganser. The area is also an important staging area for black duck. Two properties adjacent to CBPPL managed lands in the Northwest Gander River area (Long Island and The Narrows) consist of wetland and associated upland habitat. These areas are important breeding habitat for dabbling ducks such as black ducks and ring-necked ducks, and for Canada geese. The locations of these areas have been mapped with CBPPL's DFA (Map 1).

Identified Exceptional Conservation Values (IBAs and Ramsar Sites) –

Gros Morne National Park

Cook's Marsh

Middle Ridge Wildlife Reserve (No additional management required)

The DFA contains habitat for seasonal concentrations of Woodland Caribou. Eight caribou herds are found on the DFA: Gros Morne, Aides Pond, Hampden Downs, Gaff Topsails, Buchans, Mount Peyton, Pot Hill, and Middle Ridge. They are found on the forest or adjacent areas year round. The Provincial Wildlife Division has indicated that the most sensitive periods for caribou are calving/post-calving, wintering, and migration. The division has developed woodland caribou core and calving areas and identified them for both conservation and restoration herds. The mapping includes no go, restricted road building, and seasonal restricted areas. All 5-Year Plans must be approved by government prior to harvest including in or adjacent to caribou habitat.

The DFA also contains spawning habitat for Atlantic Salmon (*Salmo salar*). They live in the sea and spawn in freshwater streams, in gravel-bottom rapid areas above or below a pool (DFO 2010). The young stay in fresh water for 2-3 years, journey to the sea for a year or two before returning to spawn in freshwater.

Identified Exceptional Conservation Values (Seasonal Concentration of Species)

Woodland Caribou

Atlantic Salmon

4. Does the forest contain critical habitat for regionally significant species (e.g., species representative of habitat types naturally occurring in the management unit, focal species, or species declining regionally)?

Meta-population is defined by the Encyclopaedia Britannica as “a regional group of connected populations of a species... continually being modified by increases (births and immigrations) and decreases (deaths and emigrations) of individuals, as well as by the emergence and dissolution of local populations contained within it.” Meta-population viability is affected by the population trends of regional species.

Sources of Assessment Information

Atlantic Canada Conservation Data Center (ACCDC)

Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)

International Union for Conservation of Nature (IUCN) Red List of Threatened Species

NatureServe Explorer

Committee on the Status of Endangered Wildlife in Canada (COSEWIC)

Species Status Advisory Committee (SSAC)

Interviews with local experts, NL Dept. of Fisheries and Land Resources²⁸

Regionally significant species were determined using the same criteria as for species at risk in Question 1. Therefore, regionally specific species are those listed in Table 4.

Population trends in the species listed in Table 4 are not well documented for all species. Based on the information available, a number of the species listed in Table 4 are in decline regionally: Woodland Caribou, Barrow's Goldeneye, Harlequin Duck, Peregrine Falcon, Red Crossbill, Red Knot, Rusty Blackbird, and American Eel. However, these population declines are not a result of forest management, or solely forest management. The decline of Woodland Caribou on the island of Newfoundland has been primarily associated with declines in calf recruitment affected by increased mortality of calves by predators, e.g., Black Bears (*Ursus americanus*), coyotes (*Canis latrans*), and Golden eagles (*Aquila chrysaetos*) (Mahoney and Weir 2009). Although harvesting could impact calving and wintering areas Forest Management Guidelines for Woodland Caribou (*Rangifer tarandus caribou*) for the Island of Newfoundland, implemented since October 2006, are being followed to ensure sufficient caribou habitat. The breeding areas of Barrow's Goldeneye and Harlequin Duck are currently protected by the required buffers on waterways. Peregrine Falcon, Red Knot, and Rusty Blackbird do not frequent forests, preferring coastlines, estuaries, and shores of streams, bogs, and wet areas. The exact reason for the decline of Red Crossbill is currently unknown, (although red pine stands are important sources of cones) and the Recovery Strategy is focused on reducing the knowledge gap.

It appears American Marten populations are slowly recovering on the island of Newfoundland (Hearn per com.). A map of the critical habitat for the marten can be seen in the *Recovery Plan American marten (Martes americana atrata) in Newfoundland*, and we have mapped the critical habitat in relation to the DFA (Map 2).

Identified Exceptional Conservation Values (Critical Habitat)

Habitat for American Marten

²⁸ Shelley Moores, Sr. Mgr. Endangered Species and Biodiversity
Bruce Rodrigues, Ecosystem Management Ecologist, Biodiversity

5. Does the forest support concentrations of species at the edge of their natural ranges or outlier populations?

Vulnerability against range contraction and potential genetic variation at range edge are relevant conservation issues. Outlier and edge of range populations may also play a critical role in genetic/population adaptation to global warming.

Sources of Assessment Information

Atlantic Canada Conservation Data Center (ACCDC)
Committee on the Status of Endangered Wildlife in Canada (COSEWIC)
Species Status Advisory Committee (SSAC)
Interviews with local experts²⁹

Most North American endemic species are at their easterly range in Newfoundland. The majority of these species found on the DFA is not at risk, and is widely distributed and found abundantly throughout the remainder of the island of Newfoundland.

There are, however, commercial tree species on the DFA that have been categorized by ACCDC as at risk: white pine, red pine, black ash, and yellow birch are all species at the northeastern limit of their range, and are considered rare. White pine has declined in abundance since the late 1800s, initially due to harvest for lumber, and subsequently by the introduction at the turn of the century of the white pine blister rust fungus (*Cronartium ribicola*). To prevent further decline, provisions were implemented in 1999 and white pine was protected. No permits are issued for the harvest of red pine and black ash. However, permits are issued by the provincial government for the harvest of hardwood species including yellow birch, and they are a preferred browse species for moose.

The status of red pine in the province is Endangered. Natural populations of red pine exist on or adjacent to CBPPL limits, at West Brook Ecological Reserve and Sandy Lake. There also exist areas where it is suspected that healthy white pines of various ages are interspersed with other naturally existing species, in the Grand Lake South operating area.

Arctic hares (*Lepus arcticus*) are at the southern limits of their continental range on the island of Newfoundland. They are found primarily in upland alpine forests, at elevations >1500m and in rugged topography. It is unlikely they occur in productive forests of commercial interests, so there is likely very little potential conflict with forest harvesting.

Identified Exceptional Conservation Values (Edge of Range/Outlier Species)

Red Pine
White Pine
Yellow Birch
Black Ash

6. Does the forest lie within, or adjacent to, or contain a conservation area: a) designated by an international authority,

²⁹ Claudia Hanel, Ecosystem Management Ecologist, Botanist, NL Dept. of Fisheries and Land Resources
Basil English, Supervisor, Silviculture and Research Section, NL Dept. of Fisheries and Land Resources
Dr. Brian Hearn, Wildlife Ecologist, Canadian Forest Service, Natural Resources Canada

- b) legally designated or proposed by relevant federal/provincial/ territorial legislative body, or,
c) identified in regional land use plans or conservation plans?**
-

This indicator looks at designated conservation areas within or adjacent to the forest, to ensure compliance with the intent of the conservation area, and to ensure regionally significant forests are evaluated for consistency with the conservation intent.

Sources of Assessment Information

Canadian Commission for UNESCO

Wetlands International

Parks Canada

Canadian Heritage Rivers System

Nature Conservancy of Canada

NL Parks and Natural Areas

2014-2024 Provincial Forest Management Strategy

Humber Valley Regional Planning Advisory Authority

Interviews with local experts, NL Dept. of Fisheries and Land Resources ³⁰

Internationally Designated Conservation Areas

The United Nations Educational, Scientific and Cultural Organization (UNESCO) seeks to encourage the identification, protection and preservation of cultural and natural heritage around the world considered to be of outstanding value to humanity. Identified areas are called World Heritage Sites, where industrial activities are not permitted. The Canadian Commission for UNESCO lists Gros Morne National Park as a World Heritage Site for its geological features. The DFA is adjacent to Gros Morne National Park on the park's southern and southeastern borders.

The Convention on Wetlands of International Importance, called the Ramsar Convention, is an intergovernmental treaty. The Convention's mission is "the conservation and wise use of all wetlands through local and national actions and international cooperation, as a contribution towards achieving sustainable development throughout the world". Wetlands International lists Grand Codroy Estuary as the only Ramsar site in Newfoundland. The DFA is ~ 7 km from the estuary, but potential identified threats do not include forest management activities.

The International Biological Program (IBP) (1964-1974) identified sites across the globe which represented the world's major ecosystems, giving priority to those sites that are most vulnerable to disturbance. These areas have no formal protection. Five of the seventy-seven identified IBP sites in Newfoundland fall on the DFA:

Serpentine Lake shares a 12 km boundary with the DFA on the southeast corner.

Crooked Bog is completely surrounded by the DFA.

Brownmore Bog is completely surrounded by the DFA.

Sandy Lake lies within the DFA and is bounded by the lake.

Grand Lake Brook is completely surrounded by the DFA.

Identified Exceptional Conservation Values (International Conservation Areas)

Gros Morne National Park

Serpentine Lake IBP Site

³⁰ Jeri Graham, Manager, Natural Areas Program

Crooked Bog IBP Site
Brownmore Bog IBP Site
Sandy Lake IBP Site
Grand Lake Brook IBP Site

Federally/Provincially Designated Conservation Areas

Federal/National

Gros Morne National Park is also a federally designated conservation area, part of the National Parks system. National Parks are a “country-wide system of representative natural areas of Canadian significance, and are protected by law to be maintained in an unimpaired state for future generations”. Approximately 17% of the Gros Morne National Park boundary borders the DFA.

The Canadian Heritage Rivers System, a national river conservation program, promotes, protects and enhances Canada’s river heritage, and ensures that Canada’s leading rivers are managed in a sustainable manner. The Canadian Heritage Rivers System designated Main River as a Canadian Heritage River in 2001. The Main River Canadian Heritage River is comprised of a 152 km² river corridor (the Main River Waterway Provincial Park), and a Special Management Area of 49 km². Currently, industrial activities are not permitted in the Park, and it is expected the same restrictions will soon apply for the Special Management Area. The northwestern tip of Main River Canadian Heritage River lies ~1km northeast of a small outlier parcel of the DFA, and the southeastern tip is within ~2 km of the DFA.

The Nature Conservancy of Canada (NCC) is a private organization that “works for the direct protection of Canada’s biodiversity through the purchase, donation, or placement of easements on ecological significant lands”. NCC has identified the Southwest Newfoundland Natural Area (the boundaries of the Western Newfoundland Forest Ecoregion) as one of the most diverse and significant natural regions in the province. Within the Southwest Newfoundland Natural Area, NCC has purchased properties that are included in/defined by the following conservation areas: the Grand Codroy Estuary Ramsar site mentioned previously, Sandy Point, and The Grassy Place. In relation to the DFA, the Sandy Point area lies ~12km north, and the Grassy Place is ~10km east. Another NCC property, Lloyd’s River, is 48km east of the DFA. NCC’s Natural Area Conservation Plan for the Southwest Newfoundland Natural Area outlines seven biodiversity targets. Three of the targets (representative forests (balsam fir), riparian and wetlands ecosystems, and Newfoundland marten) are considered threatened by forest harvesting, while the remaining (limestone and serpentine heath barrens, sandy beaches and dunes, piping plover, and salt marshes) are threatened by access which could include forest resource roads. NCC has also identified old growth forests as another value in existing and proposed Natural Area Conservation Plans.

Provincial

In Newfoundland and Labrador, there are four main types of protected areas for which the Province is responsible: provincial parks, wilderness reserves, ecological reserves, and wildlife reserves. The protection of these areas is legislated under a variety of Acts, and industrial activities are not permitted in all of the protected areas within the DFA, with the exception of Special Management Areas outside the Main River Waterway Provincial Park. The following provincial protected areas are within or adjacent to the DFA:

West Brook Ecological Reserve (both parcels) are inside the DFA.

Little Grand Lake Provisional Ecological Reserve is adjacent to the DFA. The western boundary of the south block of the Reserve borders on the DFA for ~ 16 km and the southeastern tip of the north block of the Reserve borders on the DFA for ~ 55 km; 50% of the reserve boundary borders the DFA

Little Grand Lake Wildlife Reserve shares a 1 km boundary with the DFA in the south block and ~ 6 km of boundary in the north block; 19% of the reserve boundary borders the DFA

Middle Ridge Wildlife Reserve shares 7 km of its northwestern border with the DFA; 6% of the reserve boundary borders the DFA

Flatwater Pond Provincial Park Reserve lies completely within the DFA.

Jonathan's Pond Provincial Park Reserve is almost totally within the DFA; 76% of the park boundary borders the DFA.

Barachois Pond Provincial Park is adjacent to the DFA sharing ~3 km along the eastern border of the park; 19% of the park boundary borders the DFA.

Sir Richard Squires Memorial Provincial Park is surrounded on the north, east, and south by the DFA, sharing ~ 15 km of boundary; 82% of the park boundary borders the DFA.

T'Railway Provincial Park runs through the DFA in most forest management districts; 25% of the park boundary borders the DFA. (The T'Railway Provincial Park is essentially a linear recreational trail with little conservation value *per se*.)

Main River Waterway Provincial Park and Special Management Area – see above.

The Department of Environment and Climate Change has a Natural Areas Program that is responsible for administration and management of Newfoundland and Labrador's wilderness and ecological reserves, provincial protected area network planning, and the Canadian Heritage Rivers Program, whose vision is "To protect, in an unimpaired condition, large wilderness areas, representative areas of all provincial ecoregions, and areas that contain rare natural phenomena, in order to preserve the diversity and distinctiveness of the Province's rich natural heritage and to support an ecologically sustainable future for the benefit of present and future generations" (ECC 2025).

A "Connectivity Working Group" did exist between 2000 and 2006, bringing together stakeholders to develop scientifically-based solutions that ensured Gros Morne National Park remains ecologically connected to its broader landscape, much of which was in the DFA. In March 2001, CBPPL ended clear cutting in the adjacent Main River Watershed and several months later, in May, the Minister of Canadian Heritage approved the designation of the Main River as a Canadian Heritage River.

Conditions for harvesting near any protected area are contained in the Certificate of Managed Land issued by the Department of Natural Resources. This certificate is required before CBPPL can conduct any forestry activities. The Certificate of Managed Land states: *All roads proposed within 500m of a protected or proposed protected area must be referred to Parks and Natural Areas Division prior to approval.*

Two of the provincial parks/reserves listed above (Main River Waterway Provincial Park/ Special Management Area, and Little Grand Lake Wildlife Reserve/ Little Grand Lake Provisional Ecological Reserve) were once part of the timber limits of CBPPL. As a result of concerns from outfitters and the public over the harvesting of old growth forest in the Main River area, CBPPL announced a no-harvesting policy for a large area along the main stem of the river. This led to the acceptance of the designation of the Main River as a Heritage River, and later its designation as a Canadian Heritage River (the Main River Waterway Provincial Park). A Special Management Area was also delineated around the no-harvesting zone where there would be no clearcut harvesting. Although harvesting trials were conducted in the watershed in 2001, outside of the Provincial Park and special management area, there was no further harvesting in the entire watershed after that date. In 2010, the timber rights for entire Main River watershed were sold back to the Crown.

Little Grand Lake Wildlife Reserve, Little Grand Lake Provisional Ecological Reserve, and Glover Island Public Reserve were all also once part of CBPPL timber limits. In 1985, CBPPL was asked by the provincial government to exclude these areas in any of their harvesting plans in the future, so they could become reserves to protect habitat, areas with white pine, and ecological diversity. CBPPL agreed, and the areas became "provisional" reserves. In 2010 CBPPL's timber rights for these areas and for an additional corridor along the north side of Grand Lake were returned to the Crown as part of a larger sale of timber rights. This enabled the progression from provisional reserves to legal reserves.

A map of the protected and conservation areas on and near the DFA is in Appendix 8 (Map 3).

In 2014, the Centre for Forest Science and Innovation (then of the NL Department of Natural Resources), released the 2014-2024 Provincial Sustainable Forest Management Strategy. As an important component of planning for ecological values, the Forest Service Branch identified 3,993,431 ha of Large Intact Landscape (LIL) on the island of Newfoundland. Industrial forest harvest activity, defined as involving the construction of harvest roads, the use of mechanical harvesting equipment, and the creation of cuts over five hectares, was deferred in this area for a period of 10 years (2014-2024). However, the deferral period has expired and harvesting can now occur in these areas.

Identified Exceptional Conservation Values (Federal/Provincial Conservation Areas)

Gros Morne National Park
West Brook Ecological Reserve
Little Grand Lake Provisional Ecological Reserve
Little Grand Lake Wildlife Reserve
Middle Ridge Wildlife Reserve
Flatwater Pond Provincial Park Reserve
Jonathan's Pond Provincial Park Reserve
Barachois Pond Provincial Park
Sir Richard's Squires Provincial Park
T'Railway Provincial Park

Regional Land Use or Conservation Plans

A portion of CBPPL's DFA resides within the Humber Valley Regional Planning area. Although the timber rights for the whole Humber Valley were owned by CBPPL until 2010, no harvesting has occurred in the area since 2005 when CBPPL voluntarily ceased operations there. The main concern in this area was the viewscape. The timber rights to a corridor from Deer Lake to Gros Morne National Park was relinquished to the Crown for similar reasons at the same time.

The Humber Valley Regional Planning Advisory Authority released a draft Humber Valley Regional Land Use Plan in January 2011. The Plan's proposed land use policy for forestry stated "Existing practices and policies concerning domestic harvesting on Corner Brook Pulp and Paper Ltd. Limits will remain the same". There have been no further developments and it appears the authority is no longer active.

Identified Exceptional Conservation Values (Regional Conservation Areas)

None identified.

CATEGORY 2) FOREST AREAS CONTAINING GLOBALLY, REGIONALLY OR NATIONALLY SIGNIFICANT LARGE LANDSCAPE LEVEL FORESTS, CONTAINED WITHIN, OR CONTAINING THE MANAGEMENT UNIT, WHERE VIABLE POPULATIONS OF MOST IF NOT ALL NATURALLY OCCURRING SPECIES EXIST IN NATURAL PATTERNS OF DISTRIBUTION AND ABUNDANCE.

7. Does the forest constitute or form part of a globally-, nationally-, or regionally-significant forest landscape that includes populations of most native species and sufficient habitat such that there is a high likelihood of long-term species persistence?

Large landscape level forests must be large enough to potentially support most or all native species, and so that long-term, large-scale natural disturbances can take place without losing their resilience to maintain the full range of ecosystem processes and functions.

Sources of Assessment Information

Global Forest Watch

Newfoundland Forest Inventory

Interviews with local experts, NL Dept. of Fisheries and Land Resources³¹

Newfoundland Regional Working Group of the Canadian Boreal Agreement

Global Forest Watch (GFW) defines an intact forest landscape as a contiguous mosaic of natural ecosystems in a forest ecozone, essentially undisturbed by human influence, including both treed and naturally treeless areas (Lee et al, 2010). An intact forest landscape must be large enough to contain and support natural biodiversity and ecological processes, and to provide a buffer against human disturbance from surrounding areas. In their Canadian study using high resolution satellite imagery, GFW examined forest tracts of 50,000 hectares or larger that were at least 10 kilometres wide (intact forest landscapes). The study found that most intact forest landscapes are found in northern Canada and at higher elevations in western Canada: Quebec, the Northwest Territories, Ontario, and British Columbia contain two-thirds of these large landscape forests. Newfoundland, with 6.9% of Canada's total intact forest landscapes, was shown as having 86.3% of its forest as intact forest landscape.

Using parameters outlined in Lee et al (2010), CBPPL conducted a GIS exercise with provincial forest inventory data on the timber limits of CBPPL, to identify intact forest landscape. The inventory data provided a higher standard of reliability than the use of high resolution satellite imagery. The results showed no intact forest landscape of 50,000ha or over on CBPPL's DFA (see Table 5 question 10).

Protected areas exist within the island of Newfoundland, and those of note with reference to landscape level forests are the Bay du Nord Wilderness Reserve (289,500 ha) and three reserves surrounding Little Grand Lake (a total of 147,700 ha), previously part of the DFA: Little Grand Lake Provisional Ecological Reserve, Little Grand Lake Wildlife Reserve, and Glover Island Public Reserve. The first two of these areas are adjacent to CBPPL limits.

Protected areas planning has previously been undertaken by CBPPL and two local environmental organizations, Canadian Parks and Wilderness Society and Canadian Boreal Initiative, through the Canadian Boreal Forest Agreement, with the help of local experts (provincial and federal governments, and Indigenous groups). This Newfoundland Regional Working Group identified potential benchmarks – large, intact, hydrologically-connected protected areas – whose size criteria is set by the natural disturbances fire, spruce budworm, and hemlock looper. The Group also identified conservation features of interest.

Identified Exceptional Conservation Values (Intact Forest Landscape)

No ECV

CATEGORY 3) FOREST AREAS THAT ARE IN OR CONTAIN RARE, THREATENED OR ENDANGERED ECOSYSTEMS.

8. Does the forest contain naturally rare ecosystem types?

³¹ Jeri Graham, Manager, Natural Areas Program

These forests contain many unique species and communities that are adapted only to the conditions found in these rare forest types.

Sources of Assessment Information

Conservation International
World Wildlife Fund Ecoregion Conservation Assessment
Atlantic Canada Conservation Data Center (ACCDC)

Conservation International does not identify any biodiversity hotspots or areas of conservation concern in Canada. NatureServe has nothing identified in the database for Newfoundland as G1, G2, or G3 (critically imperiled, imperiled, or vulnerable). The Atlantic Canada Conservation Data Centre does not yet have information on an ecosystem or community basis for Newfoundland.

In the absence of any rare ecosystem types identified locally, the FECV Assessment Committee previously suggested parameters that could be used to search the provincial forest inventory for rare ecosystem types. These could include Damman forest types (based on soil moisture and fertility), seral stages, intact forest with no history of harvesting (age class 5+ and a considerable distance from habitation), and slope class.

Identified Exceptional Conservation Values (Rare Ecosystem Types)

No ECV at this time

9. Are there ecosystem types within the forest or ecoregion that have significantly declined?

This indicator concerns vulnerability and meta-population viability, and includes anthropogenically rare forest ecosystem types (e.g. late seral red and white pine in eastern Canada).

Sources of Assessment Information

World Wildlife Fund Ecoregion Conservation Assessmentⁱ
Interviews with local experts³²

While the World Wildlife Fund Ecoregion Conservation Assessment and the Parks and Natural Areas division have identified no ecosystem types on the DFA that have significantly declined, there are individual species of trees that have declined in abundance, namely white pine and red pine. As discussed in question 5, red pine is protected as endangered in Newfoundland and Labrador. There also exists an area where CBPPL suspects healthy white pines of various ages occur, interspersed with other naturally existing species, in the Grand Lake South operating area. This particular area may more closely resemble the forest before white pine was harvested for lumber and before the arrival of white pine blister rust. This very rich site is wet and not easily accessed, which is why it was never harvested. It boasts very tall trees and plants uncommon elsewhere in the province. The boundaries of this area have not been specifically outlined nor has there been a detailed survey of the area.

During the assessment process, there was much discussion about the definition of “ecosystem types” as the definition is somewhat broad. To determine ecosystem types in decline, suggestions were made for parameters such as, Damman forest types (based on soil moisture and fertility), seral stages, intact

³² Dave Poole, Forester

forest with no history of harvesting, anything age class 5 + and some distance from habitation, and slope class.

Identified Exceptional Conservation Values (Ecosystem Types in Decline)

Sandy Lake Red Pine

Grand Lake South Area (balsam fir/spruce/healthy white pine of all ages)

10. Are large landscape level forests (i.e., large unfragmented forests) rare or absent in the forest or ecoregion?

In regions or forests where large functioning landscape level forests are rare or do not exist, many of the remnant forest patches require consideration as potential ECVs (best of the rest).

Sources of Assessment Information

Newfoundland Forest Inventory

This attribute applies to forests where large landscape level forests are absent. A GIS exercise using provincial forest inventory data of the timber limits of CBPPL identified no intact forest landscape over 50,000ha. However, unfragmented forest areas (both treed and naturally treeless) of significance were found on the DFA (table below). A map of these areas can be seen in Appendix 8 (Map 5).

Table 5. Area of unfragmented forest on CBPPL's DFA.

Area	Hectares
Cat Arm	12,713
Hampden Downs	25,511
Little Codroy	12,252
Total	50,476

Identified Exceptional Conservation Values (Unfragmented Forest Areas)

Cat Arm

Hampden Downs

Little Codroy

11. Are there nationally/regionally significant diverse or unique forest ecosystems?

This indicator concerns vulnerability, species diversity and significant ecological processes.

Sources of Assessment Information

World Wildlife Fund Ecoregion Conservation Assessment

Interviews with local experts, NL Dept. of Fisheries and Land Resources³³

The World Wildlife Fund has identified 238 ecoregions worldwide (The Global 200), and estimated the conservation status for each one (Olson and Dinerstein 2002). Newfoundland falls within the Boreal Forest /Taiga, Nearctic, Canadian Boreal Forests, whose conservation status is estimated to be relatively stable or intact.

At the regional level however, in the Main River watershed area of the northern peninsula, a portion of the forest is characterized by small-scale, gap-dynamic old-growth boreal forest stands (McCarthy and Weetman 2006). This ecological process differs from balsam fir stands on the rest of the island whose life cycles are naturally insect-driven. The Main River watershed area is currently protected through its designation as the Main River Waterway Provincial Park. Some of the DFA which lies to the north of Main River, the Cat Arm Operating Area, may contain similar examples of this gap-dynamic old-growth boreal forest. This has not been confirmed to date however.

The DFA does include unique geological areas that strongly influence vegetation cover. There are serpentine soils on the DFA in the North Arm Hills and on the Bay D'Espoir Highway near the bridge over the Northwest Gander River. Some of these areas are lightly forested, dominated either by eastern larch and white pine, or by black spruce, but the quality of the trees is low and they are unlikely to be targeted for harvesting. Any rare species would be located in the openings where the serpentine soil is exposed. This type of geology, however, is well represented in the bordering Gros Morne National Park to the northeast, and by Blow Me Down Provincial Park.

There are regionally significant limestone areas on the DFA which have a number of rare plants, several not found on the limestone barrens of the Northern Peninsula. They are found on steep cliffs and the talus slopes underneath, and they would not be forested. In the DFA such areas are located on the south shore of Serpentine Lake, and in the Goose Arm area. Outside the DFA they are found in the Humber Gorge.

Other ecosystems of significance due to high plant diversity and rare plant occurrences are riparian ecosystems. The most significant ones in the DFA include the Upper Humber River, Bottom Brook near the Burgeo Highway turnoff, and Harry's River. However, the significant areas would not be forested, and would be within the buffers on scheduled Salmon Rivers. The estuary of Hughes Brook near Corner Brook would also be significant and inside the DFA, but again not forested.

Another significant ecosystem on the DFA would be red pine stands as red pine is an endangered tree species in Newfoundland. Only found on dry sites in central Newfoundland, these ecosystems are somewhat fragile in the sense that they are dependent on fire for renewal and fire has generally been suppressed. There are a few of these stands in the Howley area, at Birchy Narrows at the east end of Sandy Lake.

Identified Exceptional Conservation Values (Diverse or Unique Forest Ecosystem)

Serpentine Areas – North Arm Hills, Northwest Gander River near bridge on Bay D'Espoir Hwy

Limestone Areas – South shore of Serpentine Lake, Goose Arm area

Riparian Ecosystems – Upper Humber River, Bottom Brook, Harry's River

Red Pine areas – Sandy Lake (Birchy Narrows), Howley

³³ Claudia Hanel, Ecosystem Management Ecologist, Botanist

CATEGORY 4) FOREST AREAS THAT PROVIDE BASIC SERVICES OF NATURE IN CRITICAL SITUATIONS (E.G., WATERSHED PROTECTION, EROSION CONTROL).

12. Does the forest provide a significant source of drinking water?

Although many forests provide a source of drinking water for communities, this indicator is concerned with a potential impact so significant as to be catastrophic, leading to significant loss of productivity, or sickness and death, and there are no alternative sources of drinking water.

Sources of Assessment Information

NL Water Resources Management Division³⁴

Corner Brook Pulp and Paper's DFA covers approximately 1.4 million hectares, and as such covers a large area. It is no surprise then that all or a portion of 36 public water supply areas servicing 32 communities (Table 6) coincide with the DFA (see Map 6, Appendix 8)

Table 6. Communities serviced by public water supply areas occurring on the DFA.

Appleton	Gander Bay South	Pynn's Brook
Baie Verte	Glenwood	Reidville
Benton	Hampden	Seal Cove (White Bay)
Black Duck*	Howley§	South Brook§
Centreville-Wareham-Trinity	Humber Arm South	Steady Brook
Corner Brook	Indian Bay	St. Jude's
Cox's Cove	Irishtown-Summerside	Stephenville*
Deer Lake	King's Point	Stephenville Crossing
Fleur de Lys	Massey Drive	Westport
Gallants	Mount Moriah	Wild Cove*
Gander	Pasadena	

* Unprotected ground water zone

§ Unprotected surface water natural drainage

The majority of drinking water sources on the island are from surface water, due to the abundance of, easy access to, and the reliability of lakes, ponds, and rivers. In order to have their drinking water sources protected, municipal authorities must apply to the provincial Department of Municipal Affairs and Environment, which works through the Water Resources Act to designate Protected Public Water Supply Areas (PPWSA). Anyone wishing to conduct a development activity in a PPWSA must complete an application for a Permit for a Development Activity in a Public Water Supply Area. The permit is processed by the Water Resources Management Division, and sent to the municipal authority responsible for the drinking water supply, which may object to the activity, or it may agree to the activity but add operating conditions to the permit. Commercial forest harvesting may be (and has been) permitted in a PPWSA if it does not impair water quality, and must adhere to the *Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador* which includes a section Guidelines for Forestry Operations in Protected Water Supply Areas. Included are buffers ranging from a minimum of 30m to 150m, depending on the size of the waterbody and the proximity to the intake. Many municipalities require conditions in addition to the guidelines.

³⁴ Annette Tobin, Environmental Scientist
Christa Ramsay, Environmental Scientist

Research has shown that the most significant impacts to forested watersheds following timber harvest are changes in water table levels and stream flow. It is likely that similar changes occur after fire. As a general rule, harvesting impacts on stream flow regimes are usually short-lived and less severe than those brought about by land-use changes, provided that forests soils are protected and vegetation recovery is rapid. In more well-watered areas, rapid revegetation often limits meaningful water yield increases to the first 3-5 years after treatment (Megahan & Hornbeck 2000). Literature reviews also indicate that effects of forest harvesting are negligible on water quality when disturbance levels are below 30% (Kotak et al 2005), and minor on water yield (quantity) at harvesting levels of 30-40% of a watershed area (Rothwell 1997). This level of disturbance is generally used as an upper limit for harvesting in watersheds in forest management plans.

Public water supply areas on the DFA range in size from 0.2 km² to 812 km², servicing populations ranging from a maximum of 22,000 (Corner Brook) to a little as 25. CBPPL sits on the watershed monitoring committees for Corner Brook, Steady Brook, and Gander. Water sampling results from 1989 to the present are available for the Corner Brook watershed, where second-cut harvesting has been on-going since 1987. The nine parameters tested are the same as those tested by the provincial Water Resources Management Division for all drinking water sources. In all the years of testing, the only parameter that did not meet the Canadian Drinking Water Quality Guidelines was color, which is naturally a “tea” color. This applied whether there was harvesting activity or not.

Identified Exceptional Conservation Values (Significant Source of Drinking Water)

All PPWSAs occurring on CBPPL’s DFA as listed in Table 6 were determined to be FECV.

13. Are there forests that provide a significant ecological service in mediating flooding and/or drought, controlling stream flow regulation, and water quality?

Forest areas play a critical role in maintaining water quantity and quality, and breakdown of this service has catastrophic impacts or is irreplaceable.

Sources of Assessment Information

Geological Disasters in Newfoundland and Labrador
NL Water Resources Management Division³⁵

The island of Newfoundland is a mosaic of forest, water, bogs, and barrens with very little cleared land, and the productive forest is naturally highly fragmented. All forests and treed wetlands play a significant role in mediating flooding and/or drought, stream flow regulation, and filtering groundwater and runoff. This indicator, however, deals with ecological services in critical situations.

Flooding is part of natural environmental processes. In Newfoundland it is often caused by heavy rainfalls in combination with rapid melting of snow on the ground, and less frequently by ice jams in rivers, and high tides with storm surges along the coast. (Liverman et al 2001). Two flood risk areas, Black Duck and Appleton, are within CBPPL’s DFA. A number of other flood risk areas such as Stephenville, Steady Brook and Deer Lake are downstream of CBPPL’s DFA. However there is no

³⁵ Dr. Ali Khan, Manager, Hydrologic Modeling Section

information to indicate that the harvested areas in these watersheds at any one time are large enough to have an effect on these flood zones.

Identified Exceptional Conservation Values (Significant Ecological Service)

No ECV

14. Are there forests critical to erosion control?

This indicator relates to the impacts of soil, terrain or snow stability in the control of erosion, sedimentation, landslides, or avalanches.

Sources of Assessment Information

NL Geochemistry, Geophysics and Terrain Sciences Section³⁶

Movement of land mass down slope is influenced by gravity. Landslides are widespread in Newfoundland. They are usually triggered by excess quantities of either rainfall or snowfall to the slope. Other factors that commonly cause mass movement that could be influenced by forest operations include changes in slope angle (road construction across steep slopes), and changes in vegetation (forest harvesting). Generally slopes in excess of 25° (46%) are prone to slope failure, depending on the underlying material.

Identified Exceptional Conservation Values (Erosion Control)

Slopes over 25° (46%)

15. Are there forest landscapes (or regional landscapes) that have a critical impact on agriculture or fisheries?

Forests mediate wind and microclimate at the ecoregion scale, affecting agricultural and fisheries production. Riparian forests play a critical role in maintaining fisheries by providing bank stability, sediment control, nutrient inputs, and microhabitats.

Sources of Assessment Information

There are no significant agricultural production areas within CBPPL's DFA. These areas were removed from the DFA in the sale of limits back to the Crown in 2010. However, watersheds within the DFA provide irrigation on agriculturally developed land, primarily in the Deer Lake, Cormack, Reidville, Howley, Green Bay, and Robinsons to St. Fintan's areas, with smaller hobby type farms dispersed

³⁶ Dr. Martin Batterson, Section Manager

throughout. Given the expanse of forest adjacent to these agricultural areas, no significant impacts are anticipated.

There are many lakes and ponds on the DFA – 7.4% of the limits is in waterbodies of some sort. These waterbodies are home to a number of species of fish that provide a sport fishery for both locals and tourists. Major fishing rivers on the DFA include: Humber, Gander, Harry's, and Indian. Fish habitat is currently protected by riparian widths outlined in the *Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador*.

Identified Exceptional Conservation Values (Critical Impact on Agriculture or Fisheries)

Freshwater fishes
Atlantic Salmon

CATEGORY 5) FOREST AREAS FUNDAMENTAL TO MEETING BASIC NEEDS OF LOCAL COMMUNITIES (E.G., SUBSISTENCE, HEALTH).

16. Are there local communities? Is anyone within the community making use of the forest for basic needs/ livelihoods?

Sources of Assessment Information

CBPPL Sustainable Forest Management Plan, 2011
CBPPL Socio-Economic Impact Assessment Report, 2011

There are many communities (~65) of varying sizes within or in close proximity to the DFA that use and benefit from the forest. Uses/benefits of the forest include: employment (~500 employees directly in the mill, woodlands, and power house), sawlogs for sawmills, tourism and recreation, timber for wharves and boats, firewood for homes and businesses, and cabins, hunting, fishing, trapping, and berry picking. Although some of these uses may be recreational, for a portion of residents these uses fill basic needs. Den Otter & Beckley (2002) found that hunting, berry-picking, fishing, and wood-cutting supplement incomes, particularly in smaller communities.

Identified Exceptional Conservation Values (Critical to Basic Needs)

Recreation
Cutting timber for wharves and boats
Income from working in the forest
Firewood to heat homes and businesses
Company roads for access to hunting, berry picking, and cabins
Sawlogs for local sawmills

CATEGORY 6) FOREST AREAS CRITICAL TO LOCAL COMMUNITIES' TRADITIONAL CULTURAL IDENTITY (AREAS OF CULTURAL, ECOLOGICAL, ECONOMIC OR RELIGIOUS SIGNIFICANCE IDENTIFIED IN COOPERATION WITH SUCH LOCAL COMMUNITIES).

17. Is the traditional cultural identity of the local community particularly tied to a specific forest area?

Sources of Assessment Information

Qalipu Mi'kmaq First Nation Band
Miawpukek First Nation

Indigenous people on the island of Newfoundland are primarily Mi'kmaq. They have identified culturally important areas for burial and sacred sites, spirit areas and medicine plants but maintain proprietary ownership and privacy of the data within their organizations.

Both Miawpukek First Nation and Qalipu Mi'kmaq First Nation engage in the government consultation process for CBPPL's 5-yr. Operating Plans. All Annual Operating Plans are provided to Qalipu Mi'kmaq First Nation for review and feedback. Annual Operating Plans in District 6 are provided to Miawpukek First Nation for review and feedback.

Identified Exceptional Conservation Values (Traditional Cultural Identity)

Burial Sites
Sacred Sites
Spirit Areas
Medicine Plants

18. Is there a significant overlap of values (ecological and/or cultural) that individually did not meet ECV thresholds, but collectively constitute ECVs?

A comprehensive review of candidate ECVs for the DFA was conducted with many of the candidates identified as ECVs. CBPPL feels there is no significant overlap of values that would lead to new ECVs or FECVs.

Identified Exceptional Conservation Values (Collective Overlap of Values)

No ECV

5. MANAGEMENT AND MONITORING STRATEGIES FOR ECV'S AND FECV'S

Management strategies for ECVs and FECVs vary, from total protection of a species or area, to special strategies undertaken that allow harvesting, road building and silviculture operations but with conditions. Some of the management strategies are found in the 5-Year Operating Plans, others in the Sustainable Forest Management (SFM) Plan within Indicator profiles. All applicable management strategies and special considerations are relayed to CBPPL contractors during Pre-work Meetings.

Monitoring of ECV management strategies can be through a number of processes. Superintendents monitor harvesting operations by daily contact with contractors and by checking their progress through on-line operating maps that are updated daily. They also perform on-site monitoring at least weekly. Another process is through the Environmental Management System (EMS) compliance inspections. Contractors monitor their own work through monthly #1 EMS Compliance Inspections. CBPPL conducts a comprehensive #2 EMS Compliance Inspection of harvesting contractor activity twice per year, and road building activity once per year. CBPPL conducts a final #3 EMS Compliance Inspection or a Final Road Inspection after the operating area activity (harvesting or road building) is completed. A Cutover Assessment Survey is done in conjunction with the #3 inspection to determine utilization rates, amount of soil disturbance, and number of wildlife/snag trees. Any ECVs that are to be monitored through the EMS Compliance Inspections and Cutover Assessment Surveys have been added to the inspection forms and guidelines. The annual EMS and SFM Standards internal and external environmental audits also serve as monitoring processes. To determine the effectiveness of the management strategies, in many cases CBPPL relies on agencies with which it has "agreements" related to the Exceptional Conservation Values.

For each ECV/ FECV identified in the previous section, the following table indicates at least one management strategy, the monitoring responsibilities, and the effectiveness of the management strategies.

Table 7. Management and Monitoring Strategies for ECVs and FECVs on Corner Brook Pulp Paper's DFA.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
Habitat for Newfoundland Marten	No net loss of suitable habitat for marten district by district	<p>If proposed operating areas in the 5-Year Plan coincide with areas currently occupied by marten, CBPPL will adhere to directives issued by the provincial Wildlife Division, Endangered Species Section, concerning the <i>Recovery Plan for the Threatened Newfoundland Population of American Marten (Martes americana atrata)</i>. The 5-Year Plan proposal also goes through the Environmental Assessment process. Newly available predictive marten occupancy models are used to assess the proposal.</p> <p>During the prework meeting CBPPL advises harvesting contractors of special concerns in the operating area, such as marten habitat, and what restrictions apply.</p> <p>More detailed management strategies for pine marten are found in the 5-Year Operating Plans, Section 4.2.1, and in the Sustainable Forest Management Plan, Indicator 1.2.2 Pine Marten Habitat</p>	<p>1) Hectares of suitable habitat, based on baseline information, determined by habitat model.</p> <p>2) Evidence of marten occupying adjacent areas.</p>	<p>CBPPL Operations Superintendents monitor their contractors to ensure they are not harvesting outside the designated operating area. This is done through the #2 and #3 EMS inspections, and environmental auditing of Indicator 4.1.5</p> <p>Conservation Officers conduct similar but separate monitoring of CBPPL contractors.</p>	<p>The Gov NL Wildlife Division will determine hectares of suitable habitat and hectares of habitat occupied. (Need to rely on trends as well)</p> <p>CBPPL will ask GovNL Wildlife Division to continue conduct hair snag surveys.</p>
Habitat for Little Brown Myotis and Northern Myotis	Ensure suitable habitat	The wood supply model removes 15% old growth from the productive forest before determining AAC. CBPPL commits to leaving 20% in age class 81+ years.	Percentage of 81+ age class on the DFA	CBPPL Planners will ensure that a minimum of 20% of the DFA is in the 81+ years age class through the 5-Year and Annual Operating Plans.	The GovNL Wildlife Division will assess if enough suitable habitat remains for bats across the landscape.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
Habitat for Woodland Caribou	To ensure that adequate caribou habitat is available at all times on the DFA.	<p>Follow the Forest Management Guidelines for Woodland Caribou (Rangifer tarandus caribou) For the Island of Newfoundland as developed by the provincial Wildlife Division. New guidelines under development have been incorporated into the 5-Year Operating Plans since 2012:</p> <ul style="list-style-type: none"> • No harvesting in Tier 1 core areas unless permissions are given; • No harvesting during calving season in Tier 2 habitat (April 15 – July 15). <p>Through the Canadian Boreal Forest Agreement CBPPL is also involved in the maintenance of self-sustaining woodland caribou populations within the DFA.</p> <p>More detailed management strategies for woodland caribou are found in the 5-Year Operating Plans, and in the Sustainable Forest Management Plan, Indicator 1.2.1/4.1.1 Caribou Habitat</p>	Compliance with caribou guidelines	<p>CBPPL Operations Superintendents monitor contractors to ensure they are not harvesting outside the designated operating area through the #2 & #3 EMS inspections. The Report of Past AOPs, as well as soft updates of cutovers, report on the previous year's harvesting. The environmental auditing process monitors Indicator 1.2.1/4.1.1.</p> <p>Conservation Officers conduct similar but separate monitoring of CBPPL contractors. Additional monitoring required in the new guidelines will be followed.</p>	<p>On an annual basis, the Environment and Climate Change will assess habitat availability on the DFA and on the Island.</p> <p>The Wildlife Division will assess the effectiveness of the guidelines and update the kernels.</p>
Habitat for: Harlequin Duck Barrow's Goldeneye	Protect suitable habitat	<p>Buffers required in the Environmental Protection Guidelines for Ecologically Based Forest Resource Management (Stand Operations Level) cover the management of nesting Harlequins. Sensitive waterfowl habitat is protected through increased buffers of 50m on</p>		<p>CBPPL Operations Superintendents monitor that contractors are leaving the required buffers on waterways through the #2 & #3 EMS inspections.</p> <p>Conservation Officers conduct similar but</p>	<p>Environment Canada's Canadian Wildlife Service (EC-CWS) will assess the effectiveness of existing buffer requirements.</p>

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		<p>breeding, moulting, and staging areas identified by Environment Canada's Canadian Wildlife Service. These will be incorporated into the 5-Year Plans (which are reviewed by the Canadian Wildlife Service) and the Annual Operating Plans. A <i>Management Plan for the Harlequin Duck (Histrionicus histrionicus) Eastern Population, in Atlantic Canada and Québec</i> proposes 30m buffers in waterfowl breeding, moulting, and staging areas.</p> <p>More detailed management strategies for harlequin duck are found in the 5-Year Operating Plan for Districts 15 & 9.</p>		separate monitoring of CBPPL contractors.	
Mature Coniferous Habitat for: Gray-Cheeked Thrush, Northern Goshawk, Olive-sided Flycatcher, Red Crossbill, Winter Wren	Protect suitable habitat	<p>Buffers currently required in the <i>Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador</i> provide some protection for most if not all of the species listed. If additional management strategies are required they will be specified by the Wildlife Division during the 5-Year Planning process, and/or incorporated into the Annual Operating Plans.</p> <p>The wood supply model removes 15% old growth from the productive forest before determining AAC. CBPPL commits to leaving 20% in age class 81+ years.</p>	Percentage of 81+ age class in each forest management district on the DFA.	<p>Through environmental compliance inspections, CBPPL Operations Superintendents will determine if contractors are following the Environmental Protection Guidelines.</p> <p>CBPPL Planners will ensure that a minimum of 20% of the DFA is in the 81+ years age class through the 5-Year and Annual Operating Plans.</p>	<p>The Endangered Species and Biodiversity Section of the Wildlife Division will assess the effectiveness of the management strategy.</p> <p>EC-CWS will also review the effectiveness of the management strategy, particularly for Red Crossbill and Olive-sided Flycatcher.</p>

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		Other habitat management strategies include: management of patch size and shape; raptor nest protection; snag retention and creation; Structural complexity enhancement (thinning); variable retention; and deferred harvesting areas.			
Open Forest Habitat for Northern Shrike	Protect suitable habitat	The wood supply model removes 15% old growth from the productive forest before determining AAC. CBPPL commits to leaving 20% in age class 81+ years.	Percentage of 81+ age class in each forest management district on the DFA	CBPPL Planners will ensure representative amount of all age classes exists on the DFA through the 5-Year and Annual Operating Plans.	The Endangered Species and Biodiversity Section of the Wildlife Division will assess the effectiveness of the management strategy.
Open Barrens and Grasslands Habitat for: Peregrine Falcon ssp. <i>Anatu</i> ;; Short-eared Owl	Protect suitable habitat	CBPPL cannot impact open barrens and grasslands habitat during harvesting operations. When locating forest access roads it is more economically viable for CBPPL planners to follow strings of forest and avoid more expensive road building costs in open barrens (rock substrate) and grasslands (wet areas). To protect habitat for Peregrine Falcon and Short-eared Owl, CBPPL planners will avoid locating forest access	Avoidance of operating in open barrens and grasslands habitat.	CBPPL Planners will avoid locating forest access roads in barrens and grasslands whenever possible.	The Endangered Species and Biodiversity Section of the Wildlife Division will assess the effectiveness of the management strategy.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		roads in open barrens and grasslands. CBPPL follows the <i>Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador</i> . If additional management strategies are required they will be specified by the Wildlife Division during the 5-Year Planning process, and/or incorporated into the Annual Operating Plans.			
All Age-classes Forest Habitat for Sharp-shinned Hawk	Protect suitable habitat	CBPPL will ensure a representative amount of all age classes exists on the DFA.	Representation of all age classes	CBPPL Planners will ensure representative amount of all age classes exists on the DFA through the 5-Year and Annual Operating Plans.	GovNL Forestry Branch conducts a wood supply analysis using the 5YP timing windows (varying between zones) that indicates the age class structure of the forest in Newfoundland and Labrador.
Shoreline Habitat for: Piping Plover; Red knot		CBPPL cannot impact shoreline habitat.			
Wet Forest Habitat for Rusty Blackbird	Protect suitable habitat	Buffers currently required in the <i>Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador</i> provide protection for nests of Rusty Blackbird. If additional management strategies are required they will be specified by the Wildlife Division during the 5-Year Planning process, and/or incorporated into the Annual Operating Plans.	Availability of habitat for these species on the DFA.	Through environmental compliance inspections, CBPPL Operations Superintendents will determine if contractors have left the required buffers on waterways.	The Endangered Species and Biodiversity Section of the Wildlife Division will assess the effectiveness of the management strategy.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
Habitat for: Laurentian Dandelion Rock Dwelling Sedge Serpentine Stitchwort Tradescant's Aster Other S1-S3 Species	Protect occupied habitat and adjacent suitable habitat	<p>Buffers currently required in the <i>Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador</i> provide protection for most of the species listed. If additional management strategies are required they will be specified by the Wildlife Division during the 5-Year Planning process, and/or incorporated into the Annual Operating Plans.</p> <p>The Endangered Species and Biodiversity Section of the Wildlife Division has been identified "hot spots" on the DFA, defining the general location of the rarest of the rare plant species, and areas with an abundance of rare plant species. These "hot spots" will be removed from areas to be harvested during the planning stage.</p>	Availability of habitat for these species on the DFA.	Through environmental compliance inspections, CBPPL Operations Superintendents will determine if contractors have left the required buffers on waterways.	The Endangered Species and Biodiversity Section of the Wildlife Division will assess the effectiveness of the management strategy.
Boreal Felt Lichen	Maintain existing populations of Boreal Felt Lichen.	Follow <i>A 5 Year (2006 – 2011) Management Plan For the Boreal Felt Lichen (Erioderma pedicellatum) In Newfoundland and Labrador</i> by considering a Landscape Management approach based on ecological forest site types whereby suitable sites for potential Boreal Felt Lichen colonization adjacent to fertile Boreal Felt Lichen thalli are left to cycle naturally.	Presence or absence of lichen	<p>Implementation of the management plan will be monitored and evaluated by the Endangered Species Section every year for level of completion and where necessary, resources will be redirected.</p> <p>CBPPL Planners are trained to identify Boreal Felt Lichen and suitable</p>	<p>1) Reconnaissance surveys on areas identified by the Gov NL Forestry Department in the 5-Year Planning Process and review of Annual Operating Plans (DEC, FSB)</p> <p>2) Regeneration surveys will determine if balsam fir is regenerating in</p>

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		<p>CBPPL Planners and Contractor Foremen were trained to identify Boreal Felt Lichen and suitable sites for potential Boreal Felt Lichen colonization. Training occurred where Boreal Felt Lichen is known to occur. A presentation was given to CBPPL staff and contractors on Boreal Felt Lichen identification, habitat and mitigation.</p> <p>In 2013, five sites in Districts 15 & 16 most likely to have Boreal Felt Lichen were identified through mapping exercises. Qualified ecologists spent six working days searching for Boreal Felt Lichen on these sites but none were found.</p> <p>It is expected that priorities may change as the understanding of Boreal Felt Lichen biology and management increases. Memorial University (Grenfell), College of the North Atlantic, NL Government and Canadian Forestry Service (NL) have developed a working predictive model to identify areas of suitable habitat for Boreal Felt Lichen on the central Avalon Peninsula. Future research will investigate the population ecology of Erioderma and other lichen species in the lab and in the field with the aim to restore lichen populations in managed landscapes.</p>		sites for potential Boreal Felt Lichen colonization.	areas with Boreal Felt Lichen (CBPPL)

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		If proposing to harvest in Districts 5 or 6, confer with Endangered Species and Biodiversity Section during planning stage, through submission of proposed Annual Operating Plans.			
Red Pine and Red Pine Stands at: Sandy Lake, Birchy Narrows, and Birchy Lake	Protect existing red pine on the DFA from harvesting by CBPPL.	<p>CBPPL can only protect existing red pine on their limits by following provincial legislation to exclude red pine trees and stands from harvesting plans. In addition, CBPPL has included in their Domestic Wood Cutting Regulations (distributed with each fire wood permit) that cutting of red pine is not permitted</p> <p>More detailed management strategies for red pine are found in the 5-Year Operating Plans.</p>	Incident reports of harvesting red pine on the DFA by CBPPL.	Through environmental compliance inspections and cutover assessment surveys, CBPPL Operations Superintendents will determine if contractors have harvested any red pine on CBPPL limits.	Persistence of red pine on the DFA as indicated by Dept. of Fisheries and Land Resources through the provincial forest inventory or monitoring programs.
White Pine	Protect existing white pine on the DFA from harvesting by CBPPL.	<p>No harvesting of white pine is permitted except in road right-of-ways (ROW) or where they pose a safety hazard. Where possible, leave a 5m "no-machine" zone around white pine >50cm dbh (to protect roots), and leave hardwoods and unmerchantable softwoods within that radius. In addition, CBPPL has included in their Domestic Wood Cutting Regulations (distributed with each fire wood permit) that cutting of white pine is not permitted</p> <p>More detailed management strategies for white pine are</p>	Incident reports of Noncompliance with regulations	Through environmental compliance inspections and cutover assessment surveys, CBPPL Operations Superintendents will determine if contractors have harvested any white pine on CBPPL limits outside of ROWs/safety requirements, and if no-machine zones have been left around white pine >50cm dbh.	Persistence of white pine on the DFA as indicated by Dept. of Fisheries and Land Resources through the provincial forest inventory or monitoring programs.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		found in the 5-Year Operating Plans.			
Black Ash	Protect existing black ash on the DFA from harvesting by CBPPL.	No permits are issued for the harvest of black ash. CBPPL has included in their Domestic Wood Cutting Regulations (distributed with each fire wood permit) that cutting of black ash is not permitted	Incident reports of harvesting black ash on the DFA by CBPPL.	Through environmental compliance inspections and cutover assessment surveys, CBPPL Operations Superintendents will determine if contractors have harvested any black ash on CBPPL limits	Persistence of black ash on the DFA as indicated by FSB through the provincial forest inventory or monitoring programs.
Yellow Birch	No harvesting of birch >50cm dbh when encountered on the DFA.	<i>SOP H-06 Leaving Wildlife Trees for Biological Diversity</i> , to instruct operators to leave yellow birch trees >50cm dbh (as seed trees) when encountered. CBPPL has included in their Domestic Wood Cutting Regulations (distributed with each fire wood permit) that cutting of yellow birch >50cm dbh is not permitted. PCT employees are instructed to favour yellow birch over other hardwoods when thinning.	Incident reports of harvesting yellow birch >50cm dbh on the DFA by CBPPL.	Through environmental compliance inspections and cutover assessment surveys, CBPPL Operations Superintendents will determine if contractors have harvested any yellow birch >50cm dbh on CBPPL limits. On areas of the DFA where hardwood management has been transferred to Crown, FSB staff will monitor for compliance.	Persistence of yellow birch on the DFA as indicated by regeneration assessment surveys.
Ducks Unlimited Conservation Area: Cook's Marsh	Respect the boundaries of this property.	CBPPL has a Stewardship Agreement with Ducks Unlimited Canada in which the area shall be preserved for the enhancement and protection of waterfowl and other wildlife, subject to the condition that CBPPL carry on normal harvesting and other related	Incident reports of noncompliance by CBPPL of the Stewardship Agreement with Ducks Unlimited.	Geospatial data from equipment will be reviewed to ensure that all harvesting and road construction activities are in compliance with the Agreement.	CBPPL will consult with Ducks Unlimited Canada to ensure the Stewardship Agreement aimed at waterfowl protection is not being compromised by CBPPL activities.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		activities in specific designated areas. CBPPL will continue to adhere to this agreement.			
Riparian Areas/Habitat for: Atlantic Salmon; Banded Killifish; American Eel; Freshwater Fishes; Song Sparrows	Maintain integrity of the boundary of riparian areas on the DFA that have been legislated and/or established by agreements and through the 5-year planning process.	CBPPL follows the <i>Environmental Protection Guidelines Forestry Operations in Newfoundland and Labrador</i> to protect the values of riparian areas, and therefore habitat for salmon and freshwater fishes. Management strategies include buffers along streams, no-grub zones at stream crossings, and guidelines for the installation of culverts during road building. The federal <i>Fisheries Act</i> requires an application for authorization be submitted to the Fisheries and Oceans Canada (FOC) for any stream crossing where serious harm to fish cannot be avoided by following standard measures. Environmental Work Instructions, Standard Operating Procedures, ¹ and Measures to Avoid Harm to Fish and Fish Habitat guide employees in the protection of riparian areas. FOC has a self-assessment process to help proponents determine if their work requires a review by FOC. Each project requires an application to be sent and a Letter of Advice (LoA) will be issued to CBPPL by FOC for installation of single culverts ≤ 2000mm in diameter. FOC must be notified ten days prior to the start of the project. A similar	Compliance with buffers that have been legislated or agreed to with organizations.	Through environmental compliance inspections, CBPPL Operations Superintendents will determine if contractors are leaving the required buffers on waterways. Conservation Officers conduct similar but separate inspections of CBPPL contractors. Geospatial harvesting data from FPDat-equipped harvesters will be reviewed to ensure that no harvesting occurs within designated and/or regulated buffers.	Consult with regulatory bodies (DFO, DEC) and agencies with whom CBPPL has agreements, to determine their satisfaction with the effectiveness of the established buffers in maintaining integrity of riparian areas, for the management of salmon and freshwater fish stocks in waterways on CBPPL limits.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		<p>Blanket of Approval from the Provincial Water Resources Management Division (WRMD) requires submission of a Notification Form two days prior to installation.</p> <p>CBPPL leaves a 100m buffer on the main stems of scheduled salmon rivers and specific spawning areas designated by the FOC. CBPPL has also left 30m buffers on smaller tributaries identified by salmon groups such as SPAWN.</p> <p>CBPPL has voluntary agreements with local conservation groups such as the Indian Bay and Freshwater-Alexander Bays Ecosystem Corporations (developed through the 5 Year Planning process) concerning access, road building, and extended buffers (100-200m). Requests for buffers wider than required are assessed on an individual case basis. Buffers for salmon spawning areas, wildlife habitat, etc., wider than required in the EPG, are removed from the timber supply when calculating the Annual Allowable Cut.</p> <p>Sustainable Forest Management Plan Indicator 5.1.4 Access Management Plan details management strategies to ensure waterbodies are protected during road construction. Additional</p>			

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		management strategies for freshwater fishes are found in the 5-Year Operating Plans. Management strategies for riparian areas are found in the 5-Year Operating Plans.			
All Provincial and Federal Parks and Reserves listed as ECV in Question 6.	Respect the boundaries of all provincial and federal parks and reserves.	Management strategies for protected areas are found in the 5-Year Operating Plans, Section 4.2.2. CBPPL will not harvest inside parks and reserves. Also, a 500m “no roads” buffer is to be maintained around all proposed provincial protected areas to reduce access and resulting damage from motorized vehicles.	No harvesting inside the park/reserve and no road construction within the 500m no-road buffer adjacent to a proposed protected area.	CBPPL Planners will ensure that boundaries are clearly indicated and no roads are located within 500m of proposed protected areas through the 5-Year and Annual Operating Plans. Road construction inspections by CBPPL staff will determine if roads are being built within the 500m buffer adjacent to the proposed protected area boundary. Geospatial harvesting data will be reviewed to ensure that no harvesting occurred within the park/reserve.	A review of past annual operations will verify if park boundaries have been respected and if roads were built within 500 m of a proposed protected area, thus respecting the boundary.
Unfragmented Forest Areas (CBPPL); includes old growth	Contribute to the conservation of unfragmented intact forest landscape, including old growth.	CBPPL has identified 3 areas to contribute to an intact forest landscape.	Absence of harvesting in CBPPL’s 3 deferral areas.	CBPPL Planners ensure that no harvesting is scheduled in these areas.	CBPPL deferral areas remain unfragmented and intact during the monitoring period.
Open Serpentine and Limestone Areas	Minimize potential impacts from forest road construction activities on rare	CBPPL consults with the Wildlife Division to identify and digitally map the limestone (i.e., steep cliffs and talus slopes) and	Compliance with recommendations from the Wildlife Division on road	CBPPL Planners will implement management strategies agreed upon with the Wildlife Division.	The Endangered Species and Biodiversity Section of the Wildlife Division will

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
	species in serpentine and limestone areas on the DFA.	serpentine areas of concern. CBPPL works with the Wildlife Division to prevent/minimize the potential impacts on rare species in these areas from road building.	construction in limestone and serpentine areas		assess the effectiveness of the management strategy.
Protected Public Water Supply Areas (PPWSAs)	Prevent/minimize adverse effects from forest harvesting activities to public drinking water supplies on the DFA.	CBPPL will follow the <i>Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador</i> and the PPWSA permit. Management strategies include buffers along streams and no-grub zones at stream crossings. Sustainable Forest Management Plan Indicator 3.2.2 contains a management strategy to limit the proportion of watershed with recent stand-replacing disturbance within the DFA to 25%.	Compliance with the EPG and the PPWSA permit.	CBPPL Operations Superintendents will monitor that all requirements in PPWSAs are being followed by CBPPL contractors, through the #2 & #3 EMS inspections. Indicator 3.2.2 is monitored through the environmental auditing process. Conservation Officers conduct similar but separate monitoring of CBPPL contractors.	CBPPL will consult with Water Resources Management Division; Dept. of Fisheries and Land Resources annually to ensure the management strategy is effective for protecting public drinking water supplies. CBPPL's annual updates information of the SFM Plan, including the regeneration of cutovers within 5 years.
Slopes Greater Than 25° (46%)	To maintain forests valuable to the prevention of erosion.	CBPPL will avoid harvesting on slopes greater than 46% that are prone to slope failure because of the underlying material. There are no recorded incidents of significant slope erosion due to harvesting, and following this restriction will prevent such occurrences.	Absence of harvesting on slopes greater than 46% that are prone to slope failure because of the underlying material	Through environmental compliance inspections, CBPPL Operations Superintendents will determine if contractors avoid harvesting on slopes greater than 46% that are prone to slope failure because of the underlying material.	The NL Geochemistry, Geophysics and Terrain Sciences Section has confirmed that avoiding harvesting on slopes greater than 46% that are prone to slope failure because of the underlying material will prevent soil erosion.
Recreation: access to the wilderness for hunting, fishing berry picking; access to cabins for, e.g., pleasure,	To allow access to the DFA for recreation.	Make CBPPL resource roads 100% accessible to the public, subject to natural reclamation, removal of temporary infrastructure such as bridges and culverts, safety and agreements with agencies.	Amount of resource roads on the DFA under CBPPL control that are accessible to the public.	Through environmental compliance inspections and regular on-site visits, CBPPL Operations Superintendents will determine adherence to the 5-Year Plan and the SFM Plan.	CBPPL's annual updates of the SFM Plan, including the amount of roads accessible to the public.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
relaxation and hunting		<p>Recreation values are managed through the 5-Year Planning Process, and CBPPL's Sustainable Forest Management (SFM) Plan.</p> <p>Sustainable Forest Management Plan Indicator 5.1.3 ensures CBPPL resources roads are available to the public, subject to natural reclamation, removal of temporary bridges and culverts, safety, and agreements. Indicator 1.4.1 identifies Special Places on the DFA that are unique and important to the public. Indicator 5.1.1 tracks and updates agreements with forest-dependent businesses, forest users (recreational groups), and the local community. Management strategies for recreation are found in the 5-Year Operating Plans.</p>			
Income from working in the forest	Where economically feasible and beneficial to all parties, allow opportunities for income associated with the forest on the DFA.	<p>This value is managed through the 5-Year Planning Process, and CBPPL's Sustainable Forest Management (SFM) Plan.</p> <p>Sustainable Forest Management Plan Indicator 5.1.2 helps maintain commercial opportunities of outfitters by limiting the amount and location of harvesting around their camps.</p> <p>Indicator 5.2.3 maintains wages for CBPPL union workers in line with the provincial average.</p>	Individuals and businesses deriving income from working in the forest.	Through environmental compliance inspections and regular on-site visits, CBPPL Operations Superintendents will determine adherence to the 5-Year Plan and the SFM Plan.	CBPPL's annual updates of the SFM Plan, including outfitters with camps on the DFA, wages for CBPPL union workers compared to the provincial average, and agreements with forest-dependent businesses.

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		<p>Indicator 5.1.1 tracks and updates agreements with forest-dependent businesses.</p> <p>Management strategies for ensuring the existence of timber for harvesting are found in the 5-Year Operating Plans.</p>			
Sawlogs for local sawmills	Where economically feasible and beneficial to all parties, make sawlogs available to sawmills.	<p>This value is managed through the 5-Year Planning Process, and CBPPL's Sustainable Forest Management (SFM) Plan.</p> <p>Management strategies for ensuring the existence of timber for harvesting are found in the 5-Year Operating Plans.</p> <p>Sustainable Forest Management Plan Indicator 5.2.1: CBPPL makes a percentage of CBPPL annual harvest available as sawlogs, as stated in the SFM Plan.</p>	Availability of sawlogs for sawmills.	Through environmental compliance inspections and regular on-site visits, CBPPL Operations Superintendents will determine adherence to the 5-Year Plan and the SFM Plan.	CBPPL's annual updates of the SFM Plan, including the volume of sawlogs exchanged with sawmills.
Cutting timber for wharves, stages, and boats	Allow legitimate fisherperson license-holders access to timber for wharves, stages and boats on the DFA.	These values are managed through the 5-Year Planning Process, and CBPPL's Sustainable Forest Management (SFM) Plan. CBPPL issues permits for timber for these values. Management strategies for ensuring the existence of timber for harvesting are found in the 5-Year Operating Plans.	Permits for legitimate fisherperson license-holders to access timber for wharves, stages and boats on the DFA.	CBPPL Operations Superintendents will monitor adherence to the 5-Year Plan and the SFM Plan. The number of permits issued for this purpose on CBPPL limits will be recorded.	Availability of permits for legitimate fisherperson license-holders to access timber for wharves, stages and boats on the DFA.
Firewood to heat homes and businesses	Continue to allow domestic and commercial firewood cutting on the DFA.	This value is managed through the 5-Year Planning Process, and CBPPL's Sustainable Forest Management Plan (SFM Plan).	The existence of domestic and commercial firewood	The number of firewood permits issued on CBPPL limits will be recorded, and	Availability of domestic and commercial firewood cutting permits issued by CBPPL, and response to concerns

ECV	Management Objective	Management Strategy	Indicator and Threshold	Operational Monitoring	Strategic Monitoring (Effectiveness of Management Strategies)
		Firewood permits are made available, and CBPPL enters into agreements for domestic cutting with the government. Management strategies for ensuring the existence of timber for harvesting are found in the 5-Year Operating Plans.	cutting permits issued by CBPPL.	agreements for domestic cutting will be stored.	about availability of firewood.
Indigenous Burial Sites, Sacred and Spirit Areas, and Medicine Plants	Respect known Indigenous Burial Sites, Sacred and Spirit Areas, and Medicine Plants	CBPPL works with the Qalipu Mi'kmaq First Nation Band (QMFN) and Miawpukek First Nations (MFN) to review Annual Operating Plans. Both First Nations participate in the government consultations for 5-Year Operating Plans. Management strategies for historic resources are found in the 5-Year Operating Plans.	No harvesting of timber or road building in known Indigenous Burial Sites, Sacred and Spirit Areas, and Medicine Plants.	CBPPL Operations Superintendents will monitor adherence to the 5-Year and Annual Operating Plans through the #2 & #3 environmental inspections.	The QMFN and MFN will be asked for feedback concerning CBPPL's 5-Year Operating Plans, and provides a copy of the Annual Operating Plans.

6. REFERENCES

- Canadian Heritage Rivers System. 2013. [Website] <http://www.chrs.ca/en/main.php> (Last Accessed June 17, 2015)
- Conservation International. 2013. An international conservation organization. [Website] <http://www.cepf.net/resources/hotspots/North-and-Central-America/Pages/default.aspx> (Last accessed on June 17, 2015)
- BCI, 2015. BCI Species Profiles: *Myotis septentrionalis*. [Website] <http://www.batcon.org/index.php/all-about-bats/species-profiles.html?task=detail&species=2306&country=43&state=57&family=all&limitstart=0> (Last Accessed on June 17, 2015)
- Birdlife International. 2015. A global partnership of conservation organizations. [Website] www.birdlife.org/datazone/index.html. (Last accessed on June 17, 2015)
- Bird Studies Canada. Maintains information on Important Bird Areas in Canada. [Website] <http://www.ibacanada.org/?lang=EN> (Last accessed on January 11, 2013)
- COSEWIC 2006. [COSEWIC assessment and status report on the American eel *Anquilla rostrata* in Canada](#). Committee on the Status of Endangered Wildlife in Canada. Ottawa. X + 71 pp. (www.sararegistry.gc.ca/status/status_e.cfm).
- COSEWIC. 2010. COSEWIC assessment and status report on the Blue Felt Lichen *Degelia plumbea* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. x + 42 pp. (<http://sararegistry.gc.ca/default.asp?lang=En&n=24F7211B-1>).
- Damman, A.W. H. 1983. An ecological subdivision of the island of Newfoundland. Monographiae Biologicae 48:163-206. Biogeography and Ecology of the Island of Newfoundland. Edited by G.R. South, Dr. W. Junk Publishers, The Hague.
- den Otter, M. A. and T. M. Beckley. 2002. "This is Paradise": Community Sustainability Indicators for the Western Newfoundland Model Forest. Information Report M-X-216E. Natural Resources Canada. 53 pp.
- Department of Environmental and Conservation, 2008. Department of Environmental and Conservation Strategic Plan 2008-2011. http://www.env.gov.nl.ca/env/publications/departamental_pubs/strategic_plan_2008_11.pdf
- Department of Environment and Conservation, 2010. Piping Plover (*Charadrius melodus melodus*) http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/piping_plover.pdf. (Last accessed on January 14, 2013)
- Department of Environment and Conservation. 2010. Peregrine Falcon (*Falco peregrinus tundrius/anatum*) http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/perigrine_falcon.pdf (Last accessed on January 14, 2013)
- Department of Environment and Conservation. 2010. Harlequin Duck (*Histrionicus histrionicus*) [Website] http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/harlequin_duck.pdf

(Last accessed on January 14, 2013)

Department of Fisheries and Oceans, 2010. Fish Fact Sheet Atlantic Salmon (*Salmo salar*) [Website] <http://www.dfo-mpo.gc.ca/species-especes/aquatic-aquatique/atl-salmon-saumon-atl-eng.htm> (Last accessed January 14, 2013)

Encyclopaedia Britannica. 2010. [Website] <http://www.britannica.com/Ebchecked/topic/377988/metapopulation> (Last accessed on January 14, 2013).

Environment and Climate Change Canada. 2025. "About Us" [Website] <https://www.gov.nl.ca/ecc/about/> (Last accessed April 4, 2025).

Environment Canada. 2006. [Recovery Strategy for the Red Crossbill, *percna* subspecies \(*Loxia curvirostra percna*\), in Canada](#). Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa. Vii + 29 pp.

Environment Canada. 2007. [Management Plan for the Harlequin Duck \(*Histrionicus histrionicus*\) Eastern Population, in Atlantic Canada and Québec](#). Species at Risk Act Management Plan Series. Environment Canada. Ottawa. vii + 32 pp.

Fisheries and Oceans Canada. 2011. [Management Plan for the Banded Killifish \(*Fundulus diaphanus*\), Newfoundland Population, in Canada \[Final\]](#). Species at Risk Act Management Plan Series. Fisheries and Oceans Canada, Ottawa. v + 23 pp.

Garland, S. and P. Thomas. 2009. [Recovery Plan for Red Knot, *rufa* subspecies \(*Calidris canutus rufa*\), in Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation, Government of Newfoundland and Labrador, Corner Brook, NL. Iv + 12 pp

Goossen, J.P., D.L. Amirault, J. Arndt, R. Bjorge, S. Boates, J. Brazil, S. Brechtel, R. Chiasson, G.N. Corbett, R. Curley, M. Elderkin, S.P. Flemming, W. Harris, L. Heydens, D. Hjertaas, M. Huot, B. Johnson, R. Jones, W. Koonz, P. Laporte, D. McAskill, R.I.G. Morrison, S. Richards, F. Shaffer, C. Stewart, L. Swanson and E. Wiltse. 2002. National Recovery Plan for the Piping Plover (*Charadrius melodius*). National Recovery Plan No. 22. Recovery of Nationally Endangered Wildlife. Ottawa. 47 pp.

Government of Newfoundland & Labrador. 2003. Provincial sustainable forest management strategy. 77 pp. + appendices.

Government of Newfoundland & Labrador. 2014. Provincial sustainable forest management strategy. 64 pp. http://www.nr.gov.nl.ca/nr/pdf/psfms_171114.pdf

Government of Newfoundland & Labrador. 2021. [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#).

Hearn, B. J., D. J. Harrison, A. K. Fuller, C. G. Lundrigan, and W. J. Curran. 2010. Paradigm shifts in the habitat ecology of threatened Newfoundland Martens. Journal of Wildlife Management 74(4): 719-728.

Keeping, B. and C. Hanel. 2006. [A 5 Year \(2006 – 2011\) Management Plan For the Boreal Felt Lichen \(*Erioderma pedicellatum*\) In Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation. Corner Brook, NL. Vii + 44p.

Kotak, B. G., A. Selinger, and B. Johnston. 2005. Influence of watershed features and disturbances quality in Boreal Shield streams and rivers of eastern Manitoba. Manitoba Model Forest Report. 04-2-63. 161p.

- Kunz TH, Reichard JD. 2010. Status review of the little brown Myotis (*Myotis lucifugus*) and determination that immediate listing under the Endangered Species Act is scientifically and legally warranted. Boston University. Available: <http://www.bu.edu/cecb/files/2010/12/Final-Status-Review.pdf> (February 2011).
- PG Lee, Hanneman M, Gysbers JD, Cheng R, Smith W. 2010. Atlas of Canada's Intact Forest Landscapes. Edmonton, Alberta: Global Forest Watch Canada 10th Anniversary Publication #1. 74 pp. [Website] <http://www.globalforestwatch.ca/publications/20100913A> (Last accessed January 14, 2013).
- Lambeck, R.J. (1997): "Focal species: a multi-species umbrella for nature conservation." *Conservation Biology* 11: 849-856
- Liverman, D., M. Batterson, and D. Taylor. Geological disasters in Newfoundland and Labrador. Adapted from Liverman D.G.E., Batterson M.J., Taylor D., and Ryan J. 2001. Geological hazards and disasters in Newfoundland. *Canadian Geotechnical Journal*, 38: 936-956. [Website] <http://www.nr.gov.nl.ca/mines&en/geosurvey/disasters/> (Last accessed January 14, 2013).
- Mahoney, S.P. and J.N. Weir. 2009. *Caribou Data Synthesis*-Progress Report. (Reprint May 2010). Overview of the status of woodland caribou in insular Newfoundland: research methodology, results, interpretations and future projections. Sustainable Development and Strategic Science, Government of Newfoundland and Labrador, St. John's, NL.
- McCarthy, J. W. & G. Weetman. 2006. Age and size structure of gap-dynamic, old-growth boreal forest stands in Newfoundland. *Silva Fennica* [40\(2\)](#): 209–230
- Meades, S. J., S. G. Hay, & L. Brouillet. 2000. Annotated Checklist of the Vascular Plants of Newfoundland and Labrador. [Website] <http://www.digitalnaturalhistory.com/meades.htm> (Last accessed January 14, 2013).
- Megahan, Walter F. and J. Hornbeck, 2000. Lessons Learned in Watershed Management: A Retrospective View. USDA Forest Service Proceedings RMRS-P-13. 2000.
- Model Forest of Newfoundland and Labrador. 2010. [Website] http://www.modelforest.net/index.php?option=com_k2&view=item&layout=item&id=37&Itemid=38&lang=en (Last accessed January 14, 2013).
- Newfoundland Marten Recovery Team. 2010. [Recovery plan for the threatened Newfoundland population of American marten \(*Martes americana atrata*\)](#). Wildlife Division, Department of Environment and Conservation, Government of Newfoundland and Labrador, Corner Brook, Canada. lii + 31 pp.
- Nature Conservancy of Canada. 2010. [Website]. <http://www.natureconservancy.ca/en/where-we-work/newfoundland-and-labrador/our-work/> (Last accessed January 14, 2013).
- NatureServe. 2015. NatureServe Explorer An online encyclopedia of life [Web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at <http://www.natureserve.org/> (Accessed May 20, 2016).
- Olson, D. M. and E. Dinerstein. 2002. The Global 200: Priority Ecoregions for Global Conservation. *Ann. Missouri Bot. Gard.* 89: 199-224.
- Osborne D. R. and J Brazil. 2006. [\[Draft\] Management Plan for the Banded Killifish \(*Fundulus diaphanus*\) in Newfoundland](#). Fisheries and Oceans Canada, and Newfoundland and Labrador Department of Environment and Conservation. 16pp.

Parks and Natural Areas. 2010. Dept. of Environment and Conservation {Website}
<http://www.env.gov.nl.ca/env/parks/index.html> (Last accessed January 14, 2013).

Parks Canada 2010. [Website] <http://www.pc.gc.ca/pn-np/nl/grosmorne/index.aspx> (Last accessed on January 14, 2013).

Rothwell, R.L. 1997. Saskfor MacMillan Limited Partnership Twenty-Year Forest Management Plan and Environmental Impact Statement for the Pasquia-Porcupine Forest Management Area: Forest Hydrology and Watershed Management. 70pp.

Schmelzer, I. (2005) [A management plan for the Short-eared owl \(*Asio flammeus flammeus*\) in Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation. Corner Brook, NL.

Schmelzer, I. (2006) [A management plan for Barrow's Goldeneye \(*Bucephala islandica*; Eastern population\) in Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation. Corner Brook, NL.

SSAC. 2006. The Status of Rattlesnakeroot (*Prenanthus racemosa*) in Newfoundland and Labrador Report #5. [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/Rattlesnake_Root_SSAC.pdf
(Last accessed January 14, 2013)

SSAC. 2006. The Status of Northern Bog Aster (*Symphyotrichum boreale*) in Newfoundland and Labrador Report #6. [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/northern_bog_aster.pdf
(Last accessed January 14, 2013)

SSAC. 2008. The Status of Rock Dwelling Sedge (*Carex petricosa* var. *misandroides*) in Newfoundland and Labrador. Report #10. [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/rock_dwelling_sedge.pdf . (Last accessed January 14, 2013)

SSAC. 2008. The Status of Cutleaf Fleabane (*Erigeron 66omposites*) in Newfoundland and Labrador. Report # 12. [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/cutleaf_fleabane.pdf. (Last accessed January 14, 2013)

SSAC. 2008. [The Status of Tradescant Aster \(*Symphyotrichum tradescantii*\) in Newfoundland and Labrador](#). Report #16.

SSAC. 2008. The Status of Gmelin's Watercrowfoot (*Ranunculus gmelinii*) in Newfoundland and Labrador. Report #17 [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/Gelins_Water_Crowfoot_SSAC.pdf (Last accessed January 14, 2013)

SSAC, 2009. The Status of Vreeland's Striped Coralroot *Corallorhiza striata* var. *vreelandii* in Newfoundland and Labrador. Report #20 [Website]
http://www.env.gov.nl.ca/env/wildlife/endangeredspecies/ssac/Vreelands_Striped_Coralroot_SSAC.pdf
(Last accessed January 16, 2013)

SSAC. 2010. The Status of Gray-cheeked Thrush in Newfoundland and Labrador. Report #24. [Website]
<https://www.gov.nl.ca/ffa/files/wildlife-endangeredspecies-ssac-gray-cheeked-thrush-2010-ssac.pdf> (Last accessed April 15, 2025)

Schmelzer, I. (2005) [A management plan for the Short-eared owl \(*Asio flammeus flammeus*\) in Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation. Corner Brook, NL.

Schmelzer, I. (2006) [A management plan for Barrow's Goldeneye \(*Bucephala islandica*; Eastern population\) in Newfoundland and Labrador](#). Wildlife Division, Department of Environment and Conservation. Corner Brook, NL.

Stenhouse, I.J. (2004) [Canadian management plan for the Ivory Gull \(*Pagophila eburnea*\)](#). Canadian Wildlife Service, St. John's, NL.

The IUCN Red List of Threatened Species. Version 2015-4. www.iucnredlist.org. Downloaded on 02 May 2016.

UNEP-WCMC (Comps.) 2015. The Checklist of CITES Species Website. CITES Secretariat, Geneva, Switzerland. Compiled by UNEP-WCMC Cambridge, UK. Available at: <http://checklist.cites.org>. [Accessed 02/05/2016].

USSR Academy of Sciences. 1988. Red book – Russian Soviet Federal Socialist Republic. 2 volumes. USSR Academy of Science, Moscow, Russia.

Wetlands International. 2010. Delivers the Ramsar Sites information Service. [Website] <http://www.wetlands.org/> (Last accessed January 14, 2013).

Wildlife Division. 2010. [Management Plan for the American Eel \(*Anguilla rostrata*\) in Newfoundland and Labrador](#). Department of Environment and Conservation, Government of Newfoundland and Labrador, Corner Brook. Canada. v + 29 pp.

WWF Global 200. [Website] <http://www.worldwildlife.org/science/ecoregions/global200.html> (Last accessed on January 14, 2013).

7. GLOSSARY³⁷

Aboriginal rights: A practice, custom or tradition integral to the distinctive culture of the aboriginal group claiming the right. Aboriginal rights, including site specific rights may exist, even if specific title does not exist.

Adaptive management: An approach to organizing management so that explicit hypotheses are tested as management activities proceed. A monitoring program tracks outcomes and, depending on how and why actual outcomes differ from expected outcomes, the management approach is reviewed and adjusted.

Affected community: A human community that is affected by the activities on the forest being considered for certification. This will likely include all local communities as well as communities with forest product processing facilities that obtain a high proportion of their furnish from the forest.

Afforestation: The action of converting non-forest land to forest land, which may occur by natural regeneration, seeding, or planting.

Age-class: A distinct group of trees or portion of the growing stock of a forest recognized on the basis of being of similar age.

Biological (bio)diversity: The variability among living organisms from all sources including, *inter alia*, terrestrial, marine and other aquatic ecosystems, and the ecological complexes of which they are a part. This includes diversity within species, between species and of ecosystems (see Convention on Biological Diversity, 1992).

Buffer: A strip or area of vegetation that is left (often unharvested) or managed to reduce the impact of a treatment or action on neighbouring areas.

Canopy closure: The extent to which the upper layer of foliage in a stand or forested area prohibits the passage of sunlight to lower levels, or screens the view of the sky. Also used as an index of competition between adjacent dominant and co-dominant trees.

Clearcut: An area of forest in which all or most of the trees have been harvested; also the harvesting technique that removes all or most of the trees on a site. There is considerable debate within the ecological and forestry communities regarding; how to precisely define a clearcut based on size and configuration of the harvested forest area, proximity to other recently-harvested forest areas, the height or age of the regenerating vegetation both within the harvested area and proximal to the harvested area, etc.

Coarse woody debris: Logs, stumps, and tree limbs on the forest floor in various states of decomposition. Coarse woody debris provides habitat for many wildlife species.

Connectivity: The degree to which different habitat patches or environments are linked by single or multiple corridors of vegetation that provide habitat suitable for dispersal or seasonal movement of particular species, or the migration between ecosystems in response to long-term environmental change. Conditions necessary for connectivity and its effectiveness will depend on the specific purpose of the connectivity and the requirements of species or ecosystems considered.

Conservation attributes: For the purpose of the National Boreal Standard a conservation attribute is an element, structure or process associated with an Exceptional Conservation Value, that can be monitored and managed to ensure its persistence over time. For example, if the ECV designation within a

lk³⁷ From Forest Stewardship Council Working Group. 2004. National Boreal Standard, Version August 6, 2004.

management unit is a consumptive watershed, the associated conservation attributes might include water quality and quantity, flow regimes, integrity of water courses and condition of seeps and springs. These conservation attributes would be identified during the ECV assessment and management strategies to maintain and/or enhance them would be developed, implemented, and monitored, as appropriate.

Conservation biology: Conservation biology is the applied science of maintaining the earth's biological diversity. It integrates and applies the principles of ecology, biogeography, population genetics, economics, sociology, anthropology, philosophy, and other theoretically based disciplines to the maintenance of biodiversity. In the context of protected reserve network planning, applicable concepts from conservation biology include: complete ecosystem representation; protection of core habitats to ensure the maintenance of viable populations of all native species in natural patterns of distribution and abundance; sustaining ecological and evolutionary processes; and ,the maintenance of a landscape that is resilient to environmental change. Many conservation biology practitioners translate these principles into the need for a network of well-distributed protected reserves, combined with adequate buffers and linkage areas to provide for dispersal, seasonal movement, and adaptation to environmental change. The required size and distribution of the reserve network depends on the ecosystems and species present, landscape complexity, and the extent and intensity of human disturbance in the surrounding landscape.

COSEWIC: Committee on the Status of Endangered Wildlife in Canada. The Committee determines the national status of wild Canadian species, subspecies and separate populations suspected of being at risk. COSEWIC bases its decisions on the best up-to-date scientific information and Indigenous Traditional Knowledge available. All native mammals, birds, reptiles, amphibians, fish, molluscs, lepidopterans (butterflies and moths), vascular plants, mosses and lichens are included in its current mandate.

Critical habitat: An ecosystem or particular ecosystem element occupied or used by a species, or local population, that is necessary for their maintenance and/or long-term persistence, and where appropriate, recovery of a species or population.

Customary rights: Rights which result from a long series of habitual or customary actions, constantly repeated, which have, by such repetition and by uninterrupted acquiescence, acquired the force of a law within a geographical or sociological unit

Criterion (pl. Criteria): 1. A means of judging whether or not a Principle (of forest stewardship) has been fulfilled. 2. A distinguishing element or set of conditions by which a forest characteristic or management is judged. 3. A second-order principle that adds meaning and operationality to a principle without itself being a direct measure of performance.

Deforestation - The action of converting forest land to non-forest land. Deforestation implies a permanent conversion of land use; an area of mature forest that is harvested and will be renewed back to forest is not considered to be deforested.

Disturbance: A disruption in the growth and development of an individual, population or community due to natural or anthropogenic factors such as herbivory, forest fires, road building, disease infestation, or tree harvesting.

Disturbance mosaic: The landscape level, spatial pattern of disturbance. The mosaic includes not only areas that have actually been cut, but also inclusions of uncut forest (insular residual), peninsular residual patches, other cuts in close proximity, and forest separating cuts.

Disturbance regime: The characteristic manner in which forests are altered by disturbances. Disturbance regimes are characterized by the nature (e.g., pest, insect, windstorm, etc.), the periodicity, and severity of disturbance events.

Ecodistrict: A part of an ecoregion characterized by a distinctive pattern of relief, geology, geomorphology, vegetation, soils, water and fauna

Ecological integrity: The quality of a natural, unmanaged or managed ecosystem in which the natural ecological processes are sustained, with genetic, species, and ecosystem diversity assured for the future.

Ecoregion: A unit of ecological classification characterized by distinctive ecological responses to climate as expressed by vegetation, soils, water, and fauna.

Ecosite: A unit of ecological classification which is characterized primarily by soil and hydrological conditions.

Ecosystem: A community of all plants and animals and their physical environment, functioning together as an interdependent unit.

Ecosystem diversity: The variety of biomes or habitats occurring within a designated area.

Ecosystem integrity: The diversity of organisms at all levels, including genetic variation, species, populations, ecosystems, landscapes and their physical environments; the ecological patterns, structural attributes, functions and processes that are responsible for that biological diversity and also responsible for ecosystem resilience, allowing for recovery following disturbance.

Ecosystem representation: Inclusion within a reserve network of the full spectrum of biological and environmental variation, including genotypes, species, ecosystems, habitats, and landscapes.

Economic viability: The capability of an entity to be economically self-sustaining. In the long term, this means that the entity must at least break even and, more likely be profitable. In the short term, entities can run at a loss depending on their access to financial backing and the value of cash and assets held.

Enduring feature: A landscape element or unit within a natural region characterized by relatively uniform origin of surficial material, texture of surficial material, and topography.

Ephemeral stream: A stream that flows briefly only in direct response to precipitation in the immediate locality and whose channel is at all times above the water table.

Exotic species: An introduced species not native or endemic to the area in question.

Focal species: Focal species builds on the concept of umbrella species, whose habitat requirements are believed to encapsulate the needs of other species (Lambeck 1997). The focal species approach assumes that meeting the requirements of the most demanding species will result in a landscape design encompassing the needs of a wider range of species.

Forest: 1. A plant community dominated by trees and other woody vegetation, growing more or less closely together. 2. An area managed for the production of timber and other forest products or maintained under woody vegetation for such indirect benefits as protection of site or recreation. 3. An aggregate of stands.

Forest conversion: The substantial or severe modifications to the structure and dynamics of a forest, as a result of management activities, resulting in a significant reduction in the complexity of the forest system; or the transformation of a forest into a permanently non-forested area.

Forest management activities: Any or all of the operations, processes or procedures associated with managing a forest, including, but not limited to: planning, consultation, harvesting, access construction and maintenance, silvicultural activities (i.e., planting, site preparation, tending), monitoring, assessment, and reporting.

Forest unit: An aggregation of forest stands for management purposes which have similar species composition, develop in a similar manner (both naturally and in response to silvicultural treatments), and are managed under the same silvicultural system.

Forests with Exceptional Conservation Value (FECV): Critically imperiled (G1) and imperiled (G2) species and ecological communities.

Critically imperiled: A plant or animal or community, often referred to as G1, that is globally extremely rare or, because of some factor(s), especially vulnerable to extinction. Typically five or fewer occurrences or populations remain, or very few individuals ($\leq 1,000$), acres ($\leq 2,000$ acres or 809 hectares), or linear miles (≤ 10 miles or 16 kilometers) exist.

Imperiled: A plant or animal or community, often referred to as G2, that is globally extremely rare or, because of some factor(s), especially vulnerable to extinction or elimination. Typically six to 20 occurrences, or very few remaining individuals (1,000 to 3,000), or acres (2,000 to 10,000 acres or 809 to 4,047 hectares), or linear miles (10 to 50 miles or 16 to 80.5 kilometers) exist.

Fragile ecosystems: Ecosystems (at any scale) which are prone to disruption from even modest management interventions or natural disturbance events.

Gap analysis: An assessment of the protection status of biodiversity in a specified region, which looks for gaps in the representation of species or ecosystems in protected areas.

Genetic diversity: Variety within individuals within a species or a population, or more specifically the variety of DNA or alleles within a species or population.

Habitat: 1. Those parts of the environment (aquatic, terrestrial, and atmospheric) often typified by a dominant plant form or physical characteristic, on which an organism depends, directly or indirectly, in order to carry out its life processes. 2. The specific environmental conditions in which organisms thrive in the wild.

Home range: The area over which an animal roams during the course of its usual wanderings and spends most of its time. When home ranges are marked and defended they are referred to as territories. In vertebrates, the size of an animal's home range is roughly proportional to its body size.

Indicator: A specific requirement in the certification standards, subordinate to the Performance Measures and Objectives.

Indigenous: In this standard, the term "Indigenous" will be understood to be inclusive of those groups constitutionally-recognized as being Aboriginal People, including Indian, Métis and Inuit.

Intact: The maintenance of (i.e., no readily determinable changes to) ecological functions at a forest or landscape scale.

Landscape: A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area.

Landscape level: At a spatial scale above a single plant community or forest stand and below a region (See also definition of Landscape).

Late seral stage: A late stage in succession (the process of community development after disturbance) where the forest canopy starts to open up, and the amount of vertical and horizontal structural diversity increases. The time since disturbance at which a late seral stage could be said to exist varies from forest unit to forest unit.

Local community: Any (human) community that is on or adjacent to the forest that is being audited for certification. If no communities meet this criterion, then the scope of "local" should be expanded to cover communities within a reasonable daily commuting distance from the boundary of the forest being certified.

Local People: are considered local where they permanently reside within daily commuting distance by car or boat from the management unit, or where they are part of the Indigenous people whose lands and territories contain or are contained within the management unit.

Long term: The length of time consistent with sustainability. Ideally, this is the time-scale of the forest owner or manager as manifested by the objectives of the management plan, the rate of harvesting, and the commitment to maintain permanent forest cover. The length of time involved will vary according to the context and ecological conditions, and will be a function of how long it takes a given ecosystem to recover its natural structure and composition following harvesting or disturbance, or to produce mature or primary conditions.

Merchantable: A log or tree which meets or exceeds minimum size requirements and contains a proportion of sound wood in excess of minimum requirements, as determined according to applicable scaling (wood measurement) standards.

Native species: A species that occurs naturally in the region; endemic to the area.

Natural cycle: Nutrient and mineral cycling as a result of interactions between soils, water, plants, and animals in forest environments that affect the ecological productivity of a given site.

Natural forest: A forest area where many of the principal characteristics and key elements of native ecosystems such as complexity, structure and diversity are present.

Non-forested land: Land that is classed as being used for a purpose other than supporting forest growth, such as agriculture, roads, trails, landings, gravel pits, and camps. Deforestation is the process of converting forest land to non-forest land; afforestation is the reverse process.

Non-timber forest products: All forest products except timber, including other materials obtained from trees, such as resins and leaves, as well as any other plant and animal products produced by the forest. In the boreal forests of Canada, there are many commercial enterprises based on non-timber forest products, such as hunting and fishing lodges, trapping operations, outfitting, remote tourist operations, and youth camps.

Old forest: Later stage(s) in forest development which may be distinctive in composition but are always distinctive in structure from earlier (young and mature) successional stages.

Optimal or highest and best value: In many cases, a log or tree can be used for a variety of purposes, with the sale price of the wood dependent on the use to which it will be put. Optimal (or highest and best) value is obtained when the highest price is obtained, or the wood is used for the purpose which best meets financial or socio-economic objectives.

Peer review: An independent or external review by experts on the subject being considered.

Precautionary approach: An approach that tends to refrain from actions where the outcome is not known. In a forest management context it refers to situations in which a forest manager will often be required to act with incomplete knowledge of cause and effect relationships, and therefore a precautionary approach includes the following:

- The manager avoids actions that may lead to irreversible changes to ecosystem function and resilience;
- Alternative management strategies are developed and evaluated, including the alternative of no management intervention, to identify alternatives that are least likely to impair the viability of the species or ecosystem;
- The onus is on the manager to demonstrate that proposed management activities are not likely to impair ecosystem function and resilience;
- When previously unanticipated threats to ecosystem integrity are identified or knowledge of ecosystem processes increases, the manager takes timely, efficient and effective corrective actions; and,
- The manager remains mindful of the needs of future generations.

Pre-industrial forest: 1. A native forest which has not been subjected to large scale harvesting or other forms of human management. 2. A forest area such as existed prior to human settlement in the region occupied by the forest.

Protected area: An area protected by legislation, regulation, or land-use policy to control human occupancy or activity. Protection can be of many different forms. The International Union for the Conservation of Nature (IUCN) identified six main categories of protected areas.

Protected area network: The total network of places and locations protected by various means within a forest or an area, including riparian reserves, habitat reserves, parks, and all other protected areas.

Principle: An essential rule or element.

Public Advisory Group (PAG): A committee with a diversity of interests that represents the public's views during forest management planning and implementation.

Public participation process: A formal process of public involvement. A public participation process ordinarily involves a defined membership, established ground rules, opportunities for interaction among participants and the provision for ongoing involvement. It may involve establishing a new process, building on an existing process or reviving and adapting a previously existing process. A public participation process is recommended on all forest lands, and is required on Crown lands.

Remote: Areas without motorized access because roads to the area are either non-existent, seasonal, closed, abandoned or re-vegetated.

Riparian area: 1. The area related to the bank or shore of a water body. 2. The area of forest having qualities influenced by proximity to a water body.

Sensitive sites: Sites with soils prone to erosion and/or nutrient loss as a result of normal management activities or natural disturbances. Sensitivity may be linked to human activity, disruption of water flow, alteration of stand structure or composition, or some other factor. For conducting forest operations, sensitive sites often include areas with steep slopes, shallow soils, or easily rutted soils.

Silviculture: The technique of producing and tending a forest by manipulating its establishment, composition and growth to best fulfill the objectives of the owner. This may, or may not, include timber production.

Sites of special cultural, ecological, economic, or religious significance: Include, but are not limited to, sites relating to or associated with the following:

- Ceremonial/Spiritual/Religious (e.g., vision/spirit quest area, repository for the dead, gathering place, sacred places);
- Traditional Oral History (e.g., origin of a story, legend);
- Cultural Landforms (e.g., named places, marker sites, legendary landforms);
- Supernatural Beings (e.g., supernatural areas);
- Transportation (e.g., grease trail, trading route, water route, portage area);
- Habitation (e.g., permanent village, seasonal residence, storage area);
- Recreational (e.g., gathering place, games or competition place);
- Cross-Cultural Interaction (e.g., first contact, trade with Europeans, or other Indigenous People(s); and,
- Education and Training (e.g., where traditional skills, values or knowledge are conveyed).

Species at Risk: Although this term is also used by COSEWIC, in this standard it is used in a more generic sense to refer to all species about which concern exists regarding their viability at regional, provincial, or a national scale and/or which were formerly referred to as rare, threatened or endangered.

Species diversity: The variety of different organisms at the species taxonomic level.

Stakeholder: An individual or organization with an interest in the state and/or management of a forest as a result of economic, social, spiritual or conservation-oriented ties to the forest.

Stand: A community of trees possessing sufficient uniformity in composition, constitution, age, arrangement or condition to be distinguishable from adjacent communities.

Steep slopes: Slopes with an incline such that normal forest operations would result, or would have the potential to result in moderate or severe erosion.

Structural diversity: The diversity of forest structure, both vertical and horizontal, that provides for a variety of forest habitats for plants and animals. The variety results from layering or tiering of the canopy and die-back, death, and ultimate decay of trees. In aquatic habitats, structural diversity results from the presence of a variety of structural features such as logs and boulders, that create a variety of habitats.

Structure: 1. The various horizontal and vertical physical elements of the forest. 2. In landscape ecology, the spatial inter-relationships between ecosystems including energy fluxes, distribution of materials and species relative to the sizes, shapes, numbers, kinds and configurations of the ecosystems. 3. The distribution of trees in a stand or group by age, size or crown classes (e.g., all even-aged, uneven-aged, regular, and irregular structures).

Subspecies: A taxonomic designation below the level of species. For some species there is considerable uncertainty between the distinctions between species, subspecies, genus and populations.

Succession: Progressive changes in the species composition and structure of a forest community caused by non-catastrophic natural processes (nonhuman) over time.

Surrounding lands: Lands which abut the management unit.

Tenure: Socially defined agreements held by individuals or groups, recognized by legal statutes or customary practice, regarding the "bundle of rights and duties" of ownership, holding, access and/or usage of a particular land unit or the associated resources there within (such as individual trees, plant species, water, minerals, etc).

Traditional Ecological Knowledge (TEK): An accumulated body of knowledge that is rooted in the spiritual health, culture, and experiences of those who are close to the lands. It is based on an intimate knowledge of the land, its physiographic and natural features, climate, and wildlife, and the relationships between all aspects of the environment. Although in many uses it refers to knowledge of Indigenous peoples, others with intimate knowledge and experience of the land also have developed traditional ecological knowledge.

Traditional use: The use of land or the pursuit of activities on a forest.

Tree: A tree is considered to be a woody perennial plant that grows to a height of at least 4.5m.

Unique ecosystems: Rare or uncommon ecosystems of any scale within the management unit or forest being considered for certification. For example, these may include disjunct ecological communities, breeding grounds of uncommon species, etc.

Use rights: Rights for the use of forest resources that can be defined by local custom, mutual agreements, or prescribed by other entities holding access rights. These rights may restrict the use of particular resources to specific levels of consumption or particular harvesting techniques.

Utilization: The amount of potentially merchantable timber on a site that is actually used.

Verifier: Data or information which provides specific details or measures which enhance the ease, specificity or precision of assessment of an indicator. In this Standard the Verifiers: noted are not mandatory means by which to assess indicators, but suggested or useful means.

Vertical structure: The amount and orientation of above-ground biomass in a stand or forest area.

Watershed: An area of land through which water drains into other streams or waterways via underground or surface streams and rivers.

Wetland: Lands transitional between terrestrial and aquatic systems where the water table is at or near the surface, or the land is covered by shallow water at some time during the growing season. Wetlands are characterized by poorly drained soils and predominantly hydrophilic or water tolerant vegetation.

Wildlife: Any species of amphibian, bird, fish, mammal and reptile found in the wild, living unrestrained or free roaming and not domesticated.

APPENDIX 1 CANDIDATE ECVS (FORMERLY HCVS) ON CBPPL'S DFA

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
3. Does the forest include critical habitat containing globally, nationally or regionally <u>significant seasonal concentration of species</u> (one or several species, e.g., concentrations of wildlife in breeding sites, wintering sites, migration sites, migration routes or corridors – latitudinal as well as altitudinal)?							
Middle Ridge Wildlife Reserve	Upland plateau of extensive barrens, kalmia heaths, bogs, and fens which provides calving and wintering grounds for Woodland Caribou, and breeding grounds for a number of waterfowl species.	Adjacent; shares ~7km boundary	Protected Important Bird Area (IBA)	Negligible		Reserve is very large and only 6% of it borders on the DFA	ECV
Codroy Valley	Balsam fir forest mixed with yellow birch and mountain maple thickets; contains red crossbill and ovenbird	Separates Codroy Valley Estuary IBA from DFA	Important Bird Area (IBA)	No impact from forest operations on the DFA		At 8km from the DFA it is not close enough to be impacted.	Not ECV
Gros Morne National Park	Mountainous barrens, forest, and diverse coastal lowlands; provides calving and wintering grounds for Woodland Caribou.	Borders DFA	Protected Important Bird Area (IBA)	Remaining ecologically connected to its broader landscape	Protected		ECV
Codroy Valley Estuary	Important breeding and staging site for numerous waterfowl species	~ 7 km west of DFA	Important Bird Area (IBA)	No impact from forest operations on the DFA	Closed to hunting	At 7km from the DFA it is not close enough to be impacted.	Not ECV
Terra Nova National Park	Forested with black spruce, balsam fir, white birch, and tamarack; also features numerous lakes and upland plateau fens and bogs.	~ 27 km southeast of DFA	Protected	No impact from forest operations on the DFA	Protected	At 27km from the DFA it is not close enough to be impacted.	Not ECV

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Cook's Marsh	Marsh valuable for waterfowl	Part of DFA just west of Corner Brook	N/A	Disturbance of habitat and nesting birds	Conservation agreements between DUC and CBPPL include wetland and associated upland areas.		ECV
Upper Humber Wetlands Complex (Birchy Basin)	Reportedly has the highest known inland waterfowl breeding densities on the island of Newfoundland; important staging area for black duck	Inside the DFA in District 16	N/A	Disturbance of habitat and nesting birds	No harvesting in Birchy Lake Management Unit, Gales Bottom/Alder Brook Management Unit, and Neds Steady Management Unit as defined by GovNL. These areas are within the Wetlands Complex.		ECV
Long Island and The Narrows Properties	Important breeding habitat for dabbling ducks and Canada Geese	Adjacent to CBPPL managed lands in the Northwest Gander River area	N/A	Disturbance of habitat and nesting birds		No evidence of significant seasonal concentrations of species; can be dealt with through 5-Yr. Planning process	Not ECV
Woodland Caribou <i>Rangifer tarandus</i>	Caribou prefer mostly barren land during the summer months, moving to areas of mixed forests during the colder months.	Eight herds on DFA: Gros Morne, Aides Pond, Hampden Downs, Gaff	Populations in decline although not yet classified as at risk.	Can disturb calving areas, wintering areas, and travel corridors	Restrictions on harvesting in "core" areas and buffer areas, and during calving ³⁸	Provincial government is revising guidelines and the Canadian Boreal Forest	Habitat for Woodland Caribou ECV

³⁸ Forest Management Guidelines for Woodland Caribou (*Rangifer tarandus caribou*) for the Island of Newfoundland, 2007

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
		Topsails, Buchans, Mount Peyton, Pot Hill, and Middle Ridge				Agreement is working on guidelines across Canada	
Atlantic Salmon <i>Salmo salar</i>	Spawn in freshwater streams, in gravel-bottom rapid areas above or below a pool	Throughout the DFA	Not at risk	Spawning areas could be impacted by harvesting	Minimum 30m buffers on streams, wider on spawning areas ^{39,40}		Habitat for Atlantic Salmon ECV
4. Does the forest contain critical habitat for regionally significant species (e.g., species representative of habitat types naturally occurring in the management unit, focal species, or species declining regionally)?							
See Table 4 in Assessment Report							
5. Does the forest support concentrations of species at the edge of their natural ranges or outlier populations?							
Red Pine <i>Pinus resinosa</i>	Deep sands or gravel soils	Sandy Lake, Howley, Birchy Narrows, Birchy Lake, West Brook	Endangered	Harvesting not permitted in red pine stands	No permits issued for red pine		ECV
White Pine <i>Pinus strobus</i>	Moist sandy soil	Particularly in Grand Lake South and Southwest Gander areas	Rare	Risk from scarring during road construction and harvesting operations	White pine protected		ECV
Yellow Birch <i>Betula alleghaniensis</i>	Variety of sites, often in association with balsam fir and other hardwoods	Bottom Brook	Rare	Access to yellow birch for firewood is increased when roads are built to harvest operations	None		ECV
Black Ash <i>Fraxinus nigra</i>	Wet sites along rivers or the margins of swamps	South of Bonne Bay; Bottom Brook	Rare	Negligible	None		ECV

³⁹ [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#), 2021

⁴⁰ Policy for the Management of Fish Habitat, 1986

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Balsam Poplar <i>Populus balsamifera</i>	Found in association with white birch, balsam fir, and various willows and alders	Crabbes River (TCH) Lomond	Rare	Could possibly be cut for hog fuel	None	Insufficient information to assess	Possible ECV
Arctic Hare <i>Lepus arcticus</i>	Primarily in upland alpine forests, at elevations >1500m and in rugged topography	Possibly on DFA but not in commercial forest	Rare	None as do not occur in commercial forest	Protected only in Gros Morne National Park	Determined not ECV as there is no threat from harvesting	Not ECV
6. Does the forest lie within, or adjacent to, or contain a conservation area: a) designated by an international authority, b) legally designated or proposed by relevant federal/provincial/ territorial legislative body, or, c) identified in regional land use plans or conservation plans?							
Gros Morne National Park	Variety of habitats	Borders DFA	UNESCO World Heritage Site	Connectivity to broader landscape, Main River in particular			FECV
Grand Codroy Estuary (Codroy Valley Estuary IBA)	Wetlands; important breeding and staging site for numerous waterfowl species	~7 km from the DFA	RAMSAR site	Forestry not listed as a threat	Closed to hunting	Well outside of the DFA	Not FECV
Serpentine Lakes	Forest types of the region, subarctic and serpentine barrens habitats	Shares a 12 km boundary with DFA on the southeast corner	IBP Site				FECV
Crooked Bog	Ombrotrophic bogs, oceanic raised bogs, fens and black spruce forest	Completely surrounded by DFA	IBP Site				FECV
Brownmore Bog	Western blanket bogs	Completely surrounded by DFA	IBP Site				FECV
Sandy Lake	Red pine forests	Completely surrounded by DFA and is bounded by the lake	IBP Site	Harvesting not permitted in red pine stands			FECV

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Grand Lake Brook	Rich fen indicator species	Completely surrounded by DFA	IBP Site				FECV
Main River (Waterway Provincial Park and Special Management Area)	Watershed	Lies ~1km northeast of a small outlier parcel of DFA, and the southeastern tip is within ~2 km DFA	Canadian Heritage River Provincial Park/Special Management Area	None	Protected	DFA not adjacent to watershed and there is already a 49-km ² special management area buffering the park	Not ECV
Sandy Point (NCC)	Important beach system for providing nesting sites for Piping Plover, Common Tern and Arctic Tern. It is the only breeding site in the Province for the Willet and is a roosting site for the Caspian Tern. Largest spartina salt marsh and one of the largest eel-grass beds in the province	Lies ~12 km north of the DFA	1,000 ha covering nearly all of Sandy Point is zoned as "Conservation" under the St. George's Municipal Plan.	None from CBPPL operations	No development is permitted that will exploit the natural resources of the area.		Not FECV
The Grassy Place (NCC)	Wide grassy fluvial wetland, forested slopes, and sparsely vegetated mountaintops.	Lies ~10 km east of the DFA	Owned fee simple by NCC	None from CBPPL operations	NCC permits only low-impact historical uses such as hiking, hunting, and berry picking. Cutting and ATV use are prohibited.		Not FECV
Lloyd's River (NCC)	Mature forest, fen, marsh and stream habitats	Lies ~48 km east of the DFA	Owned fee simple by NCC	None from CBPPL operations	NCC permits only low-impact historical uses such as hiking, hunting, and berry picking. Cutting		Not FECV

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
					and ATV use are prohibited.		
West Brook Ecological Reserve	Natural stands of red pine	Completely surrounded by DFA	Protected	None	Protected		ECV
Little Grand Lake Provisional Ecological Reserve	Protects extensive bogs and barrens, as well as mature boreal forest	Borders the DFA in two areas, total of ~ 70km	Protected	None	Protected		ECV
Little Grand Lake Wildlife Reserve		Shares ~ 7km of boundary with DFA	Protected	None	Protected		ECV
Middle Ridge Wildlife Reserve	Upland plateau of extensive barrens, kalmia heaths, bogs, and fens	Shares ~ 7 km of boundary with DFA	Protected	None	Protected		ECV
Flatwater Pond Provincial Park Reserve	Similar to surrounding forest	Completely surrounded by DFA	Protected	None	Protected		ECV
Jonathan's Pond Provincial Park Reserve	Preserves a majestic stand of white birch	Almost completely surrounded by DFA	Protected	None	Protected		ECV
Barachois Pond Provincial Park	Balsam fir forest with an understory of ferns and moss; some white pine	Shares ~3 km of boundary with the DFA	Protected	None	Protected		ECV
Sir Richard Squires Memorial Provincial Park	Larch, birch, fir and spruce	Completely surrounded by DFA	Protected	None	Protected		ECV
T'Railway Provincial Park	Many of the Island's representative natural and scenic landscapes	Runs through the DFA	Protected	None	Protected		ECV
7. Does the forest constitute or form part of a globally-, nationally-, or regionally-significant forest landscape that includes populations of most native species and sufficient habitat such that there is a high likelihood of long-term species persistence?							
No large landscape-level forests.							No ECV
8. Does the forest contain naturally rare ecosystem types?							
Nothing identified by international agencies: Conservation International, WWF Ecoregion Conservation Assessments, or NatureServe. Atlantic Canada Conservation Data Centre doesn't yet have info on an ecosystem or community basis.							No ECV
9. Are there ecosystem types within the forest or ecoregion that have significantly declined?							

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Sandy Lake Red Pine	Red pine	Within DFA	Rare	No permits are issued for red pine	No permits are issued for red pine		ECV
Grand Lake South White Pine	White pine, balsam fir, spruce	Within DFA	Rare	White pine protected	None	Area will never be harvested due to terrain and soil moisture	ECV
10. Are large landscape level forests (i.e., large, unfragmented forests) rare or absent in the forest or ecoregion?							
Cat Arm							FECV
Hampden Downs							FECV
Little Codroy							FECV
11. Are there nationally/regionally significant diverse or unique forest ecosystems?							
Serpentine areas: rare plant species located in openings where serpentine soil is exposed	Lightly forested, dominated by larch and white pine or by black spruce	North Arm Hills, Northwest Gander River near bridge on Bay D'Espoir Hwy	Rare plant species	Risk from construction of forest access roads	None		ECV
Limestone Areas: rare plant species not found on limestone barrens	Steep cliffs and the talus slopes underneath – not forested	South shore of Serpentine Lake; Goose Arm Area	Rare	Risk from construction of forest access roads	None		ECV
Red pine areas	Already covered in questions 1, 5, 6 & 9						
Riparian ecosystems	Significant areas would not be forested and would be within the buffers on the scheduled salmon rivers	Most significant areas: Upper Humber, Bottom Brook near the Burgeo turnoff, Harry's River, and Hughes Brook estuary		Riparian ecosystems could be compromised by forest operations with no restrictions	Minimum 30m buffers are required on all streams identified on 1:50,000 topographic maps and those greater than 2m wide ⁴¹		ECV
12. Does the forest provide a significant source of drinking water?							

⁴¹ [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#), 2021

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Thirty-six Protected Public Drinking Water Supplies servicing 32 communities		Within DFA	All legally protected except small portions of three	Negligible with management guidelines	Guidelines for Forestry Operations Within Protected Public Water Supply Areas; Municipal Watershed Monitoring Committees		ECV
13. Are there forests that provide a significant ecological service in mediating flooding and/or drought, controlling stream flow regulation, and water quality?							
Black Duck and Appleton are flood risk zones within DFA						No information to indicate that the harvested areas in these watersheds at any one time are large enough to have an effect on these flood zones	Not ECV
Stephenville, Steady Brook and Deer Lake are flood risk zones downstream of DFA						As above	Not ECV
14. Are there forests critical to erosion control?							
Slopes greater than 25 degrees (46%)		Throughout the DFA		Depending on the underlying material, mass movement of earth could occur	Generally slopes greater than 40% are not harvested. Infrequently, very small areas of slopes up to 45% are harvested.		ECV
15. Are there forest landscapes (or regional landscapes) that have a critical impact on agriculture or fisheries?							

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
Freshwater Fishes	Freshwater streams	Throughout the DFA	Banded Killifish are vulnerable		Minimum 30m buffers on streams, wider on spawning areas. A 30m no-grub zone is maintained around any waterbody ^{42,43}		ECV
Atlantic Salmon <i>Salmo salar</i>	Spawn in freshwater streams, in gravel-bottom rapid areas above or below a pool	Throughout the DFA	Not at risk	Spawning areas could be impacted by harvesting	Minimum 30m buffers on streams, wider on spawning areas. A 30m no-grub zone is maintained around any waterbod ^{34,35}		ECV
16. Are there local communities? Is anyone within the community making use of the forest for basic needs/ livelihoods?							
Recreation	Various habitats	Throughout the DFA		Harvesting could impact some aspects of recreation, i.e. viewscapes		Forest operations roads create access	ECV
Income from working in the forest		Throughout the DFA		None		Dependent on forest operations	ECV
Cutting timber for houses, wharves, stages, and boats		Throughout the DFA		None		Forest operations roads create access	All ECV except timber for houses
Firewood to heat homes and businesses		Throughout the DFA		None	Where markets exist, non-commercial tree species that are harvested should	Forest operations roads create access	ECV

⁴² [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#), 2021

⁴³ Policy for the Management of Fish Habitat, 1986

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
					be brought to roadside ⁴⁴		
Company roads to gain access to the wilderness for hunting, fishing, berry picking or tree species that are of no value for the paper mills.		Throughout the DFA		None		Forest operations roads create access	ECV
Cabins for pleasure, relaxation, and hunting		Throughout the DFA		Harvesting could impact viewscape	A minimum 100-metre, no-cut buffer is to be left between operations within approved cabin development areas. ⁴⁵	Forest operations roads create access	ECV
Sawlogs for local sawmills		Throughout the DFA		If not separated for sawmills, sawlogs could be used for pulpwood	CBPPL has Saw log exchange agreements to supply integrated sawmills with sawlogs		ECV
17. Is the traditional cultural identity of the local community particularly tied to a specific forest area?							
Burial Sites, Sacred Areas, Spirit Areas and Medicine Plants	Mostly along rivers and near ponds	In FMDs 6, 14, & 15	Generally not Protected	Unsure	The Historic Resources Division reviews Five-year Operating Plans to determine the location of historic resources; appropriate mitigation measures are then designed.	Specific areas can be dealt with through the 5 Year Planning process	ECV

⁴⁴ [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#), 2021

⁴⁵ [Environmental Protection Guidelines for Forestry Operations in Newfoundland and Labrador](#), 2021

Candidate ECV	Habitat	DFA Occurrence	Status Info	Risk From Forest Operations	Current Management	Comments	Decision
					Annual Operating Plans are provided to Qalipu Mi'kmaq First Nation for review and feedback. Annual Operating Plans in District 6 are provided to Miawpukek First Nation for review and feedback.		
18. Is there a significant overlap of values (ecological and/or cultural) that individually did not meet ECV thresholds, but collectively constitute ECVs?							
There was no apparent overlap of values that would lead to new ECV'S, and as most of the submitted values in this question were identified as ECVs, further analysis was unnecessary.							No ECV

APPENDIX 2 INITIAL MEETINGS WITH STAKEHOLDERS/INTERESTED PARTIES FOR HCVF REPORT

Stakeholder	Meeting Date	
	Introduction to HCVF	Submission of HCVFs
Western Environment Centre Simon Jansen	December 15, 2010	N/A
Grenfell Campus, MUN, Environmental Policy Unit Dr. Erin Kelly	December 15, 2010	N/A
Salmon Preservation Association for the Waters of Newfoundland (SPAWN) John McCarthy	January 11, 2011	N/A
Atlantic Salmon Federation Don Ivany	January 11, 2011	April 26, 2011
Humber Natural History Society Lois Bateman	January 14, 2011	N/A
International Appalachian Trails NL Paul Wylezol, Kevin Noseworthy	January 14, 2011	N/A
Model Forest of Newfoundland and Labrador Sean Dolter	January 14, 2011	N/A
Humber Valley Regional Planning Advisory Authority Don Downer	January 14, 2011	N/A
CBPPL Public Advisory Committee (See attendance list, meeting # 38)	January 19, 2011	N/A
Ducks Unlimited Canada Danielle Fequet	January 19, 2011	March 22, 2011
NL Environment Network Chris Hogan	January 25, 2011	April 12, 2011
Canadian Parks and Wilderness Society/Coalition for Sustainable Forestry Dr. Ian Goudie	January 25, 2011	April 12, 2011
Sierra Club of Canada, Atlantic Canada Chapter Nicole Renaud	January 25, 2011	N/A
Protected Areas Association Ruth French, Doyle Wells, Valerie Simmons, Fanta Yao	January 25, 2011	N/A
Department of Natural Resources, St. John's Bill Clark, Rod Hillyard	January 25, 2011	N/A
Wildland Associates Bruce Roberts	January 25, 2011	N/A
Nature Conservancy of Canada Doug Ballam, Randal Greene, Daniel Myers	January 26, 2011	April 11, 2011
Salmonid Council of Newfoundland and Labrador Trevor Davis	January 27, 2011	N/A
Salmon Association of Eastern NL Dr. Larry Felt	January 27, 2011	N/A
Natural History Society of Newfoundland and Labrador Dr. Rita Anderson, Dr. Allan Stein, Dr. John Jacobs	January 27, 2011	April 11, 2011
Department of Natural Resources, St. Georges Jamie Kennedy, Dick Brake, Craig Stoyles, Mike Bennett, Paul Grenier	February 15, 2011	N/A
CBPPL Woodlands Staff	February 21, 2011	N/A
College of the North Atlantic First Year Forest Resource Technician Students	February 22, 2011	N/A

Newfoundland and Labrador Snowmobile Federation Donnie O'Keefe, Bruce Nichols	March 3, 2011	
Western Sno-Riders John Alteen, Joe Callaghan	March 3, 2011	
Newfoundland & Labrador Outfitters Association Keith Payne	March 11, 2011	
Department of Natural Resources Keith Deering, Ivan Downton, Dave Cheeks, Eric Young, Dave Poole, Basil English	March 22, 2011	
Hospitality Newfoundland and Labrador Todd Wight	March 29, 2011	
Federation of Newfoundland Indians Brendan Sheppard, Roger Gallant	March 31, 2011	Sent their values - no second meeting
Town of Deer Lake Maxine Hayden (Mgr.), Councilors	May 9, 2011	
Gros Morne National Park Peter Deering, Tom Knight	May 24, 2011	

APPENDIX 3 CANDIDATE EXCEPTIONAL CONSERVATION VALUES (FORMERLY HCVS) SUBMITTED BY STAKEHOLDERS

Stakeholder	Value(s)	Corresponds to Framework Question	Decision
Ducks Unlimited Canada	Cook's Marsh	3. Significant seasonal concentration of species	ECV
	Birchy Basin		ECV
	Long Island Property		Not ECV
	The Narrows Property		Not ECV
Nature Newfoundland and Labrador (Formerly Natural History Society of Newfoundland and Labrador)	Landscape connectivity	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Forest self-regeneration	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Biodiversity	3. Significant seasonal concentration of species 6. Conservation Areas	ECV through GMNP and all Provincial Parks and Reserves on DFA
	Ecological representation	6. Conservation Areas	ECV As above
	Ecological persistence	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Arboreal lichens	1. Species at Risk or potential habitat of Species at Risk	ECV
	Newfoundland marten	1. Species at Risk or potential habitat of Species at Risk	ECV
	Caribou	1. Species at Risk or potential habitat of Species at Risk 3. Significant seasonal concentration of species	ECV
	Salmonids	16. Critical impact on agriculture or fisheries	ECV
	Drinking Water	12. Significant source of drinking water	ECV
	Recreation	17. Fundamental to meeting basic needs of local communities	ECV
	Carbon	Does not fit into any of the questions, rather it is dealt with through Criteria 6.1, 6.3, 6.5, and 8.3 of the Boreal Standard; also through Appendix 3 of the Standard: International	N/A

		Agreements Ratified by Canada, Framework Convention on Climate Change.	
Nature Conservancy of Canada	Representative Forests	6. Conservation Areas	Specific conservation areas are designated ECV
	<u>NCC Focal Areas:</u> Sandy Point Lloyd's River Grassy Place Grand Codroy River Estuary	6. Conservation Areas	Not ECV Not ECV Not ECV Not ECV
	Newfoundland Marten	1. Species at Risk or potential habitat of Species at Risk	ECV
	Riparian and Wetland Areas	11. Significant diverse or unique forest ecosystems	Riparian Ecosystems ECV
	Limestone and Serpentine Barrens	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Piping Plover	1. Species at Risk or potential habitat of Species at Risk	ECV No management required
	Old Growth	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Caribou	1. Species at Risk or potential habitat of Species at Risk 3. Significant seasonal concentration of species	ECV
	Arboreal Lichens	1. Species at Risk or potential habitat of Species at Risk	Boreal Felt Lichen ECV Other arboreal lichens Possible ECV
Canadian Parks and Wilderness Society/ Coalition for Sustainable Forestry/ NL Environment Network	Old growth forest	7. Significant large landscape level forest, such that there is a high likelihood of long-term species persistence	ECV through large landscape level forest (being identified)
	Increased buffers Around parks/protected areas	6. Conservation Areas	Specific conservation areas are designated ECV
	Riparian areas	11. Significant diverse or unique forest ecosystems	ECV
	Caribou habitat	3. Significant seasonal concentration of species	ECV

Atlantic Salmon Federation	Protection of critical salmon spawning habitat	3. Significant seasonal concentration of species 16. Critical impact on agriculture or fisheries	ECV
Federation of Newfoundland Indians	Burial areas Sacred areas Spirit areas Medicine Plants	18. Traditional cultural identity of local community	ECV
Lew Hounsell	Corner Brook Watershed (drinking water supply)	12. Significant source of drinking water	ECV
Wilf Bartlett	Income from working in the forest	17. Fundamental to meeting basic needs of local communities	ECV
	Cutting timber for houses, wharves, stages, and boats		All ECV except timber for houses
	Firewood to heat homes and businesses		ECV
	Company roads to gain access to the wilderness for hunting, fishing, berry picking or tree species that are of no value for the paper mill.		ECV

APPENDIX 4 HIGH CONSERVATION VALUE FOREST ASSESSMENT COMMITTEE (FOR HCVF REPORT)

Dr. Andre Arsenault	Forest Ecologist, Canadian Forest Service, Natural Resources Canada
Dr. Brian Hearn	Wildlife Ecologist, Canadian Forest Service, Natural Resources Canada
Dr. Erin Kelly*	Post-doctoral Fellow in Social Values in Forestry, Grenfell Campus, MUN
Dr. Blair Adams	Regional Ecologist, Newfoundland Department of Natural Resources
Dr. Ivan Emke**	Associate Professor, Social/Cultural Studies, Grenfell Campus, MUN
Barry Elkins	Planning and Development Superintendent, Corner Brook Pulp and Paper Ltd.
Faron Knott	Environmental Management Representative, Corner Brook Pulp and Paper Ltd.

Sean Dolter, Model Forest of Newfoundland and Labrador, facilitated the meetings.

Debbie Hearn, Hearn Consulting Inc., provided resources and recorded decisions for the FECV Assessment Report.

*Dr. Kelly served on the Committee for the initial assessment of potential FECVs. Dr. Kelly resigned from the Committee in July 2012.

**Dr. Emke joined the Committee in November 2012.

APPENDIX 5 POSSIBLE EXCEPTIONAL CONSERVATION VALUES (FORMERLY HCVS)

There are a number of “species at risk as listed by international, national or provincial authorities” that occur on the island of Newfoundland. However, for some of these species, their occurrence on CBPPL limits is not clear so they have been identified as Possible ECV. Because there is a knowledge gap in their occurrence, habitat preference, food availability, etc., it is impossible to develop management strategies for their maintenance. These species and other candidate values have been included in the table below until enough information becomes available that they can be assessed.

Candidate Values	Scientific Name
Terre-nueve Vallonia	<i>Vallonia terraenovae</i>
Vole Ears	<i>Erioderma mollissimum</i>
Cutleaf Fleabane	<i>Erigeron composites</i>
Whorled Aster	<i>Ocelmena acuminate</i>
Other arboreal lichens	
Mosses except Bryum	
Montane Trematodon Moss	<i>Trematodon montanus</i>
Philonotis Moss	<i>Philonotis yezoana</i>
Balsam Poplar	<i>Populus balsamifera</i>

APPENDIX 6 VASCULAR PLANTS OCCURRING ON THE DFA THAT HAVE BEEN RANKED S1-S3 AT THE SUBNATIONAL (S) LEVEL IN THE GENERAL STATUS OF WILD SPECIES

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
1. <u>Shoreline/Aquatic</u> (not usually in forest)				
tall scouring rush	Equisetum hyemale subsp. affine	S1	Secure	1
thimbleweed, riverbank anemone	Anemone virginiana var. alba	S1	Apparently secure	1
lake sedge	Carex lacustris	S1	Secure	1
stellate sedge	Carex radiata	S1	Secure	1
bald spike-rush, bald spikerush, spikesedge	Eleocharis erythropoda	S1	Secure	1
slender spikerush, quill spikerush	Eleocharis nitida	S1	Apparently secure	1
Englemann's rush	Juncus subcaudatus	S1	Secure	1
southern mudwort	Limosella australis	S1	Secure	1
tufted loosestrife, swamp loosestrife	Lysimachia thyrsiflora	S1	Secure	1
leafy pondweed	Potamogeton foliosus subsp. foliosus	S1	Secure	1
bluntleaf pondweed	Potamogeton obtusifolius	S1	Secure	1
Robbins' pondweed	Potamogeton robbinsii	S1	Secure	1
Proserpinaca pectinata	Proserpinaca pectinata	S1	Secure	1
hooked crowfoot	Ranunculus recurvatus	S1	Secure	1
Broadleaf Arrowhead, wapato, duck potato	Sagittaria latifolia	S2	Secure	1
stalked bulrush	Scirpus pedicellatus	S1	Apparently secure	1
slender wedgescale, early bunchgrass	Sphenopholis intermedia	S1	Secure	1
Tradescant's or shore aster	Symphyotrichum tradescantii	S1	Apparently secure	1
twigrush	Cladium mariscoides	S1	Secure	1, 2
Labrador milkvetch	Astragalus alpinus var. brunetianus	S2S3	Vulnerable	1, 3
Farwell's watermilfoil	Myriophyllum farwellii	S1	Secure	1
Fries pondweed, flatstalk pondweed	Potamogeton friesii	S2	Secure	1
upland bent	Agrostis perennans	S2	Secure	1
northern waterstarwort	Callitriche hermaphrodita	S2	Secure	1
cyperus sedge	Carex pseudocyperus	S2	Secure	1
fireberry hawthorn, roundleaf hawthorn	Crataegus chrysocarpa var. chrysocarpa	S2	Secure	1
small waterwort	Elatine minima	S2	Secure	1
pale St. Johnswort	Hypericum ellipticum	S3	Secure	1
knotted rush	Juncus nodosus	S2	Secure	1
creeping rush	Juncus subtilis	S2	Secure	1
auricled twayblade	Listera auriculata	S2	Vulnerable	1
spiked watermilfoil	Myriophyllum sibiricum	S2	Secure	1
whorled watermilfoil	Myriophyllum verticillatum	S2	Secure	1
nodding waternymph, slender naiad	Najas flexilis	S2	Secure	1
Kotzebue's grass-of-Parnassus	Parnassia kotzebuei	S2	Secure	1
large purple fringed orchid	Platanthera grandiflora	S3	Secure	1

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
broadleaf pondweed	Potamogeton amplifolius	S2	Secure	1
northern snailseed pondweed	Potamogeton spirillus	S2	Secure	1
Ball's willow	Salix ballii	S2	Secure	1
threesquare, canemaker's rush	Schoenoplectus pungens	S3	Secure	1
great bulrush, softstem bulrush	Schoenoplectus tabernaemontani	S2	Secure	1
floating burreed	Sparganium fluctuans	S2	Secure	1
sago pondweed	Stuckenia pectinata	S2	Secure	1
lanceleaf aster, panicked aster	Symphyotrichum lanceolatum var. lanceolatum	S2	Secure	1
purple bladderwort	Utricularia purpurea	S2	Secure	1
tall mannagrass	Glyceria grandis	S2	Secure	1, 2
Dudley's rush	Juncus dudleyi	S2	Secure	1, 2
purple false oats	Trisetum melicoides	S2	Apparently secure	1, 2
tawny sedge	Carex hostiana	S2	Secure	1, 2, 5
New England sedge	Carex novae-angliae	S2	Secure	1, 7
bristly crowfoot	Ranunculus pensylvanicus	S2	Secure	1, 7
Indian Hemp, clasping leaf dogbane	Apocynum cannabinum	S3	Secure	1
black grass, saltmarsh rush	Juncus gerardii	S2	Secure	1
whitestem pondweed	Potamogeton praelongus	S3	Secure	1
woolgrass	Scirpus cyperinus	S3	Secure	1
American burreed	Sparganium americanum	S3	Secure	1
lesser brown sedge	Carex adusta	S3	Secure	1, 6
hay sedge	Carex foenea	S3	Secure	1, 6
leathery grapefern	Botrychium multifidum	S3	Secure	1, 7
blue skullcap, maddog skullcap	Scutellaria lateriflora	S3	Secure	1, 7
bayberry willow	Salix myricoides var. myricoides	S3S4	Apparently secure	1, 3
necklace sedge	Carex projecta	S3	Secure	1
bulbous waterhemlock	Cicuta bulbifera	S3	Secure	1
marsh horsetail	Equisetum palustre	S3	Secure	1
winterberry	Ilex verticillata	S3	Secure	1
bayonet rush, jointed bog rush	Juncus militaris	S3	Secure	1
cutleaf bugleweed, waterhorehound	Lycopus americanus	S3	Secure	1
slender watermilfoil	Myriophyllum tenellum	S3	Secure	1
fragrant waterlily, water nymph	Nymphaea odorata subsp. odorata	S3	Secure	1
grass-leaf arrowhead; grassy arrowhead	Sagittaria graminea	S3	Secure	1
cottony willow, stiff willow	Salix eriocephala	S3	Secure	1
hardstem bulrush	Schoenoplectus acutus var. acutus	S3	Secure	1
water Parsnip	Sium suave	S3	Secure	1
freshwater cordgrass, sloughgrass	Spartina pectinata	S3	Secure	1
hiddenfruit bladderwort	Utricularia geminiscapa	S3	Apparently secure	1
northern yelloweyed grass	Xyris montana	S3	Secure	1
fewflowered spikerush	Eleocharis quinqueflora	S3	Secure	1, 2
marsh fern	Thelypteris palustris var. pubescens	S3	Secure	1, 2
northern bog violet	Viola nephrophylla	S3	Secure	1, 2, 3, 6
dogmint, wild basil	Clinopodium vulgare	S3	Secure	1, 3
variegated sedge	Carex stylosa	S3	Secure	1, 5

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
milky willowherb	<i>Epilobium lactiflorum</i>	S3	Sensitive	1, 5
white dock, seabeach dock	<i>Rumex pallidus</i>	S2S4	Apparently secure	1
northern water plantain	<i>Alisma trivale</i>	S2	Secure	1, 2
elegant milkvetch	<i>Astragalus eucosmus</i>	S3	Secure	1, 3
least moonwort	<i>Botrychium simplex</i>	S2	Apparently secure	1, 3, 7
curved sedge	<i>Carex maritima</i>	S2	Secure	1
retorse sedge	<i>Carex retrorsa</i>	S1	Secure	1
sea-beach sedge	<i>Carex silicea</i>	S2	Secure	1
woolly panic grass	<i>Dichanthelium acuminatum</i> var. <i>fasciculatum</i>	S2S3	Secure	1
rough horsetail	<i>Equisetum hyemale</i>	S1	Secure	1
Arctic red fescue	<i>Festuca rubra</i> subsp. <i>arctica</i>	S2S3	Secure	1
purple false oats	<i>Graphephorum melicoides</i>	S2S3	Secure	1
purple rattlesnake-root	<i>Prenanthes racemosa</i>	S1	Secure	1
white water lily	<i>Nymphaea odorata</i>	S3	Secure	1
field oxytrope	<i>Oxytropis campestris</i>	S3	Secure	1
Saint John River oxytrope	<i>Oxytropis campestris</i> var. <i>johannensis</i>	S3	Secure	1
grass-of-parnassus	<i>Parnassia palustris</i>	S2S3	Secure	1, 2
sharp-fruited knotweed	<i>Polygonum oxyspermum</i> subsp. <i>raii</i>	S2	Apparently secure	1, 2
leafy pond weed	<i>Potamogeton foliosus</i>	S1S2	Secure	1
hard-stemmed club-bulrush	<i>Schoenoplectus acutus</i>	S3	Secure	1
water bulrush	<i>Schoenoplectus subterminalis</i>	S3	Secure	1
horned pondweed	<i>Zannichellia palustris</i>	S2S3	Secure	1, 2
2. Wetlands				
slenderleaf sundew	<i>Drosera linearis</i>	S2	Apparently secure	2
close-sheathed cottongrass	<i>Eriophorum brachyantherum</i>	S2	Secure	2
fen grass-of-Parnassus	<i>Parnassia glauca</i>	S2S3	Secure	2
bog willow	<i>Salix pedicellaris</i>	S2S3	Secure	2
autumn willow	<i>Salix serissima</i>	S2S3	Secure	2
green addersmouth	<i>Malaxis unifolia</i>	S3	Secure	2
pod grass, marsh scheuchzeria, rannoch rush	<i>Scheuchzeria palustris</i>	S3S4	Secure	2
curly-grass fern	<i>Schizaea pusilla</i>	S3S4	Vulnerable	2
Fernald's serviceberry	<i>Amelanchier fernaldii</i>	S1	Vulnerable	1, 2
Virginia screwstem	<i>Bartonia virginica</i>	S1	Secure	2
water arum	<i>Calla palustris</i>	S2S3	Secure	1, 2
Northeastern sedge	<i>Carex cryptolepis</i>	S1	Apparently secure	2
marsh straw sedge	<i>Carex hormathodes</i>	S3	Apparently secure	2
pointed broom sedge	<i>Carex scoparia</i>	S3	Secure	2, 7
English sundew	<i>Drosera anglica</i>	S3	Secure	2
variegated scouring-rush	<i>Equisetum variegatum</i>	S3	Secure	1, 2
slender cottongrass	<i>Eriophorum gracile</i>	S1S2	Secure	2
russet cottongrass	<i>Eriophorum russeolum</i>	S3	Secure	2
roundleaf orchid	<i>Galearis rotundifolia</i>	S2	Secure	2

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
long-styled rush	<i>Juncus longistylis</i>	S1	Secure	1, 2
Carolina sea lavender	<i>Limonium carolinianum</i>	S2S3	Secure	2
auricled twayblade	<i>Neottia auriculata</i>	S2	Vulnerable	1, 2, 4
seaside goldenrod	<i>Solidago sempervirens</i>	S2S3	Secure	2
saltmeadow cordgrass	<i>Spartina patens</i>	S2	Secure	2
smooth cordgrass	<i>Sporobolus alterniflorus</i>	S2	Secure	2
seablite	<i>Suaeda calceoliformis</i>	S1S2	Secure	2, 7
herbaceous seablite	<i>Suaeda maritima</i>	S3	Secure	2
northern bog aster	<i>Symphyotrichum boreale</i>	S1	Secure	2
Gaspé Peninsula arrowgrass	<i>Triglochin gaspensis</i>	S3	Apparently secure	2
marsh valerian	<i>Valeriana dioica</i>	S3	Secure	2
3. Cliffs and Talus Slopes				
maidenhair spleenwort	<i>Asplenium trichomanes</i> subsp. <i>trichomanes</i>	S1S2	Secure	3
rock dwelling sedge	<i>Carex petricosa</i> var. <i>misandroides</i>	S1	Vulnerable	3
pendantpod oxytrope	<i>Oxytropis deflexa</i> var. <i>foliolosa</i>	S1S2	Secure	3
Laurentian dandelion	<i>Taraxacum laurentianum</i>	S1	Critically imperiled	3
alpine cliffbrake, northern woodsia	<i>Woodsia alpina</i>	S1S2	Apparently secure	3
prolific fescue, proliferous red fescue	<i>Festuca prolifera</i>	S3	Apparently secure	3
cutleaf anemone, cliff anemone	<i>Anemone multifida</i> var. <i>multifida</i>	S3	Secure	3
bulblet bladder fern	<i>Cystopteris bulbifera</i>	S2S3	Secure	3
yellow mountain avens, Drummond's dryad	<i>Dryas drummondii</i>	S2	Secure	3
Rocky Mountain fescue	<i>Festuca saximontana</i> subsp. <i>saximontana</i>	S2	Secure	3
serpentine sandwort, dryleaf sandwort	<i>Minuartia marcescens</i>	S2	Vulnerable	3
coastal cinquefoil	<i>Potentilla litoralis</i>	S2S3	Secure	3
spreading dogbane	<i>Apocynum androsaemifolium</i>	S3	Secure	3
Steller's rockbrake	<i>Cryptogramma stelleri</i>	S2S3	Secure	3
limestone oak fern	<i>Gymnocarpium robertianum</i>	S3	Secure	3
arctic bladderpod	<i>Lesquerella arctica</i>	S3	Apparently secure	3
white addersmouth, Northern white addersmouth	<i>Malaxis monophyllos</i> var. <i>brachypoda</i>	S3	Secure	3
alpine ragwort; alpine groundsel	<i>Packera pauciflora</i>	S2S3	Secure	3, 5
Lapland rosebay	<i>Rhododendron lapponicum</i>	S3S4	Secure	3
smooth cliffbrake	<i>Woodsia glabella</i>	S3	Secure	3
Aleutian madienhair fern	<i>Adiantum aleuticum</i>	S3	Secure	3
alpine pussytoes	<i>Antennaria alpina</i>	S3	Secure	3
pulvinate pussytoes	<i>Antennaria rosea</i> subsp. <i>pulvinata</i>	S3S4	Secure	3
bristleleaf sedge	<i>Carex eburnea</i>	S3	Secure	3
rock sedge	<i>Carex rupestris</i>	S3	Secure	3
Sitka clubmoss, tufted groundceder	<i>Diphasiastrium sitchense</i>	S3S4	Secure	3
rockcress whitlowgrass	<i>Draba arabisans</i>	S3	Apparently secure	3
viviparous fescue	<i>Festuca frederikseniae</i>	S3	Vulnerable	3

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
alpine sweetvetch	Hedysarum alpinum	S3	Secure	3
alpine lady fern	Athyrium distentifolium	S2	Apparently secure	3
shaved sedge	Carex tonsa	S1	Secure	3
parasol sedge	Carex umbellata	S2	Secure	3, 6
serpentine sandwort	Cherleria marcescens	S2	Vulnerable	3, 5
laurentian fragile fern	Cystopteris laurentiana	S2	Vulnerable	3
timber oatgrass	Danthonia intermedia	S1S2	Secure	3, 5, 6
hairy rosette-panicgrass	Dichanthelium acuminatum	S2S3	Secure	3, 4, 5
fragrant woodfern	Dryopteris fragrans	S2S3	Secure	3
marginal wood fern	Dryopteris marginalis	S1	Secure	1, 3
crowded wormseed mustard	Erysimum coarctatum	S1	Apparently secure	3, 4, 6
Rocky Mountain fescue	Festuca saximontana var. saximontana	S2	Secure	3, 4
alpine sweet-vetch	Hedysarum americanum	S3	Secure	3
pine-barren false heather	Hudsonia ericoides	S2	Apparently secure	3, 4
Chinese clubmoss	Huperzia miyoshiana	S2	Apparently secure	3
mountain sorrel	Oxyria digyna	S2S3	Secure	3
arctic bladderpod	Physaria arctica	S3	Apparently secure	3
coast cinquefoil	Potentilla litoralis	S2S3	Secure	3
snow cinquefoil	Potentilla nivea	S3	Secure	3
4. Forest (at least occasionally, including openings and cutovers)				
Menzies' rattlesnake plantain	Goodyera oblongifolia	S1	Secure	4
tapering sweetcicely	Osmorhiza berteroi	S2	Secure	4
small roundleaf orchid, one leaf orchid	Amerorchis rotundifolia	S2	Secure	2, 4
pipsissewa, prince's pine	Chimaphila umbellata	S2	Secure	4
Carolina spring beauty	Claytonia caroliniana	S2	Secure	1, 4
blunt sweetcicely	Osmorhiza depauperata	S3	Secure	4
red pine	Pinus resinosa	S2	Secure	4
hollyfern	Polystichum lonchitis	S3	Secure	3, 4
Macoun's buttercup	Ranunculus macounii	S2S3	Secure	1, 4, 6, 7
Fernald's false mannagrass	Torreyochloa pallida var. fernaldii	S2S3	Secure	1, 2, 4, 7
northern valerian, marsh valerian	Valeriana dioica subsp. sylvatica	S3	Secure	2, 4
Selkirk's violet, great spurred violet	Viola selkirkii	S2	Secure	1, 3, 4, 5
woodland agrimony, roadside agrimony	Agrimonia striata	S3	Secure	1, 4, 6
forest bluegrass, weak meadowgrass	Poa saltuensis	S3S4	Secure	1, 4, 6
thinstemmed lady's mantle	Alchemilla filicaulis subsp. filicaulis	S3	Apparently secure	1, 3, 4, 5, 7
dwarf mistletoe	Arceuthobium pusillum	S3	Secure	4
Yellow Birch	Betula alleghaniensis	S3	Secure	4
northern shorthusk	Brachyelytrum septentrionale	S3S4	Secure	1, 4
graceful sedge	Carex gracillima	S3S4	Secure	1, 4
pale sedge	Carex pallescens	S3	Secure	1, 4, 5, 7
longstalked sedge	Carex pedunculata	S3	Secure	1, 4
alternatleaf dogwood, green osier	Cornus alternifolia	S3S4	Secure	1, 4

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
showy ladyslipper, queen lady's-slipper	Cypripedium reginae	S3	Apparently secure	2, 4
bog willowherb	Epilobium leptophyllum	S3	Secure	2, 4
meadow horsetail	Equisetum pratense	S3	Secure	
black ash	Fraxinus nigra	S2S3	Apparently secure	3, 4
northern wild licorice	Galium kamtschaticum	S3S4	Apparently secure	4, 5
checkered rattlesnake plantain	Goodyera tessellata	S3S4	Secure	4
piresap, false beechdrops	Monotropa hypopitys	S3	Secure	4
whitegrain mountainrice	Oryzopsis asperifolia	S3	Secure	4
interrupted fern	Osmunda claytoniana	S4	Secure	1, 4, 6
Arctic sweet coltsfoot	Petasites frigidus var. palmatus	S3S4	Secure	2, 4
eastern white pine	Pinus strobus	S3	Secure	4
plumboy, arctic bramble	Rubus arcticus subsp. acaulis	S3S4	Secure	3, 4
Canada yew, American yew, ground-hemlock	Taxus canadensis	S3S4	Secure	4
nodding trillium, nodding wake-robin	Trillium cernuum	S4	Secure	1, 4
kidneyleaf violet	Viola renifolia	S3	Secure	3, 4
Canada anemone	Anemonastrum canadense	S1	Secure	4, 7
daisy-leaved moonwort	Botrychium matricariifolium	S2S3	Secure	4
stubby-stalked cladonia	Cladonia caespiticia	S3	Secure	4
Vreeland's striped coralroot	Corallorhiza striata var. vreelandii	S1	Apparently secure	2, 4
fireberry hawthorn	Crataegus chrysocarpa	S2	Secure	4
fan clubmoss	Diphasiastrum digitatum	S2	Secure	4, 7
Wintergreen	Gaultheria procumbens	S1	Secure	4, 7
black huckleberry	Gaylussacia baccata	S3	Secure	4
herb-robert	Geranium robertianum	S3	Secure	4
piresap	Hypopitys monotropa	S3	Secure	4
pale jewelweed	Impatiens pallida	S1	Secure	4
hairy woodrush	Luzula acuminata	S1	Secure	4
one-cone clubmoss	Lycopodium lagopus	S2	Secure	4, 7
large false Solomon's seal	Maianthemum racemosum	S1	Secure	4
false Solomon's seal	Maianthemum racemosum subsp. racemosum	S1	Secure	4
wood millet	Milium effusum var. cisatlanticum	S2S3	Secure	4
partridgeberry	Mitchella repens	S2S3	Secure	4
northern twayblade	Neottia borealis	S1	Secure	2, 4
whorled wood aster	Oclemena acuminata	S1	Secure	4
mountain woodsorel	Oxalis montana	S1S2	Secure	4
brown-eyed shingle lichen	Pannaria rubiginosa	S3	Apparently secure	4
Degen's felt pelt lichen	Peltigera degenii	S1S3	Apparently secure	4
a pelt lichen	Peltigera horizontalis	S1S3	Secure	4
Canadian rice grass	Piptatheropsis canadensis	S2	Apparently secure	3, 4, 6
Hooker's bog-orchid	Platanthera hookeri	S2	Apparently secure	4
yellow specklebelly	Pseudocyphellaria holarctica	S2S3	Secure	4
shinleaf	Pyrola elliptica	S2S3	Secure	4
punctured ribbon lichen	Ramalina dilacerata	S1S3	Secure	4

Habitat Group Common Name	Scientific Name	S Rank 2024	General Status 2024	Habitat Group(s)
angel's hair lichen	Ramalina thrausta	S2S3	Secure	4
red penny moss	Rhizomnium punctatum	S3	Secure	4, 7
leathery grapefern	Sceptridium multifidum	S3	Secure	4, 7
5. <u>Alpine /Subalpine</u>				
Lapland buttercup	Ranunculus lapponicus	SH	Secure	5
arctic willow	Salix arctica	S2	Secure	5
alpine fescue, shortleaf fescue	Festuca brachyphylla subsp. brachyphylla	S2S3	Secure	5
blue mountainheath	Phyllodoce caerulea	S3	Secure	5
cutleaf anemone	Anemone multifida	S3	Secure	5
glandular birch	Betula glandulosa	S3	Secure	2, 5
cutleaf daisy	Erigeron compositus	S1	Secure	5
rough fescue	Festuca altaica	S2	Secure	5
moss-plant	Harrimanella hypnoides	S2	Secure	5, 7
rock stitchwort	Sabulina dawsonensis	S2S3	Secure	5
alpine catchfly	Viscaria alpina	S3	Secure	3, 5
6. <u>Roadsides and Ditches</u>				
haysented fern	Dennstaedtia punctilobula	S1	Secure	6
mosquito bulrush	Scirpus hattorianus	S3S4	Secure	6
Houghton's sedge	Carex houghtoniana	S1	Secure	4, 6
red fescue	Festuca rubra	S2S3	Secure	6, 7
fox-tail barley	Hordeum jubatum subsp. jubatum	S2S3	Secure	6, 7
round-fruited pinweed	Lechea intermedia	S1	Secure	4, 6
Kentucky blue grass	Poa pratensis	S3	Secure	4, 6, 7
yellow rattle	Rhinanthus minor	S3	Secure	6, 7
7. <u>Meadows and Grassy Areas</u>				
As indicated in column five – Habitat Group(s)				

APPENDIX 7 SPECIES AT RISK ON THE ISLAND OF NEWFOUNDLAND

Common Name	Scientific Name	Global Conservation Status	National SARA COSEWIC Recommended	Provincial ESA SSAC Recommended	Habitat on DFA?¹	Species Found on DFA?¹,⁶
Mammals						
American Marten ^{2,4}	<i>Martes americana atrata</i>	G5T1	Threatened	Threatened	Yes	Yes
Little Brown Myotis ^{4,5}	<i>Myotis lucifugus</i>	G3	Endangered		Yes	Yes
Hoary Bat ⁴	<i>Lasiurus cinereus</i>	G3	Endangered		No	No
Northern Long-eared Myotis ^{4,5}	<i>Myotis septentrionalis</i>	G2	Endangered		Yes	Yes
Woodland Caribou (Nfld. Pop'n) ^{4,5}	<i>Rangifer tarandus</i>	G5TNR	Special Concern		Yes	Yes
Birds						
Bank Swallow ^{2,4,5}	<i>Riparia riparia riparia</i>	G5	Threatened		Yes	Yes
Barn Swallow ^{4,5}	<i>Hirundo rustica</i>	G5	Special Concern		Yes	Yes
Barrow's Goldeneye ^{2,4,5}	<i>Bucephala islandica</i>	G5T3	Special Concern	Vulnerable	Yes	Yes
Bobolink ^{2,4,5}	<i>Dolichonyx oryzivorus</i>	G5	Special Concern	Vulnerable	Yes	Yes
Chimney Swift ^{2,4}	<i>Chaetura pelagica</i>	G4	Threatened	Threatened	Yes	No
Common Nighthawk ^{2,4}	<i>Chordeiles minor</i>	G5	Special Concern	Vulnerable	Yes	Yes
Eskimo Curlew ^{2,4}	<i>Numenius borealis</i>	GH	Endangered	Endangered	No	No
Evening Grosbeak	<i>Coccothraustes vespertinus</i>	G5	Special Concern		Yes	Yes
Harlequin Duck ^{2,4,5}	<i>Histrionicus histrionicus</i>	G4T4	Special Concern	Vulnerable	Yes	Yes
Ivory Gull ^{2,4,5}	<i>Pagophila eburnea</i>	G4	Endangered	Endangered	Yes	Yes
Newfoundland Gray-cheeked Thrush ^{2,4}	<i>Catharus minimus minimus</i>	G5T4	Threatened	Threatened	Yes	Yes
Northern Goshawk	<i>Accipiter gentilis atricapillus</i>	G5T5	Not at Risk		Yes	Yes
Olive-sided Flycatcher ^{2,4,5}	<i>Contopus cooperi</i>	G4	Special Concern	Vulnerable	Yes	Yes
Peregrine Falcon ^{2,4,5}	<i>Falco peregrinus</i> subsp. <i>tundrius</i>	G4T3Q	Not at Risk	Vulnerable	No	No
Peregrine Falcon ^{2,3,4}	<i>Falco peregrinus</i> subsp. <i>anatum</i>	G4T4	Not at Risk	Vulnerable	Yes	Yes
Piping Plover ^{2,4,5}	<i>Charadrius melodus melodus</i>	G3	Endangered	Endangered	Yes	Yes
Red Crossbill ^{2,4,5}	<i>Loxia curvirostra</i> subsp. <i>percna</i>	G5T1T2Q	Threatened	Threatened	Yes	Yes
Red Knot ^{2,4,5}	<i>Calidris canutus</i> subsp. <i>rufa</i>	G4T2	Endangered	Endangered	Yes	Yes

Common Name	Scientific Name	Global Conservation Status	National SARA COSEWIC Recommended	Provincial ESA SSAC Recommended	Habitat on DFA? ¹	Species Found on DFA? ^{1,6}
Red-tailed Hawk ³	<i>Buteo jamaicensis</i>	G5	Not at Risk		Yes	Yes
Rough-legged Hawk ^{2,5}	<i>Buteo lagopus</i>	G5	Not at Risk		Yes	Yes
Rusty Blackbird ^{2,4,5}	<i>Euphagus carolinus</i>	G4	Special Concern	Vulnerable	Yes	Yes
Sharp-shinned Hawk ^{4,5}	<i>Accipiter striatus</i>	G5	Not at Risk		Yes	Yes
Short-eared Owl ^{2,4,5}	<i>Asio flammeus</i>	G5	Special Concern	Threatened	Yes	Yes
Song Sparrow ^{2,5}	<i>Melospiza melodia</i>	G5			Yes	Yes
Winter Wren ^{2,5}	<i>Troglodytes hiemalis</i>	G5			Yes	Yes
Fishes						
American Eel ^{2,4,5}	<i>Anguilla rostrata</i>	G4	Threatened	Vulnerable	Yes	Yes
Atlantic Salmon ^{4,5} (South. Nfld Pop'n)	<i>Salmo salar</i>	G5TNRQ	Threatened		No	No
Banded Killifish ^{2,4,5}	<i>Fundulus diaphanus</i>	G5TNRQ	Special Concern	Vulnerable	Yes	Yes
Arthropods						
Gypsy Cuckoo Bumble Bee ⁴	<i>Bombus bohemicus</i>	G4	Endangered			
Suckley's Cuckoo Bumble Bee	<i>Bombus suckleyi</i> ^{4,5}	G2	Threatened			
Yellow-banded Bumble Bee ^{4,5}	<i>Bombus terricola</i>	G3	Special Concern			Yes
Tomah Mayfly ⁵	<i>Siphoniscia aerodromia</i>	G2				
A Mayfly ⁵	<i>Siphonurus barbaroides</i>	G3				
A Mayfly ⁵	<i>Baetisca rubescens</i>	G3				
Transverse Lady Beetle ^{4,5}	<i>Coccinella transversoguttata</i>	G5	Special Concern			Yes
Troubled Northern Caddisfly ⁵	<i>Limnephilus ademus</i>	G4				
Short-tailed Swallowtail ⁵	<i>Papilio brevicauda brevicauda</i>	G4T4T5				
A Mayfly ⁵	<i>Acentrella feropagus</i>	G4				
Molluscks						
Maritime Ambersnail ⁵	<i>Oxyloma verrilli</i>	G4				
Terre-nueve Vallonia ⁵	<i>Vallonia terraenovae</i>	G2				
Olive Vertigo ⁵	<i>Vertigo perryi</i>	G3				
Vascular Plants						

Common Name	Scientific Name	Global Conservation Status	National SARA COSEWIC Recommended	Provincial ESA SSAC Recommended	Habitat on DFA?¹	Species Found on DFA?¹,⁶
Acadian Quillwort ⁵	<i>Isoetes acadiansis</i>	G3				No
Alaska Rein Orchid ²	<i>Platanthera unalascensis</i>	G5		Endangered		No
Alberton Alkali Grass ⁵	<i>Puccinellia ambigua</i>	G3				No
Alpine Silverwort ⁵	<i>Anthelia julacea</i>	G3				No
Auricled Twayblade ⁵	<i>Listera auriculata</i>	G3			Yes	Yes
Baltic Saltbrush ⁵	<i>Atriplex nudicaulis</i>	G5			No	No
Barrens Willow ^{2,5}	<i>Salix jejuna</i>	G1	Endangered	Endangered	No	No
Black Ash	<i>Fraxinus nigra</i>				Yes	Yes
Bodin's Milkvetch ²	<i>Astragalus bodinii</i>	G4		Threatened	No	No
Crowded Wormseed Mustard ^{2,5}	<i>Erysimum inconspicuum</i> var. <i>coarctatum</i>	G4G5		Endangered	Yes	Yes
Curly-grass Fern ⁵	<i>Schizaea pusilla</i>	G3			Yes	Yes
Cutleaf Fleabane ²	<i>Erigeron compositus</i>	G5		Endangered	Yes	Yes
Dense Whitlow-grass ^{2,4,5}	<i>Draba pycnosperma</i>	G2	Special Concern	Vulnerable		No
Feathery False Solomon's Seal ^{2,5}	<i>Maianthemum racemosum</i> subsp. <i>racemosum</i>	G5T5		Endangered	Yes	Yes
Fernald's Braya ^{2,4,5}	<i>Braya fernaldii</i>	G1	Endangered	Endangered	No	No
Fernald's Milk-vetch ^{2,4,5}	<i>Astragalus robbinsii</i> var. <i>fernaldii</i>	G5T1T2	Special Concern	Vulnerable	No	No
Fernald's Serviceberry ⁵	<i>Amelanchier fernaldii</i>	G3			Yes	Yes
Gaspe Peninsula Arrow-grass ⁵	<i>Triglochin gaspensis</i>	G4				No
Gmelin's Watercrowfoot ^{2,5}	<i>Ranunculus gmelinii</i>	G5		Endangered		No
Griscom's Arnica ^{2,4,5}	<i>Arnica frigida</i> subsp. <i>Griscomii</i>	G5T1T2	Threatened	Endangered		No
Laurentian Bladderfern ⁵	<i>Cystopteris laurentiana</i>	G3			Yes	Yes
Laurentian Dandelion ⁶	<i>Taraxacum laurentianum</i>	G1			Yes	Yes
Lindley's Aster ^{2,5}	<i>Symphyotrichum ciliolatum</i>	G5		Endangered	Yes	Yes
Long's Braya ^{2,4,5}	<i>Braya longii</i>	G1	Endangered	Endangered	No	No
Low Northern Rockcress ²	<i>Neotorularia humilis</i> spp. <i>humilis</i>	G5T5		Endangered	No	No
MacKenzie's Sweet Vetch ^{2,5}	<i>Hedysarum boreale</i> subsp. <i>Mackenzii</i>	G5T5		Threatened	No	No
Mountain Bladder Fern (Island pop)	<i>Cystopteris montana</i>	G5		Endangered		No
Mountain Fern ^{2,5}	<i>Thelypteris quepaertensis</i>	G4		Vulnerable		No

Common Name	Scientific Name	Global Conservation Status	National SARA COSEWIC Recommended	Provincial ESA SSAC Recommended	Habitat on DFA?¹	Species Found on DFA?¹,⁶
Mountain Holly Fern ^{4,5}	<i>Polystichum scopulinum</i>	G4	Threatened		Yes	Yes
Newfoundland Pussytoes ⁵	<i>Antennaria eucosma</i>	G3				No
North Atlantic Fescue ⁵	<i>Festuca frederikseniae</i>	G3			Yes	Yes
Northern Bog Aster ^{2,5}	<i>Symphyotrichum boreale</i>	G5		Endangered	Yes	Yes
Northern Twayblade	<i>Listera borealis</i>	G5		Endangered	Yes	Yes
Oval-leaf Creeping Spearwort ²	<i>Ranunculus flammula</i> var. <i>ovalis</i>	G5T5		Endangered	No	No
Rattlesnakeroot ^{2,5}	<i>Prenanthes racemosa</i>	G5		Endangered	Yes	Yes
Red Pine	<i>Pinus resinosa</i>	G5		Threatened	Yes	Yes
Robinson's Hawkweed ⁵	<i>Hieracium robinsonii</i>	G3				No
Rock Dwelling Sedge ^{2,5}	<i>Carex petricosa</i> var. <i>misandroides</i>	G4		Endangered	Yes	Yes
Serpentine Stitchwort ⁵	<i>Minuartia marcescens</i>	G3	Not At Risk		Yes	Yes
Sharpleaf Aster ^{2,5}	<i>Oclemena acuminata</i>	G5		Threatened	Yes	Yes
Shaved Sedge ^{2,5}	<i>Carex tonsa</i> var. <i>tonsa</i>	G5T5			Yes	Yes
Tradescant's Aster ^{2,5}	<i>Symphyotrichum tradescantii</i>	G4		Threatened	Yes	Yes
Upward-lobed Moonwort ⁵	<i>Botrychium ascendens</i>	G4				No
Vreeland's Striped Coralroot ²	<i>Corallorhiza striata</i> var. <i>vreelandii</i>	G5T3T5		Endangered	Yes	Yes
Woolly Arnica ^{2,5}	<i>Arnica augustifolia</i> subsp. <i>Tomentosa</i>	G5T5		Endangered		No
Water Pygmyweed ²	<i>Crassula aquatica</i>	G5		Vulnerable	No	No
Mosses and Non-Vascular Plants						
Porsild's Bryum ^{2,5}	<i>Mielichhoferia macrocarpa</i>	G2	Threatened	Threatened	No	No
Splachnoid Trumpet Moss ⁵	<i>Tayloria splachnoides</i>	G2			Possibly	No
Small Four-toothed Moss ⁵	<i>Tetradontium repandum</i>	G2			Possibly	No
Short-leaved Bristle Moss ⁵	<i>Seligeria brevifolia</i>	G2			Likely	No
Yezo's Apple Moss ⁵	<i>Philonotis yezoana</i>	G3			Likely	No
Montane Trematodon Moss ⁵	<i>Trematodon montanus</i>	G2			Possibly	No
Minakata's Hump-backed Elves Moss ⁵	<i>Buxbaumia minakatae</i>	G3				No
Bluntleaf Threadmoss ⁵	<i>Pohlia obtusifolia</i>	G3				No
Red-toothed Extinguisher Moss ⁵	<i>Encalypta longicolla</i>	G3				No

Common Name	Scientific Name	Global Conservation Status	National SARA COSEWIC Recommended	Provincial ESA SSAC Recommended	Habitat on DFA? ¹	Species Found on DFA? ^{1,6}
Copper Coscinodon ⁵	<i>Coscinodon cribrosus</i>	G3				No
Large Brook Moss ⁵	<i>Hygrohypnum subeugyrium</i>	G3				No
Carrion Moss ⁵	<i>Aplodon wormskjoldii</i>	G4				No
Spring Hook Moss ⁵	<i>Warnstorfia pseudostraminea</i>	G3				No
Carolina Peat Moss ⁵	<i>Sphagnum carolinianum</i>	G3				No
Saltmarsh Bryum ⁵	<i>Bryum salinum</i>	G3				No
Schimper's Swan-neck Moss ⁵	<i>Campylopus schimperi</i>	G3				No
Blunted Earwort ⁶	<i>Diplophyllum obtusatum</i>	G2				No
Lichens						
Appalachian Matchsticks	<i>Pilophorus fibula</i>	No rank				No
Blue Felt Lichen ⁴	<i>Degelia plumbea</i>	G4	Special Concern	Vulnerable	Yes	No
Boreal Felt Lichen ^{2,4,5,7}	<i>Erioderma pedicellatum</i>	G2	Endangered	Vulnerable	Possibly	No
Vole Ears ⁵	<i>Erioderma mollissimum</i>	G4	Endangered			No
Red Firedot Lichen ⁶	<i>Polycauliona leuteominia</i>	G2				No
Witch's Hair Lichen ⁶	<i>Alectoria fallacina</i>	G2				No
Wrinkled Shingle Lichen ⁴	<i>Pannaria lurida</i>	G4	Threatened	Threatened		No
Lichen ⁶	<i>Porpidia grisea</i>	G2				No
Lichen ⁶	<i>Fuscidea lowensis</i>	G1				No
Lichen ⁶	<i>Thelocarpon epibolum</i>	G5				No
Fungus ⁶	<i>Phaeocollybia gregaria</i>	G2				No

¹ Confirmed by NL Wildlife Division Personnel: Shelley Garland

² SSAC Species Status Reports

³ CITES

⁴ COSEWIC


⁵ NatureServe Explorer

⁶ Atlantic Canada Conservation Data Centre


⁷ IUCN Red List

Common Name	Scientific Name	Global Conservation Status	<u>National</u> SARA COSEWIC Recommended	<u>Provincial</u> ESA SSAC Recommended	Habitat on DFA? ¹	Species Found on DFA? ^{1,6}
-------------	-----------------	----------------------------------	---	---	------------------------------------	--


APPENDIX 8 MAPS OF EXCEPTIONAL CONSERVATION VALUES (FORMERLY HCVs)




Map 1. Duck Unlimited Canada projects on the DFA.




Map 2. Newfoundland Marten core areas and critical habitat on the DFA.




Map 3. Protected and conservation areas on the DFA.



Map 4. Large intact landscape on Crown land



Map 5. Unfragmented forest areas on the DFA.



Map 6. Protected Water Supply Areas on the DFA.