Anxiety Disorders					
Separation Anxiety Disorder	Selective Mutism	Specific Phobia	Social Anxiety Disorder	Panic Disorder	Agoraphobia
EXCESSIVE FEAR/ANXIETY SEPARATING from attached individuals/figures (3+)  1. Distress anticipating/experiencing separation from attachment figures  2. Worry re: losing attachment figures, harm coming to them  3. Worry re: events causing separation  4. Reluctance to go out d/t separation	Consistent failure to speak in specific social situations when expected, but speaking in other situations     Not due to language difficulties	Fear about specific object/situation	Fear about social situations where exposed to possible scrutiny  Social interactions, being observed, performing In children, must be in peer settings	RECURRENT PANIC ATTACKS (4+) Unexpected, abrupt, peak in minutes 1. Palpitations 2. Sweating 3. Shaking 4. Shortness of breath 5. Choking feeling 6. Chest pain 7. Nausea 8. Dizziness 9. Chills/heat sensation 10.Parasthesias 11.Derealization/depersonalization 12.Fear of losing control 13.Fear of dying	FEAR OF 5 SPECIFIC SITUATIONS (2+)  1. Public transportation 2. Open spaces 3. Enclosed spaces 4. Crowds 5. Outside home alone
5. Reluctance being without figures 6. Reluctance to sleep without figures 7. Nightmares re: separation 8. Physical c/o if separation/anticipated			Fear of negative evaluation due to behavior/anxiety sx		Fears b/c escape may be difficult, or help may not be available (in event of developing panic sx, incapacity sx, embarrassing sx
		Phobic object almost always provokes immediate fear/anxiety	Social situations almost always provoke fear/anxiety	>1 MONTH OF RELATED SX (1+)  1. Worry about more panic attacks or their consequences	Agoraphobic situations almost always provoke fear/anxiety
		Phobic object actively avoided or endured with intense fear/anxiety	Social situations avoided or endured with intense fear/anxiety	Maladaptive behavior change related to attacks (avoidance)	Agoraphobic situations actively avoided, endured with intense fear/anxiety, or require companion
		Fear/anxiety out of proportion to actual danger of phobic object	Fear/anxiety out of proportion to actual threat of social situation		Fear/anxiety out of proportion to actual threat of agoraphobic situation
Duration >4 weeks in children     Duration >6 months in adults	Duration >1 month     Not first month of school	Duration >6 months	Duration >6 months		Duration >6 months
Sig distress/fxnal impairment	Interferes with social/school/work	Sig distress/fxnal impairment	Sig distress/fxnal impairment		Sig distress/fxnal impairment
Not due to another mental disorder (autism, psychosis, agoraphobia, GAD, IAD)	Not due to communication disorder, autism, schizophrenia, psychosis	Not due to another mental disorder (agoraphobia, OCD, PTSD, SAD, separation anxiety)	Not due to another mental disorder (panic, body dysmorphic, autism)	Not due to another mental disorder (specific phobia, OCD, PTSD, SAD, separation anxiety)  Not due to substance or AMC	Not due to another mental disorder (specific phobia, body dysmorphia, OCD, PTSD, SAD, separation anxiety)
	Not due to language difficulties		Not due to substance or AMC		
			If AMC present, fear is excessive or unrelated		If AMC present, fear is excessive
SPECIFIERS	None	<ul> <li>Animal</li> <li>Natural environment</li> <li>Blood-injection-injury</li> <li>Situational</li> <li>Other</li> </ul>	Performance only	None	Can have both panic disorder and agoraphobia

Anxiety Disorders										
	Generalized Anxiety Disorder	Substance/Medication-Induced	Due to AMC	Other specified	Unspecified					
-	Excessive anxiety/worry     About number of things     More days than not     Duration >6 months	Panic attacks or anxiety predominate clinical picture	Panic attacks or anxiety predominated clinical picture	Does not meet any full criteria Choose TO communicate specific reason  Limited-symptom attacks Generalized anxiety not occurring more days than not Khyál cap (wind attacks)  Ataque de nervios (attack of nerves)	Does not meet any full criteria     Choose NOT to communicate specific reason     Insufficient information for dx					
	Difficult to control worry  ASSOCIATED SYMPTOMS (3+) 1. Restlessness, keyed up, on edge 2. Fatigued easily 3. Concentration difficulties 4. Irritability 5. Muscle tension 6. Sleep disturbance	Hx, P/E, lab findings of both:  1. Sx onset during/soon after substance intox/withdrawal or medication exposure  2. Sub/med capable of producing sx	Direct pathophysiological consequence of AMC, as per evidence from Hx, P/E, labs							
	Sig distress/fxnal impairment	Sig distress/fxnal impairment	Sig distress/fxnal impairment							
	Not due to another mental disorder (autism, psychosis, agoraphobia, GAD, IAD)	Not due to another anxiety disorder (Sx precede sub/med use, symptoms >1 month after stopping, other evidence)	Not due to another mental disorder							
	Not due to substance or AMC	Not during delirium	Not during delirium							
SPECIFIERS	None	Indicate substance  • With onset during intoxication  • With onset during withdrawal  • With onset after medication use	Indicate name of AMC							