Trauma and Stressor-Related Disorders					
Reactive Attachment	Disinhibited Social Engagement	PTSD	PTSD (children 0-6)	Acute Stress Disorder	Adjustment Disorders
A EMOTIONALLY WITHDRAWN towards adult caregivers when distressed (2) 1. Minimally seeks comfort 2. Minimally responds to comfort	APPROACHES UNFAMILIAR PPL (2+) 1. Decr reticence approaching 2. Overfamiliar verbal/physical behave 3. Decr checking back with caregiver 4. Willingness to go off, no hesitation	EXPOSURE to trauma (1+) 1. Direct experience 2. Witnessing in person 3. To close family/friend 4. Repeated/extreme exposure to aversive details (first responders) - Exposure not from electronic media,	EXPOSURE to trauma (1+) 1. Direct experience 2. Witnessing in person 3. To parent/caregiver - Exposure not from electronic media,	EXPOSURE to trauma (1+) 1. Direct experience 2. Witnessing in person 3. To close family/friend 4. Repeated/extreme exposure to aversive details (first responders) - Exposure not from electronic media,	Emotional/behavioral symptoms Identifiable stressor Within 3 months of stressor onset
		unless work related	unless work related	unless work related	
B SOCIAL/EMOTION DISTURBANCE (2+) 1. Minimal responsiveness to others 2. Limited positive affect 3. Unexplained irritability, sadness, fearfulness (when not threatened)	Not due to impulsivity (ADHD) But includes socially disinhibition	INTRUSION symptoms (1+) 1. Distressing memories 2. Distressing dreams 3. Dissociative rxns (flashbacks) 4. Psychological distress to cues 5. Physiological reactions to cues	INTRUSION symptoms (1+) 1. Distressing memories/play 2. Distressing dreams 3. Dissociative rxns (flashbacks) 4. Psychological distress to cues 5. Physiological reactions to cues	(9+) INTRUSION symptoms Distressing memories/play Distressing dreams Dissociative rxns (flashbacks) Psychological distress or physiological reactions to cues	CLINICALLY SIGNIFICANT (1+) 1. Out of proportion to stressor sever. 2. Sig fxnal impairment
C EXTREMES OF INSUFFICIENT CARE (1+) 1. Social neglect/deprivation 2. Repeated changes of 1° caregivers 3. Rearing in unusual settings	EXTREMES OF INSUFFICIENT CARE (1+) 1. Social neglect/deprivation 2. Repeated changes of 1° caregivers 3. Rearing in unusual settings	AVOIDANCE of stimuli (1+) Distressing memories, thoughts, feelings External reminders arousing memories, thoughts, feelings	(1+) AVOIDANCE of stimuli 1. Distressing memories, thoughts, feelings 2. External reminders arousing memories, thoughts, feelings	or AVOIDANCE of stimuli 1. Distressing memories, thoughts, feelings 2. External reminders arousing memories, thoughts, feelings	
Criterion C responsible for Criteria A	Criterion C responsible for Criteria A	NEGATIVE COGNITION/MOOD (2+) 1. Amnesia about event (typically dissociative) 2. Negative beliefs about oneself, others, world 3. Distorted cognitions about cause/consequences, blame 4. Negative emotional state 5. Diminished interest/activity 6. Detached/estranged feeling 7. Inability for positive emotions	or NEGATIVE COGNITION 1. Negative emotional states 2. Diminished interest/activity 3. Socially withdrawn 4. Reduced positive emotions	or DISSOCIATIVE SYMPTOMS 1. Amnesia about event (typically dissociative) 2. Altered sense of reality of surroundings/oneself or NEGATIVE MOOD 1. Inability for positive emotions	
Not autism spectrum disorder		AROUSAL/REACTIVITY (2+) 1. Irritability, angry outbursts 2. Hypervigilance 3. Exaggerated startle response 4. Concentration problems 5. Sleep disturbance 6. Self-destructive/recklessness	AROUSAL/REACTIVITY (2+) 1. Irritability, angry outbursts 2. Hypervigilance 3. Exaggerated startle response 4. Concentration problems 5. Sleep disturbance	or AROUSAL/REACTIVITY 1. Irritability, angry outbursts 2. Hypervigilance 3. Exaggerated startle response 4. Concentration problems 5. Sleep disturbance	
Disturbance evidence before age 5		Duration >1 month	Duration >1 month	Duration 3 days to 1 month	Not another mental disorder Not exacerbation of mental disorder
Development age >9 months	Development age >9 months	Sig distress/fxnal impairment	Sig distress/fxnal impairment	Sig distress/fxnal impairment	Not normal bereavement
		Not due to substance or AMC	Not due to substance or AMC	Not due to substance or AMC or brief psychotic disorder	Symptoms do not persist >6 mos after stressor/consequences end
Persistent (>12 mo) Severe (all symptoms at high levels)	Persistent (>12 mo) Severe (all symptoms at high levels)	With dissociative symptoms Depersonalization Derealization With delayed expression (>6mo)	With dissociative symptoms Depersonalization Derealization With delayed expression (>6mo)		With depressed mood With anxiety With mixed anxiety and depressed mood With disturbance of conduct With mixed disturbance of emotions and conduct Unspecified

Trauma and Stressor-Related Disorders			
Other Specified	Unspecified		
Do not meet any full criteria	Does not meet any full criteria		
Choose to communication specific	Choose NOT to communication		
reason for not meeting	specific reason for not meeting		
	Insufficient information for dx		
Adjustment-like disorders, delayed			
onset of sx, 3 months after stressor			
Adjustment-like disorders,			
prolonged duration, >6 months,			
without prolonged stressor			
J. T. G. T.			
Ataque de nervios			
Other cultural syndromes			
F. Dominton to consult to the consul			
5. Persistent complex bereavementSevere/persistent grief/mourning			
- Severe/persistent grief/modifiling			