

Clarifying Amazonia's burning crisis

Recent fires in the Amazon have attracted much political and media attention, but it still remains unclear what has been burning. This is crucial to understanding impacts and identifying solutions. There are three broad types of fire in the Amazon (Figure 1a). First, there are deforestation fires—the process of clearing primary forest which starts with the vegetation being felled and left to dry. Fire is then used to prepare the area for agriculture. Second, there are fires in areas that have been previously cleared. For example, cattle ranchers use fire to rid pastures of weeds, and smallholders, indigenous and traditional peoples use fire in farm-fallow systems. Not all fires in previously cleared lands are intentional—some escape beyond

intended limits. Third, fires can invade standing forests, either for the first time when flames are mostly restricted to the understorey, or as repeated events, resulting in more intense fires.

Different types of fire have different drivers (Figure 1a). While weak governance may lead to more deforestation fires (Fonseca et al., 2019), climate change makes forests hotter and drier, thus more likely to sustain uncontrolled fires (Brando et al., 2019). Different fire types also have different impacts. For example, uncontrolled fires in open lands can kill livestock and destroy crops and farm infrastructure, while even low-intensity forest fires can kill up to 50% of the trees and reduce the value of the forests for local




FIGURE 1 (a) An overview of fire types, drivers and their positive feedbacks on fire prevalence. Fire types are shaded by the three broad classes of fire in the Amazon: Deforestation (dark grey), fires on previously cleared lands (grey), and forest fires (light grey). Deforestation and agricultural fires are intentional, while uncontrolled fires are either started accidentally or through malevolent intent. (b) Annual deforestation (August–July, following PRODES from TerraBrasilis & INPE 2019; yellow) and active fire counts (January–August, matching the pre-Moratoria period in 2019; grey bars, stacked to separate January–July and August). Deforestation in 2019 (orange) was estimated based on DETER-b (TerraBrasilis & INPE 2019). Active fire counts are from BDQueimadas. Years with extreme droughts are shown with an asterisk. (c) Monthly deforestation detected by DETER-b for the period April–September, comparing 2019 (red) with the mean \pm SD from 2016–2018 (blue). The vertical dashed line represents the onset of the fire moratoria [Colour figure can be viewed at wileyonlinelibrary.com]

people (e.g. Barlow et al., 2012). In contrast, farm-fallow fires are essential for the food security and livelihoods of some of the Amazon's poorest people (Carmenta, Vermeylen, Parry, & Barlow, 2013). A failure to distinguish between different fire types has contributed to the uncertainty surrounding the recent Amazonian fires, and has important implications for policy responses.

We evaluate the Brazilian government's claims that the Amazon fire situation in August 2019 was 'normal' and 'below the historical average' by assessing the longer term trends in active fires and annual deforestation and recent monthly deforestation trends. The number of active fires in August 2019 was nearly three times higher than in August 2018 and the highest since 2010 (Figure 1b). There is strong evidence this increase in fire was linked to deforestation. To examine this, we first estimate 2019 deforestation, as the numbers from the Brazilian PRODES system for measuring annual deforestation are yet to be published. The area of deforestation detected by PRODES runs from August to July each year and is, on average, 1.54 higher than near-real-time DETER-b measure of deforestation. Using this conversion factor suggests that >10,000 km² of forest was lost in the period between August 2018 and July 2019, which would make it the highest annual loss since 2008 (Figure 1b). These annual trends are mirrored by a sharp increase in monthly deforestation detected by DETER-b—deforestation in July 2019 was almost four times the average from the same period in 2016–2018 (Figure 1c).

The marked upturn in both active fire counts and deforestation (Figure 1b) in 2019 therefore refutes suggestions that August 2019 was a 'normal' fire month in the Amazon (Figure 1b). Moreover, the increase in fires has occurred in the absence of a strong drought, which can be a good predictor of fire occurrence (Aragão et al., 2018). The important contribution of deforestation-related fires was consistent with media footage of large-scale fires in deforested areas, while the enormous plumes of smoke that reached high into the atmosphere can only be explained by the combustion of large amounts of biomass. The unusual nature of 2019 was also emphasized by exceptionally high fire counts in some protected areas, such as Jamanxim National Forest where active fires increased by 355% from 2018 to 2019, 44% above the long-term average (BDQueimadas & INPE, 2019).

Some key uncertainties remain. Despite the large-scale fires seen in August 2019, there was a 35% drop in active fires in September, and it is unclear to what extent rainfall or the recent 2-month fire moratoria declared by President Bolsonaro has contributed to this. Crucially, it is also unclear what will happen now the ban has been lifted. The figures from DETER-b suggest deforestation remained well above average in September, despite the moratoria (Figure 1c). Moreover, over the last 20 years, 65% of annual fire detections occur from September to December, which is the peak of the dry season for much of the Amazon. Given the essential role of fire in clearing felled vegetation, these recently deforested areas are very likely to burn at some point in the future.

Another uncertainty relates to the extent of forest fires. These are rarely detected by active fire counts—for example, active fire

counts during the 2015 El Niño were unexceptional, even though forest fires burned 1 million ha in the Santarém region (Withey et al., 2018) and affected many other areas of the Amazon in one of the strongest El Niño-mediated droughts on record (Jiménez-Muñoz et al., 2016). However, weekly mapping in August 2019 has already revealed 8,500 ha of forest fires in the frontier region of Brazil and Peru (Sonaira Silva, personal communication) despite the lack of an unusual drought in 2019.

Managing Amazonian fires requires understanding what is burning, what drives contagion and extent, and how different drivers combine to make the Amazon more flammable (Figure 1a). Tackling deforestation is key—forest clearance is a major source of ignition, and augments the flammability of remaining forests by increasing edge density, raising regional temperatures and reducing rainfall. Brazil's successful deforestation action plan of 2004–2012 provides a clear blueprint for action (see Figure 1b), but is contrary to the current government's approach of undermining forest monitoring and cutting resources for law enforcement.

Fires on previously cleared lands provide many of the ignition sources for forest fires (Figure 1a). Some of these risks could be reduced by phasing burning out. For example, incentives and capacity building can encourage fire-free cattle ranching, which can also return higher yields than extensive fire-based approaches to pasture management. Finding equitable alternatives to the traditional fire-dependent agriculture practiced by smallholders is more challenging. 'Technological spillover' has enabled some smallholders to access tractors from capitalized landholders, but this could have perverse outcomes for sustainable land use, crop choices that affect regional food security and even social justice. Where alternatives are not practical or socially desirable, policies need to support politically and economically marginalized land users to develop farming practices adapted to changing environmental conditions.

Preventing forest fires will also require action to prevent illegal logging operations, as microclimatic changes make logged forests more flammable (Uhl & Kauffman, 1990). Near-real-time monitoring and forecasting of drought intensity and fire risk would also help, especially if linked to responsive, resourced and capable local fire brigades. Global climate change is also a key driver of change in the Amazonian system, increasing both dry season lengths and temperatures (Brando et al., 2019). Maintaining the climate change mitigation potential of the Amazon is therefore itself dependent on reducing greenhouse gas emissions across the world.

ACKNOWLEDGEMENTS

Some contributors declined authorship to maintain anonymity. We regret this was necessary and thank them for their important contribution. We also thank the editors of the journal for comments that helped improve the manuscript. The work was supported by National Environment Research Council NE/K016431/1, NE/P004512/1 and NE/S0184X/1, PELD (CNPq Process 441659/2016-0), SEM-FLAMA (441949/2018-5) and the Frank Jackson Foundation.

Jos Barlow^{1,2} 

Erika Berenguer^{1,3} 

Rachel Carmenta⁴ 

Filipe França⁵ 

¹Lancaster Environment Centre, Lancaster University, Lancaster, UK

²Universidade Federal de Lavras, Lavras, Brazil

³Environmental Change Institute, University of Oxford, Oxford, UK

⁴Department of Geography, University of Cambridge Conservation Research Institute, University of Cambridge, Cambridge, UK

⁵Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, PA, Brazil

Correspondence

Jos Barlow, Lancaster Environment Centre, Lancaster University,
Lancaster LA1 4YQ, UK.

Email: jos.barlow@lancaster.ac.uk

The full article is available in Portuguese:
see Supporting Information.

ORCID

Jos Barlow  <https://orcid.org/0000-0003-4992-2594>

Erika Berenguer  <https://orcid.org/0000-0001-8157-8792>

Rachel Carmenta  <https://orcid.org/0000-0001-8607-4147>

Filipe França  <http://orcid.org/0000-0003-3827-1917>

REFERENCES

Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G. M. G., Rosan, T. M. T. M., Vedovato, L. B. L. B., Wagner, F. H. F. H., ... Saatchi, S. (2018).

21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nature Communications*, 9(1), 1–12. <https://doi.org/10.1038/s41467-017-02771-y>

Barlow, J., Parry, L., Gardner, T. A., Ferreira, J., Aragão, L. E. O. C., Carmenta, R., ... Cochrane, M. A. (2012). The critical importance of considering fire in REDD+ programs. *Biological Conservation*, 154, 1–8. <https://doi.org/10.1016/j.biocon.2012.03.034>

BDQueimadas & INPE. (2019). *Banco de Dados de Queimadas. INPE - Programa Queimadas*. Retrieved from <http://queimadas.dgi.inpe.br/queimadas/bdqueimadas>

Brando, P. M., Paolucci, L., Ummenhofer, C. C., Ordway, E. M., Hartmann, H., Cattau, M. E., ... Balch, J. (2019). Droughts, wildfires, and forest carbon cycling: A pantropical synthesis. *Annual Review of Earth and Planetary Sciences*, 47(1), 555–581. <https://doi.org/10.1146/annurev-earth-082517-010235>

Carmenta, R., Vermeylen, S., Parry, L., & Barlow, J. (2013). Shifting cultivation and fire policy: Insights from the Brazilian Amazon. *Human Ecology*, 41(4), 603–614. <https://doi.org/10.1007/s10745-013-9600-1>

Fonseca, M. G., Alves, L. M., Aguiar, A. P. D., Arai, E., Anderson, L. O., Rosan, T. M., ... Aragão, L. E. O. e. C. (2019). Effects of climate and land-use change scenarios on fire probability during the 21st century in the Brazilian Amazon. *Global Change Biology*, 25(9), 2931–2946. <https://doi.org/10.1111/gcb.14709>

Jiménez-Muñoz, J. C., Mattar, C., Barichivich, J., Santamaría-Artigas, A., Takahashi, K., Malhi, Y., ... Schrier, G. V. D. (2016). Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–2016. *Scientific Reports*, 6, 33130. <https://doi.org/10.1038/srep33130>

TerraBrasilis, and INPE. (2019). *TerraBrasilis. PRODES (Desmatamento)*. Retrieved from <http://terrabrasilis.dpi.inpe.br/en/home-page/>

Uhl, C., & Kauffman, J. B. (1990). Deforestation, fire susceptibility, and potential tree responses to fire in the Eastern Amazon. *Ecology*, 71(2), 437–449. <https://doi.org/10.2307/1940299>

Withey, K., Berenguer, E., Palmeira, A. F., Espírito-Santo, F. D. B., Lennox, G. D., Silva, C. V. J., ... Barlow, J. (2018). Quantifying immediate carbon emissions from El Niño-mediated wildfires in humid tropical forests. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170312. <https://doi.org/10.1098/rstb.2017.0312>