

SPIROMETRY REQUISITION

INSPIRE Lung and Medical Centre

Unit 302 - 9639 137A St, Surrey, BC, V3T0M1

Phone: 604 676 7750 Fax: 604 676 7751 Website: www.inspireImc.com

Date:	Patient Name:			
Referring MD:	PHN: Cell:			
Phone:	DOB: Home:			
Fax:	Other: Work:			
Signature:	Email:			
Reason for Test: Diagnosis Follow-Up Other: Clinical Diagnosis: Smoking History: Never Former Current Pack years #: Respiratory Medications: Spirometry requested: Pre-Bronchodilator only Pre and Post-Bronchodilator (400ug salbutamol given) Withhold medications? Yes (see page 2) No	Relative Contraindications: ☐ Acute MI within the past week ☐ Systemic hypotension or severe hypertension ☐ Significant arrythmia ☐ Uncompensated heart failure ☐ Uncontrolled pulmonary hypertension ☐ Acute cor pulmonale ☐ Unstable pulmonary embolism ☐ Syncope related to cough / forced expiration ☐ Cerebral aneurism ☐ Brain surgery < 4 weeks ☐ Recent symptomatic concussion ☐ Eye surgery < 1 week ☐ Sinus surgery, middle ear surgery, or infection < 1 week ☐ Pneumothorax ☐ Thoracic surgery < 4 weeks ☐ Abdominal surgery within 4 weeks ☐ Late term pregnancy ☐ Active or suspected transmissible respiratory or systemic infection, including TB ☐ Hemoptysis			
List medications and withholding time:	Significant secretions, oral lesions, or oral bleeding			

OFFICE USE ONLY Appointment Date:

Appointment Time:

PLEASE PROVIDE THE FOLLOWING INSTRUCTIONS TO THE PATIENT

• Review detailed instructions on the website: www.inspirelmc.com

Medications to withhold, and for now long:						

Bronchodilator Medication	Withholding Time		
SABA (eg: salbutamol)	4-6h		
SAMA (eg: ipratropium bromide)	12h		
LABA (eg: formoterol, salmeterol)	24h		
Ultra-LABA (eg: indacaterol, vilanterol, olodaterol)	36h		
LAMA (eg: tiotropium, umeclidinium, aclidinium, glycopyrronium)	36-48h		

• If needed for symptom relief, a rescue inhaler should be used, and the time of use noted so that it can be reported to the technologist conducting the test

Avoid the following prior to testing:

- Smoking and/or vaping and/or water pipe within 1 hour before testing
- Consuming intoxicants within 8 hours before testing
- · Consuming alcohol or caffeine within 4 hours before testing
- Eating a large meal within 2 hours before testing
- · Performing vigorous exercise within 1 hour before testing
- Wearing clothing that restricts full chest and abdominal expansion