

Positive Double Blind Clinical Low Level Laser Therapy (LLLT) Studies

(Summary according to category)

Allergic rhinitis	1
Arthritis	16*
Epicondylitis	5*
Fibrositis	1
Herpes simplex	1
Hypersensitive dentine	3
Microcirculation	1
Mucositis	1
Nerve functions	6*
Pain	33
Paresthesia	2*
Post herpetic neuralgia	2*
Tendinitis	3*
Trigger points	4*
Sinusitis	1
Sjogren's Syndrome	1
Wound healing	8*

* Multiple positive double blind Low Level Laser Therapy (LLLT) studies performed

Airaksinen O., et al.

Effects of infra-red laser irradiation at the trigger points. Scand J of Acu & EI Therapy. 1988; 3: 56-61

Airaksinen O., et al.

Effects of laser irradiation at the treated and non-treated trigger points. Proc. 4th Intern Symposium. Acupunc & Electrother Res. 1988; 13 (4): 238-239.

Antipa C. et al.

Clinical results of the low energy laser action on distal forearm posttraumatic nerve lesions. Laser Therapy. 1996; 1: 36.

Antipa C. et al.

Comparative effects of various IR low energy diodes in the treatment of the rheumatic diseases. 1997. In press (Monduzzi Editore, Bologna)

Armino L. et al.

Laser therapy in post-episiotomic neuralgie. LASER. Journ Eur Med Laser Ass. 1988; 1(1):7.

Atsumi K. et al.

Biostimulation effect of low-power energy diode laser for pain relief.
Lasers Surg Med. 1987; 7: 77.

Barabas K. et al.

Controlled clinical and experimental examinations on rheumatoid arthritis patients and synovial membranes performed with neodym phosphate glas laser irradiation.
Proc. 7th Congr Internat Soc for Laser Surg and Med, Munich June 1987. Abstract no 216a.

Bihari I., Mester A.

The biostimulative effect of low level laser therapy of long-standing crural ulcer using Helium Neon laser, Helium Neon plus infrared lasers and non coherent light: Preliminary report of a randomized double blind comparative study.
Laser Therapy. 1989; 1(2): 97.

Boerner E. et al.

Double-blind study on the efficacy of the laser therapy.
SPIE Proc. 1996. Vol. 2929: 75-79.

Bihari I., Mester A.

The biostimulative effect of low level laser therapy of long-standing crural ulcer using Helium Neon laser, Helium Neon plus infrared lasers and non coherent light: Preliminary report of a randomized double blind comparative study.
Laser Therapy. 1989; 1(2): 97.

Carillo J. et al.

A randomized double-blind clinical trial on the effectiveness of helium-neon laser in the prevention of pain, swelling and trismus after removal of impacted third molars.
Int Dent Journ. 1990; 40: 31.

Ceccherelli F.

Diode laser in cervical myofascial pain. A double-blind study versus placebo.
The Clin J Pain. 1989; 4: 301-304.

Cheng R.

Combined treatments of electrotherapy plus soft laser therapy has synergistic effect in pain relief and disease healing. Surgical and Medical Lasers.
1990; 3 (3): 135 (abstract).

Cieslar G. et al.

Effect of low-power laser radiation in the treatment of the motional system overloading syndromes.
SPIE Proc. Vol 3198. 1997, pp. 76-82.

Cowen D. et al.

Low energy helium neon laser in the prevention of oral mucositis in patients undergoing bone marrow transplant: results of a double blind randomized trial.
Int J Radiat Oncol Biol Phys. 1997; 38 (4): 697-703.

de Bie R. A.

Effect of laser therapy on ankle sprains
Ned. T. Fysiotherapie. 1988; 95: 108-112. (in Dutch)

Eckerdal A., Lehmann Bastian H.

Can low reactive-level laser therapy be used in the treatment of neurogenic facial pain? A double-blind, placebo controlled investigation of patients with trigeminal neuralgia.
Laser Therapy. 1996; 8: 247-252.

Emmanoulidis O. et al.

CW IR low-power laser application significantly accelerates chronic pain relief rehabilitation of professional athletes. A double blind study.
Lasers Surg Med. 1986; 6: 173.

England S. et al.

Low power laser therapy of shoulder tendinitis.
Scand J Rheumatology. 1989; 18: 427.

Flöter T., Refisch H. P.

Pain treatment with laser. A double blind study. Proc. of the 4th Internat Symposium. Acupunct & Electro-Therap Res. 1988; 13(4): 236-237. Also: Schmerzbehandlung mit Laser. Eine Doppelblinde Studie. Top Medizin. 1990; 4(4): 52-56.

Fructuoso F. J. G., Moset J. M.

Estudio randomizado doble ciego sobre los efectos bioestimulantes del láser en la irradiación de glándula paratida en pacientes afectados de síndrome de Sjogren. (Double blind study on the biostimulatory effects of laser irradiation on the parotid gland in patients affected by Sjogrens syndrome).
Investigacion y Clinica Laser. 1987; 4 (1): 18-25.

Gelskey S. C. et al.

The effectiveness of the Nd:YAG laser in the treatment of dentinal hypersensitivity.
J Can Dent Assoc. 1993; 59 (4): 337-386.

Gerschman J. A. et al.

Low Level Laser in dentin hypersensitivity.
Australian Dent J. 1994;39:6.

Goldman J. A. et al.

Laser therapy of rheumatoid arthritis.
Lasers Surg Med. 1980; 1: 93-102.

Gudmundsen J. et al.

Laserbehandling av epicondylitis humeri og rotatorcuff-syndrom. Dobbelt blindstudie - 200 pasienter. (Laser treatment of epicondylitis humeri and rotator cuff syndrome. Double blind study - 200 patients. In Norwegian)
Norsk tidsskrift for idrettsmedisin. 1987; 2: 6.

Gertner C.

Analgesy by low power laser (LPL): a controlled double blind study in ankylosing spondarthritis (SPA).

Lasers Surg Med. 1989; Suppl 1:55.

Gärtte S. et al.

Doppelblindstudie zur Ueberpruefung der Wirksamkeit und Vertraeglichkeit einer niederenergetischen Lasertherapie bei Patienten mit aktiver Gonarthrose.

Jaros Orthopaedie. 1995. 12: 3034.

Haker E. et al.

Is low-energy laser treatment effective in lateral epicondylalgia? J of Pain and Symptom Management.

1991; 6(4): 241.

Hashimoto K.

Clinical applications of various lasers in oral surgery.

Lasers in dentistry. Eds. Yamamoto Y et al. 1989; p. 63-70. Elsevier Science Publishing B.V, Amsterdam

Hashimoto T. et al.

Efficacy of laser irradiation on the area near the stellate ganglion is dose-dependent: a double-blind crossover placebo-controlled study.

Laser Therapy. 1997; 1(9): 7-11.

Hopkins G. O. et al.

Double blind cross over study of laser versus placebo in the treatment of tennis elbow. Proc International. Congress on Lasers, "Laser Bologna".

1985: 210. Monduzzi Editore S.p.A., Bologna.

Hoteya K. et al.

Effects of a 1 W GaAlAs diode laser in the field of orthopedics. In: Meeting Report: The first Congress of the International Association for Laser and Sports Medicine.

Tokyo, 1997. Laser Therapy 1997; 9 (4): 185.

Kaiser C. et al.

Estudio en doble ciego randomizado sobre la eficacia del He-Ne en el tratamiento de la sinuitis maxilar aguda: en pacientes con exacerbacion de una infeccion sinusal cronica.

(Double blind randomized study on the effect of HeNe in the treatment of acute maxillary sinusitis: in patients with exacerbation of a chronic maxillary sinusitis).

Boleton CDL. 1986; 9: 15.

Kamikawa K. et al.

Double blind experiences with mid-Lasers in Japan. 1985. Proc Int Congr on Lasers, "Laser Bologna".

1985: 165-169. Monduzzi Editore S.p.A., Bologna.

Kemmotsu M. D. et al.

LLLT for pain attenuation - the current experience in the pain clinic. In: Progress in Laser Therapy.

Eds Oshiro T, Calderhead R G. 1991: 197-200. John Wiley & Sons, Chichester, Engl. ISBN 0-471-93154-3.

Khullar S. M. et al.

Low level laser treatment improves longstanding sensory aberrations in the inferior alveolar nerve following surgical trauma.

J Oral Maxillofac Surg. 1996; 54: 2-7.

Khullar S. M. et al.

Effect of low-level laser treatment on neurosensory deficits subsequent to sagittal split ramus osteotomy.

Oral Surgery Oral Medicine Oral Pathology. 1966; 82 (2): 132-8.

Kim J. W., Lee J. O.

Double blind cross-over clinical study of 830 nm diode laser and 5 years clinical experience of biostimulation in plastic & aesthetic surgery in Asians.

Lasers Surg Med. 1998; Suppl. 10: 59.

Kinoshita F. et al.

Clinical evaluation of low-energy, semi-conductor laser therapy in oral surgery - a double blind study.

Josai Shika Daigaku Kiyo. 1986; 15 (3): 735-742. (in Japanese)

Laakso E. L. et al.

Pain scores and side effects in response to low level laser therapy (LLLT) for myofascial trigger points.

Laser Therapy. 1997; 2 (9): 67-72.

Lonauer G.

Controlled double blind study on the efficacy of He-Ne-laser beams versus He-Ne- plus Infrared-laser beams in the therapy of activated osteoarthritis of finger joints.

Clin Experim Rheuma. 1987; 5 (suppl 2) : 39

Longo L. et al.

Treatment with 904 nm and 10600 nm laser of acute lumbago - double blind control.

LASER. Journ Eur Med Laser Ass. 1988; 1(3):16.

Lucas C. et al.

Low level laser therapy bij decubitus statium III.

Rapport Hoegschool van Amsterdam. 1994.

Loegdberg-Andersson M. et al.

Low level laser therapy (LLLT) of tendinitis and myofascial pains - a randomized, double-blind, controlled study.

Laser Therapy. 1997; 2 (9): 79-86.

Mach E. S. et al.

Helium-Neon (Red Light) Therapy of Arthritis.
Rheumatologia, 1983; 3: 36.

Meier J. .L, Kerkour K.

Traitement laser de la tendinite.
Med. et Hyg. 1989; 46: 907-911.

Mester A.

Biostimulative effect in wound healing by continuous wave 820 nm laser diode. Double-blind randomized cross-over study.
Lasers in Med Science, abstract issue July 1988, No 289.

Miyagi K.

Double-blind comparative study of the effect of low-energy laser irradiation to rheumatoid arthritis.
Current awareness of Excerpta Medica. Amsterdam. Elsevier Science Publishers BV. 1989; 25: 315.

Mokhtar B. et al.

A double blind placebo controlled investigation of the hypoalgesic effects of low intensity laser irradiation of the cervical roots using experimental ischaemic pain. Proc. Second Meeting of the International Laser Therapy Assn., ôLondon Laserö, Sept 1992, p 61.

Mokhtar B. et al.

The possible significance of pulse repetition rate in lasermediated analgesia: A double blind placebo controlled investigation using experimental ischaemic pain. Proc. Second Meeting of the International Laser Therapy Assn, ôLondon Laserö Sept 1992. p 62

Molina J. J. et al.

La laserterapia como coadyuvante en el tratamiento de la A.R. (Artritis Reumatoidea). Boletin C.D.L., Barcelona. 1987; 14: 4-8.

Moore K. et al.

LLLT treatment of post herpetic neuralgia.
Laser Therapy. 1988; 1: 7.

Moore K. et al

The effect of infra-red diode laser irradiation on the duration and severity of postoperative pain. A double-blind trial.
Laser Therapy. 1992; 4: 145.

Mousques T.

etude en double aveugle des effets du traitement unilateral au laser helium-neon lors de chirurgies parodontales bilaterales simultanes.
Quest Odonto-Stomatol. 1986; 11: 245.

Mousques T.

Etude en double aveugle des effets du helium-neon en chirurgie parodontale.
Quest Odonto-Stomatol 1986; 11: 223.

Neuman I. et al.

Low energy phototherapy in allergic rhinitis and nasal polyposis.
Laser Therapy. 1996. 1: 37.

Nivbrant Bo et al.

Therapeutic laser treatment in gonarthrosis.
Acta Orthop Scand. 1989; 60: 231.

Ortutay J et al.

Psoriatic Arthritis Treatment with low power laser irradiation. A double blind clinical study.
Lasermedizin - Laser in Med Surg. 1998; 13 (3-4): 140.

Oyamada Y. et al.

A double blind study of low power He-Ne laser therapy in rheumatoid arthritis.
Optoelectronics in Medicine. 1987; p 747-750. Springer Verlag, Berlin (abstract). Complete study in Boletón de CDL. 1988; 17: 8-12.

Palmgren N. et al.

Low-Power Laser Therapy in Rheumatoid Arthritis.
Lasers in Medical Science. 1989; 4: 193.

Palmgren N. et al.

Low Level Laser Therapy of infected abdominal wounds after surgery.
Lasers Surg Med. 1991; Suppl 3:11.

Palmieri B.

A double blind stratified cross over study of amateur tennis players suffering from tennis elbow using infrared laser therapy.
Medical Laser Report. 1984; 1: 3-14

Rochkind S. et al.

Double-blind Randomized Study Using Neurotube and Laser Therapy in the Treatment of Complete Sciatic Nerve Injury of Rats.
Proc. 2nd Congr World Assoc. for Laser Therapy, Kansas City, 1998.

Roumeliotis D. et al.

820nm 15mW 4J/cm², laser diode application in sports injuries. A double blind study.
Proc. Fifth Annual Congress British Medical Laser Association. 1987.

Saeki N. et al.

Double blind test for biostimulation effects on pain relief by diode laser.
1989. Laser Surgery; 1066: 93-100.

Sasaki K. et al.

A double-blind controlled study on free amino acid analysis in CO₂ laser burn wounds in the mouse model following doses of low incident infrared (830 nm) diode laser energy.
Proc. 2nd Meeting of the Internat Laser Therapy Assn., London, 1992, p.4.

Sasaki K. et al.

A preliminary double blind controlled study on free amino acid analysis in burn wounds in the mouse following 830 nm diode laser therapy.
Laser Therapy. 1997; 2 (9): 59-65.

Sato K. et al.

A double blind assessment of low power laser therapy in the treatment of postherpetic neuralgia. Surgical and Medical Lasers.
1990; 3 (3): 134 (abstract)

Saunders L.

The efficiency of low-level laser therapy in supraspinatus tendinitis.
Clin Rehab. 1995; 9: 126-134

Schindl A. et al.

Low intensity laser irradiation improves skin circulation in patients with diabetic microangiopathy.
Lasers Surg Med. 1998; Suppl. 10: 7.

Scudds R. A. et al:

A double-blind crossover study of the effects of low-power gallium arsenide laser on the symptoms of fibrositis.
Physiotherapy Canada. 1989; 41: (suppl 3): 2.

Simunovic Z., Trobonjaca T. et al.

Treatment of medial and lateral epicondylitis - tennis and golfer elbow - with low level laser therapy: a multicenter double blind, placebo controlled clinical study on 324 patients.
J Clin Laser Med & Surg. 1998; 16 (3): 145-151.

Simunovic Z., Trobonjaca T.

Soft tissue injury during sport activities and traffic accidents - treatment with low level laser therapy. A multicenter double blind, placebo controlled clinical study on 132 patients.
Proc. IXX ASLMS Congress, Orlando, Florida, April 1999.

Snyder-Mackler L. et al.

Effect of helium-neon laser on musculoskeletal trigger points.
Physical Therapy. 1986; 66: 1087.

Snyder-Mackler L. et al.

Effect of helium-neon laser irradiation on peripheral sensory nerve latency.
Physical Therapy. 1988; 68: 223.

Snyder-Mackler L. et al.

Effect of helium-neon laser irradiation on skin resistance and pain in patients with trigger points in the neck or back
Physical Therapy. 1989; 69: 336.

Soriano F. A. et al

Acute cervical pain is relieved with gallium-arsenida (GaAs) laser irradiation. A double-blind preliminary study.
Laser Therapy. 1996; 8: 149-154.

Soriano F. A. et al.

Low level laser therapy response in patients with chronic low back pain. A double blind study.
Lasers Surg Med. 1998, Suppl. 10, p. 6.

Toya S. et al.

Report on a computer-randomized double blind clinical trial to determine the effectiveness of the GaAlAs (830 nm) diode laser for pain attenuation in selected pain.
Laser Therapy 1994; 6:143.

Taguchi T. et al.

Thermographic changes following laser irradiation for pain.
Clinical Laser Med Surg. 1991; 2(9): 143.

Tsurko V. et al.

Laser therapy of rheumatoid arthritis. A clinical and morphological study.
Terap Arkh. 1983; 97. (Russian).

Volez-Gonzalez M. et al.

Treatment of relapse in herpes simplex on labial and facial areas and of primary herpes simplex on genital areas and "area pudenda" with low power HeNe-laser or Acyclovir administered orally.
SPIE Proc. 1995; Vol. 2630: 43-50

Vasseljen O. et al.

Low level laser versus placebo in the treatment of tennis elbow. Scand Scand J Rehab Med. 1992; 24: 37. Also in Physiotherapy. 1992; 5: 329.

Walker J.

Relief from Chronic Pain by Low Power Laser Irradiation.
Neuroscience Letters. 1983; 43: 339

Walker J.

Temporary suppression of clonus in humans by brief photostimulation
Brain Research. 1985; 340: 109.

Walsh D. et al.

The effect of low intensity laser irradiation upon conduction and skin temperature in the superficial radial nerve. Double-blind placebo controlled investigation using experimental ischaemic pain.

Proc. Second Meeting of the Internat Laser Therapy Association, London, sept. 1992.

Willner R. et al.

Low power infrared laser biostimulation of chronic osteoarthritis in hand.

Lasers Surg Med. 1985; 5: 149.

Wylie L. et al.

The hypoalgesic effects of low intensity infrared laser therapy upon mechanical pain threshold.

Lasers Surg Med. 1995; Suppl 7:9.

Yamaguchi M. et al.

Clinical study on the treatment of hypersensitive dentin by GaAlAs laser diode using the double blind test.

Aichi Gakuin Daigaku Shigakkai Shi - Aichi-Gakuin Journal of Dental Science. 1990; 28(2): 703-707. (in Japanese)