



Kelpie Wilson
Wilson Biochar Associates
www.wilsonbiochar.com
kelpiew@gmail.com
Office: 541-592-3083
Mobile: 541-218-9890

Biochar Return on Investment in Fruit and Nut Orchard Production

By Kelpie Wilson

Biochar is a form of highly stable carbon produced by the thermal degradation of biomass that creates value when used as a soil amendment. Biochar also provides climate benefits by transferring carbon from the atmosphere to highly stable soil carbon pools. As a soil amendment, biochar can address production bottlenecks while stimulating higher yields of an improved product that fetches a premium price. Farmers can incorporate biochar into current cultural practices for the production of walnuts, almonds, filberts, avocados, citrus, stone fruits, olives, apples, pears, pomegranates and other tree crops at a small additional cost, with variations in application methods depending on the irrigation system used. This paper discusses common nut and fruit production bottlenecks and how biochar can address these. It also proposes a method of partial budget analysis for determining whether biochar is economically viable as a soil amendment for orchards.

How Biochar Addresses Fruit and Nut Orchard Production Bottlenecks

Propagation and disease control

Control of tree crop pathogens begins in the nursery where rootstocks and cuttings are propagated. Biochar has been shown to induce resistance to a variety of pathogens and to encourage strong root growth in many different horticultural crops¹. Activated carbon (with many similar properties to biochar) has long been used in tissue culture to encourage growth and root vigor². Recommended formulations for grafting wax use powdered wood charcoal to protect against disease³.

Water management – physical effects

Biochar increases soil water holding capacity (WHC), reducing irrigation needs. In one sandy loam, biochar increased WHC by 23%⁴. Many researchers have found similar results in a variety of soil and biochar combinations and have shown that the effect on WHC increases linearly with biochar application rate⁵. Biochar can increase total field capacity for better retention of both irrigation water and winter and spring rainfall.

Water management – biological effects

Biochar buffers soil water availability and the flooding/drying cycles that cause problems with fungal diseases such as *Phytophthora*. Evidence shows that biochar can make water more available to plants under water stress conditions and increase water use efficiency (WUE)^{6,7}. This is an important benefit for young trees as they are especially sensitive to

water stress. Declining snow packs in the West are forcing land managers to consider deficit watering strategies for some crops, but this approach leads to an increased potential for problems such as reduced fruit size, greater pest problems and decreased tree health⁸. Biochar-mediated improvements in WHC and WUE can help make the most of scarce water resources. Maintaining consistent soil water can also help avoid calcium uptake fluctuations that cause fruit diseases like apple bitter pit. Water availability buffering also benefits nutrient cycling by maintaining more consistent nutrient levels. This can improve yields with less fertilizer. Researchers in California are recommending biochar for increasing yields of avocado, especially under drought conditions⁹.

Nutrient leaching and salinization

Biochar has been shown to prevent leaching of nitrogen into groundwater in diverse cropping systems, including orchards^{10, 11, 12}. As a result, producers can safely apply less fertilizer and they will avoid pollution problems in the face of proposed surcharges on nitrogen fertilizers. Cutting back fertilizer use will also reduce fertilizer salt accumulation in soil.

Fruit quality, size and yield

Biochar applications can result in bigger, healthier fruits and nuts. A five-year biochar trial of apples in Australia has produced fruits that are up to 20 grams heavier than fruits from untreated trees, with no decrease in fruit firmness or sugar content¹³. In a Costa Rican trial, using cacao, researchers observed that biochar alone increased the fresh weight production of cacao fruits (pods) per tree and per hectare and also increased the height and stem diameter of seedlings after planting. Biochar reduced the incidence of *Monilia* pod rot.¹⁴ Biochar can also lead to earlier production in new orchards. Field trials in Belize have seen a 2.5-year-old cacao tree producing fruits 18 months after a top-dressing of biochar. Most cacao orchards in the area do not produce fruit until trees are 5-7 years old¹⁵.

Trees, biochar and vesicular-arbuscular mycorrhizae

Vesicular-arbuscular mycorrhizae (VAM) are especially important for tree crops, promoting significant increases in plant productivity and survival¹⁶. Biochar supports the growth of healthy communities of beneficial microbes such as VAM that enhance soil nutrient cycling and plant nutrient uptake¹⁷. UC Davis researchers studying biochar use in a walnut orchard have found that populations of soil microorganisms increased in the biochar amended plots¹⁸. In a Japanese study, two year-old citrus trees in root boxes were used to evaluate three types of biochar as VAM inoculant carriers. All three biochars showed increased VAM infection along with increased root and shoot growth in the trees (at least 150% greater than an unamended control)¹⁹. One of the biochar types used also produced significantly higher phosphorus concentration in leaves. A second experiment by the same researchers applied biochar in trenches around 15 year-old trees in an established orchard. Root samples taken a year later found a much higher percentage of VAM in biochar-amended trees as compared to controls. In general, beneficial microbial inoculants are likely to be better supported and provide benefits over a longer period in biochar-amended soils^{20,21}.

Disease and pest resistance

Biochar has been shown to induce plant resistance to a broad spectrum of pathogens and parasites, including bacteria, viruses, molds, mildews, soil-borne fungal pathogens, mites, flies and nematodes. This induced resistance is partly due to biochar promoting the growth of beneficial fungi and bacteria that outcompete soil-borne pathogens, and partly due to induced systemic resistance in plants, possibly from increased populations of plant-colonizing microbes. There may be abiotic resistance mechanisms as well, related to organic chemical compounds present in biochar²². Two studies have demonstrated the effectiveness of biochar in suppressing bacterial wilt (*Ralstonia solanacearum*, *Pseudomonas syringae*) in tomatoes^{23,24}. One of these studies also reported a decrease in whitefly populations on greenhouse tomatoes grown in biochar-amended soil that was inoculated with *Trichoderma harzianum* and *Bacillus subtilis*, two beneficial microbes. A study of three different strawberry fungal diseases with three different infection mechanisms found that biochar was effective in suppressing all three (Grey mold caused by *Botrytis cinerea*, Strawberry black spot caused by *Colletotrichum acutatum* and Strawberry powdery mildew caused by *Podosphaera aphanis*)²⁵. An experiment to assess resistance to *Phytophthora cinnamomi* in oak and maple seedlings found that not only did biochar induce systemic resistance to canker expansion; it also reduced water stress in infected plants²⁶. Another study measured parasitic nematode abundance in a wheat field and found that biochar altered the species composition of nematodes (without reducing overall nematode abundance) toward more beneficial species and fewer plant parasites²⁷.

Plant establishment

Biochar-mediated improvement in WHC helps to maintain the higher soil moisture levels that are needed for adventitious root development and successful plant establishment, while avoiding waterlogged conditions that promote fungal infections. Biochar can also protect sensitive young trees from damage related to residual herbicides in the soil. Root dips of activated carbon have been recommended to protect young plants from herbicide toxicity since the 1950s²⁸. Biochar can be added to other amendments and fertilizers that are placed directly in tree planting holes²⁹.

Revitalizing older orchards

Some tree diseases of older orchards may respond to a biochar soil revitalization treatment. Fungal pathogens of walnuts, almonds and stone fruits in California are developing resistance to fungicides. Root aging and browning are caused by warm, dry soils and the presence of phenolic compounds. Biochar can improve soil moisture and it adsorbs phenolic compounds. If the orchard has a re-development budget, it will likely be cost effective to utilize techniques developed in Japan to successfully restore root systems of declining trees by adding biochar, phosphorus and VAM inoculant to holes or trenches around a tree's drip line³⁰.

Replant preparation

When orchards are replaced, producers are often faced with replant rejection due to an "unfriendly" soil ecosystem caused by high levels of soil pathogens that were tolerable to mature trees, but overwhelming to saplings. To restore a more balanced soil microbial environment, extension agents recommend fallowing with a cover crop for a year or more,

but this may not be economical. The alternative of soil fumigation is also problematic. Methyl bromide is a Class I stratospheric ozone-depleting chemical under the Montreal Protocol, and fumigation with methyl bromide is scheduled for phase out in California³¹. Other fumigation chemicals come with water quality and health concerns. Researchers are looking for ways to quickly re-establish a friendly microbial community that can induce systemic resistance to disease³². Two recent studies confirm the potential of biochar soil amendments to mitigate replant disease. Using replant soil from an apple orchard, researchers found that biochar adsorbed phenolic acids in the soil and increased plant height, fresh weight, and several measures of photosynthesis while promoting higher enzyme activity³³. A study of disease-susceptible peach rootstock in replant soil found that total biomass production was significantly higher with biochar³⁴.

Interactions with herbicides and pesticides

Soil organic matter in all its forms is known to adsorb soil-applied herbicides³⁵. Depending on the amount of biochar in soil and its adsorption characteristics, biochar may also reduce the effectiveness of soil-applied herbicides^{36,37,38,39}. Those biochars with less surface area will adsorb herbicides less strongly. Researchers have found that soil applied herbicides were still effective at killing weeds in biochar-amended soil, but only at the higher label application rates. Once available sorption sites on biochar are filled, biochar-amended soils behave like control soils in herbicide adsorption⁴⁰. Biochar can also beneficially reduce leaching of both soil and foliar applied herbicides and pesticides, retaining them in place where they degrade over time⁴¹.

How Biochar Produces Higher Yields of Fruits and Nuts

Higher Yields of Marketable Fruits and Nuts

Increased resistance to pests and diseases combined with better use of nutrients and water can result in greater yields, as shown by the results cited previously for apples and cacao.

Earlier and Longer Production

Orchards require a considerable investment to establish. Producers cannot afford to let poor establishment, early decline or disease reduce their returns. Biochar can produce stronger transplants that can be moved to the field earlier and survive better. Biochar may also bring on high yield production at an earlier date. Biochar can help to maintain vigor and productivity in mature and older orchards, while biochar soil improvement can make replanting more efficient as well. Longer total production time results in a reduced unit cost of production and an improved return on investment.

Biochar Impacts on Economic Returns

In order to determine the return on investment for biochar, a partial budget analysis may be used. A partial budget analysis distributes costs and benefits into four quadrants that account for additional costs and returns as well as reduced costs and returns. The net return is calculated by combining the results from each quadrant. Below are some items

that should be considered when evaluating the return on investment from adding biochar to orchard soil.

Additional Costs

- Cost of biochar – Biochar can be purchased from suppliers or it can be produced on site using orchard prunings. Several low cost methods are available to produce biochar from material that might normally be burned on site to dispose of it⁴².
- Application amount - In some cropping systems biochar can be applied in small annual increments until the desired soil saturation is reached. For tree crops it will be most effective to apply all the biochar at one time in the tree row with the cost spread over the orchard's lifetime. A small additional amount of biochar is applied in the planting holes.
- Application cost - Biochar will be applied along with other soil amendments during regular land preparation, or it can be added to planting holes when replacing trees in an older orchard. It can also be added to compost or other amendments that are surface applied in established orchards
- Propagation – adding biochar to the propagation system will incur a small additional cost for trees.

Reduced Costs

- Biochar may result in significant water use reduction based on a combination of increased water holding capacity and increased plant water availability and water use efficiency.
- Biochar may reduce the amount of NPK fertilizers required based on biochar impacts on soil nutrient cycling and retention.
- Biochar may reduce losses to disease.

Additional Returns

- Yield increase – To start with, it is reasonable to assume a yield increase of 15% based on average yield increases reported for a number of crops using biochar⁴³. Actual yield increases will depend on a number of factors, including the type of crop and soil type as well as disease pressure, climate, fertilization regime and many other factors.

Reduced Returns

- No reductions in returns are anticipated from properly applied biochar.

Conclusions

This economic model for tree crop production using biochar is based on reasonable assumptions. Biochar is new to modern agriculture, but it has a long history of use in many cultures, including 19th century farming in Europe and the United States⁴⁴. This long history of use gives assurance that biochar will provide a net benefit when added to farm soils. Actual economic returns and assumptions about specific costs and benefits will need to be proven through extensive, well-documented field trials.

-
- ¹ Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., & Crowley, D. (2011). Biochar effects on soil biota—a review. *Soil Biology and Biochemistry*, 43(9), 1812-1836.
- ² Thomas, T. D. (2008). The role of activated charcoal in plant tissue culture. *Biotechnology advances*, 26(6), 618-631.
- ³ Oberly, G. H. (1975). Top-Working and Bridge-Grafting Fruit Trees. Cornell Cooperative Extension Publication. Information Bulletin 75.
- ⁴ Basso, A. S., Miguez, F. E., Laird, D. A., Horton, R., & Westgate, M. (2012). Assessing potential of biochar for increasing water-holding capacity of sandy soils. *GCB Bioenergy*.
- ⁵ Briggs, C., Breiner, J. M., & Graham, R. C. (2012). Physical and chemical properties of *Pinus ponderosa* charcoal: implications for soil modification. *Soil Science*, 177(4), 263.
- ⁶ Pereira, R. G., Heinemann, A. B., Madari, B. E., Carvalho, M. T. D. M., Kliemann, H. J., & Santos, A. P. D. (2012). Transpiration response of upland rice to water deficit changed by different levels of eucalyptus biochar. *Pesquisa Agropecuária Brasileira*, 47(5), 716-721.
- ⁷ Kammann, C. I., Linsel, S., Gößling, J. W., & Koyro, H. W. (2011). Influence of biochar on drought tolerance of *Chenopodium quinoa* Willd and on soil-plant relations. *Plant and soil*, 345(1), 195-210.
- ⁸ Deficit Irrigation Strategies. University of California Fruit Report.
http://ucanr.org/sites/fruitreport/Irrigation/Deficit_Irrigation_Strategies/. Accessed online 3-30-13.
- ⁹ Crowley, D. (2012). Biochar as a Soil Amendment for Avocado Production. California Avocado Society 2012 Yearbook 95:44-61. *Avocadosource.com*
- ¹⁰ Lehmann J., da Silva J.P., Steiner C., Nehls T., Zech W., Glaser B. (2003b) Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil* 249:343.
- ¹¹ Ventura, M., Sorrenti, G., Panzacchi, P., George, E., & Tonon, G. (2013). Biochar Reduces Short-Term Nitrate Leaching from A Horizon in an Apple Orchard. *Journal of Environmental Quality*, 42(1), 76-82.
- ¹² Eldridge, S., ChengRong, C., ZhiHong, X., Meszaros, I., Chan, K. Y., Gilkes, R. J., & Prakongkep, N. (2010). Greenwaste biochar potentially reduces nitrogen fertiliser losses. In *Proceedings of the 19th World Congress of Soil Science: Soil solutions for a changing world*, Brisbane, Australia, 1-6 August 2010. (pp. 100-103). International Union of Soil Sciences (IUSS), c/o Institut für Bodenforschung, Universität für Bodenkultur.
- ¹³ Briscoe, T. (2012, October 18). Charcoal Apples. Australian Broadcast Corporation.
<http://www.abc.net.au/rural/tas/content/2012/10/s3613529.htm?site=northtas>. Accessed online 3-29-13.
- ¹⁴ da Silva, J. H. (2013). Impacto del uso de biocarbón sobre la calidad de suelos y producción de cacao (*Theobroma cacao* L.) en sistemas agroforestales, Reserva Indígena Bribri, Talamanca, Costa Rica por Tesis sometida a consideración de la Escuela de Posgrado como requisito para optar por el grado de Magister Scientiae en Agroforestería Tropical Turrialba, Costa Rica.
- ¹⁵ International Biochar Initiative (2012). Carbon Gold: Working with Cacao Farmers in Belize to Create a Rotating Biochar Production and Utilization System. <http://www.biochar-international.org/carbongold>. Accessed online 3/18/13.
- ¹⁶ Menge, J., Davis, R., Johnson, E., & Zentmyer, G. (1978). Mycorrhizal fungi increase growth and reduce transplant injury in avocado. *California agriculture*, 32(4), 6-7.
- ¹⁷ Warnock, D. D., Lehmann, J., Kuyper, T. W., & Rillig, M. C. (2007). Mycorrhizal responses to biochar in soil—concepts and mechanisms. *Plant and Soil*, 300(1), 9-20.
- ¹⁸ Pereira, E., Suddick, E., Six, J. (2012). Use of Walnut Biochar to improve soil properties and reduce N₂O emissions. UC Davis Agroecology Lab.
<http://www.plantsciences.ucdavis.edu/Agroecology/Outreach/Walnut.html>. Accessed online, 3-21-13.
- ¹⁹ Ishii, T., & Kadoya, K. (1994). Effects of charcoal as a soil conditioner on citrus growth and vesicular-arbuscular mycorrhizal development. *Journal of the Japanese Society for Horticultural Science*, 63.
- ²⁰ Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., & Crowley, D. (2011). Biochar effects on soil biota—a review. *Soil Biology and Biochemistry*, 43(9), 1812-1836.

-
- ²¹ Saranya, K., Krishnan, P. S., Kumutha, K., & French, J. (2011). Potential for Biochar as an Alternate Carrier to Lignite for the Preparation of Biofertilizers in India. *International Journal of Agriculture, Environment and Biotechnology*, 4(2), 167-172.
- ²² Elad, Y., Cytryn, E., Meller Harel, Y. A. E. L., Lew, B., & Graber, E. R. (2012). The biochar effect: plant resistance to biotic stresses. *Phytopathologia Mediterranea*, 50(3), 335-349.
- ²³ Nerome, M., Toyota, K., Islam, T. M., Nishijima, T., Matsuoka, T., Sato, K., & Yamaguchi, Y. (2005). Suppression of bacterial wilt of tomato by incorporation of municipal biowaste charcoal into soil. *Soil Microorganisms*, 59.
- ²⁴ Henreaux, J. Thesis, Mag Sc in Organic Farming, Tropical Agricultural Research and Higher Education, Turrialba (Costa Rica). Effect of biochar and organic fertilizers combined with beneficial microorganisms on development, productivity and plant resistance, Turrialba, Costa Rica.
- ²⁵ Meller Harel, Y., Elad, Y., Rav-David, D., Borenstein, M., Shulchani, R., Lew, B., & Graber, E. R. (2012). Biochar mediates systemic response of strawberry to foliar fungal pathogens. *Plant and soil*, 1-13.
- ²⁶ Zwart, D. C., & Kim, S. H. (2012). Biochar Amendment Increases Resistance to Stem Lesions Caused by *Phytophthora* spp. in Tree Seedlings. *HortScience*, 47(12), 1736-1740.
- ²⁷ Zhang, X. K., Li, Q., Liang, W. J., Zhang, M., Bao, X. L. and Xie, Z. B. (2013). Soil nematode response to biochar addition in a Chinese wheat field. *Pedosphere*, 23(1), 98-103.
- ²⁸ Scott, D. H., Shaw, W. C., & Ruppenthal, R. U. (1954). Evaluation of several chemicals for weed control in strawberry fields. *Weeds*, 192-207.
- ²⁹ [Biochar for Arborists](#). Kelpie Wilson. Tree Care Industry Association. September 2012.
- ³⁰ Ogawa, M., & Okimori, Y. (2010). Pioneering works in biochar research, Japan. *Soil Research*, 48(7), 489-500.
- ³¹ Pacific-Area Wide Integrated Methyl Bromide Alternatives project, USDA, ARS. <http://ucanr.edu/sites/PAWMBA/> Accessed online 2-27-2013.
- ³² Mazzola, M., Granatstein, D. M., Elfving, D. C., Mullinix, K., & Gu, Y. H. (2002). Cultural management of microbial community structure to enhance growth of apple in replant soils. *Phytopathology*, 92(12), 1363-1366.
- ³³ Wang, Y., Pan, F., Wang, G., Zhang, G., Wang, Y., Chen, X., & Mao, Z. (2014). Effects of biochar on photosynthesis and antioxidative system of *Malus hupehensis* Rehd. seedlings under replant conditions. *Scientia Horticulturae*, 175, 9–15. <http://doi.org/10.1016/j.scienta.2014.05.029>
- ³⁴ Atucha, A., & Litus, G. (2015). Effect of Biochar Amendments on Peach Replant Disease. *HortScience*, 50(6), 863–868.
- ³⁵ Sun, K., Gao, B., Ro, K. S., Novak, J. M., Wang, Z., Herbert, S., & Xing, B. (2012). Assessment of herbicide sorption by biochars and organic matter associated with soil and sediment. *Environmental Pollution*, 163, 167-173.
- ³⁶ Mesa, A. C., & Spokas, K. A. (2011). Impacts of biochar (black carbon) additions on the sorption and efficacy of herbicides. *Herbicides and the Environment*, InTech, Croatia, 315-340.
- ³⁷ Graber, E. R., Tsechansky, L., Khanukov, J., & Oka, Y. (2011). Sorption, Volatilization, and Efficacy of the Fumigant 1, 3-Dichloropropene in a Biochar-Amended Soil. *Soil Science Society of America Journal*, 75(4), 1365-1373.
- ³⁸ Graber, E. R., Tsechansky, L., Gerstl, Z., & Lew, B. (2012). High surface area biochar negatively impacts herbicide efficacy. *Plant and soil*, 1-12.
- ³⁹ Nag, S. K., Kookana, R., Smith, L., Krull, E., Macdonald, L. M., & Gill, G. (2011). Poor efficacy of herbicides in biochar-amended soils as affected by their chemistry and mode of action. *Chemosphere*, 84(11), 1572-1577.
- ⁴⁰ Martin, S. M., Kookana, R. S., Van Zwieten, L., & Krull, E. (2012). Marked Changes in Herbicide Sorption-Desorption Upon Ageing of Biochars in Soil. *Journal of Hazardous Materials*.
- ⁴¹ Jones, D. L., Edwards-Jones, G., & Murphy, D. V. (2011). Biochar mediated alterations in herbicide breakdown and leaching in soil. *Soil Biology and Biochemistry*, 43(4), 804-813.
- ⁴² [Learning to Burn and Make Biochar, Not Smoke](#). Kelpie Wilson. Tree Care Industry Magazine. February 2014.
- ⁴³ Jeffery, S., Verheijen, F. G. A., Van der Velde, M., & Bastos, A. C. (2011). A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, Ecosystems & Environment*, 144(1), 175-187.
- ⁴⁴ [Justus von Liebig and the Roots of Modern Biochar](#). Kelpie Wilson. Ithaka Journal. August 2013.