MAKING & USING A PROBLEM SOLVER DEVICE

DR. RICK HARLOW


INTRODUCTION

- Everywhere there are problems. And everyone has them.
- What I am going to show you is the SIMPLE solution to many of these problems.
- USING IT WILL NOT HURT YOU.
- It is VERY easy to use
- IT WILL COST YOU LITTLE OR NOTHING TO MAKE.

WHAT YOU NEED TO MAKE THE DEVICE

- A 5-volt AC/DC power supply (mobile phone charger) or a 6-volt battery.
 - If you use a battery, you will need approximately 1 m (3 feet) of two stranded wire.
- Two pieces of cotton fabric about 25 cm X 25 cm (10" x 10")
- Plastic bags or plastic/latex gloves.
- Water.
- Baking Soda (Sodium Bicarbonate).


DEVICE READY TO ASSEMBLE



HOW TO MAKE A THE DEVICE

- Cut off the adapter tip (the part that plugs into the cell phone)
- Remove the insulation (melt the insulation about 2-3 cm from the end of the wire, then pull it off with a rag)
- Separate the two wires
- Wet the two pieces of cotton fabric in baking soda solution (mix 1 cc (1/4 tsp) with 250 mL (1 cup) water, and wring out the excess water)
- Ball up the cotton cloth around each of the bare wires

WIRE PLACED IN MIDDLE OF CLOTH



DEVICE READY TO BE USED



HOW TO USE A PROBLEM SOLVER DEVICE

- This will not solve all the problems. However, it will help with most problems, and as I said at the beginning, it will not hurt you.
- It is normal to feel a slight tingling or burning sensation while using the device
- Use the device daily to prevent problems

SIMPLE PROBLEMS

- Plug in the adapter, turn on the adapter, or connect the wires to the battery.
- Hold the balled up cotton fabric in your hands for between 10 minutes and 30 minutes.
 - Keep increasing treatment time until you notice an improvement.
 - Repeat every 2 hours until problem is gone

SEVERE PROBLEMS

- Hold your wrists on the fabric electrodes for 30 minutes.
 - REPEAT TREATMENT EVERY 2 HOURS UNTIL PROBLEM IS GONE

PROBLEMS IN THE HEAD

- Wearing a glove or plastic bag on one hand (this is, so the electricity does not go through the hand), place the electrodes over the left and right arteries and veins in the neck below the jaw.
 - Switch sides; move the right electrode to the left side and the left electrode to the right side, every 5 minutes.
 - Treat for 30 minutes per treatment.
 - Treat every 2-8 hours (as often as possible) until the problem(s) are gone

ALL PROBLEMS

 Repeat treatments every 2 to 8 hours; until the problems are solved.

 Use more frequently for more severe problems.

WHEN YOU ARE BETTER

 Treat for 10 minutes to 20 minutes once or twice a day to help prevent problems.

Email us at info@thl-u.org with questions, problems, translation errors, or suggestions.