Franklin County Noxious Weed Control Board

Mission Statement

The Franklin County Noxious Weed Control Board strives to serve as responsible stewards by supporting the protection of Franklin County's natural resources and the preservation of its rich agricultural heritage from the degrading impacts of noxious weeds.

We help landowners, land users and managers, government entities, and the local community comply with the state of Washington's weed law (RCW 17.10).

We accomplish our mission by:

- Executing a program emphasizing education in the identification and control of noxious weeds, and encouraging the control of priority species.
- Providing a trained staff of regulatory inspectors to survey for noxious weeds and work with landowners to formulate reasonable weed control plans compatible with our goals and, when necessary, to require controls.
- Maintaining a presence in the community by participating in local events, publishing an educational website and printed materials, and conducting regularly scheduled public board meetings.
- Offering technical assistance to encourage stewardship of natural resources by educating the community on the principles of integrated weed management.

The Franklin County Noxious Weed Control Board's complete Philosophy and Policy Statements are posted online at *fcweedboard.com/board-of-directors*.

About The Noxious Weed List

Noxious weeds are certain plant species the state has determined can cause ecological or economic harm.

The Noxious Weed List is updated annually and prioritized into three classes: A, B and C.

Class A Weeds are non-native species whose distribution in Washington is still limited. Preventing new infestations and eradicating existing ones are of the highest priority. Eradication of all Class A species is required by law statewide (RCW 17.10.140).

Class B and Class B-Designate Weeds are nonnative species presently limited to portions of the state. In regions where a Class B species is already abundant, control is decided at the local level. A further classification for certain Class B Weeds that occur in regions where they are not yet widespread is a Class B-Designate, which requires the prevention of seed production and dispersal of all propagative parts capable of forming a new plant. Preventing new infestations in such areas is a high priority.

Class C Weeds are species that are already widespread in Washington or are of special interest to the state's agricultural industry. A county weed board may require landowners to control a Class C Weed if it poses a threat to agriculture. Long-term programs of suppression and control are decided at the local level.

*WEED IN TRANSITION: Kochia is a noxious weed that is intensely managed in portions of Franklin County and selected for increased management efforts in future years countywide.

Controlling noxious weeds is the landowner's responsibility.

OUR WEEDS OF FOCUS

Noxious weeds we are actively working on or monitoring to prevent further spread

| babysbreath | puncturevine |
|-----------------------------------|--------------------|
| camelthorn | Ravenna grass |
| cereal rye | rush skeletonweed |
| common reed (nonnative genotypes) | saltcedar |
| Dalmatian toadflax | spikeweed |
| hoary cress | thistle, Canada |
| houndstongue | thistle, Scotch |
| jointed goatgrass | tree-of-heaven |
| knapweed, diffuse | velvetleaf |
| knapweed, spotted | white bryony |
| *kochia | yellow flag iris |
| longspine sandbur | yellow nutsedge |
| loosestrife, purple | yellow starthistle |
| poison hemlock | |





Perennial **rush skeletonweed** gained a foothold on thousands of acres in Franklin County while ground was out of wheat production in the Conservation Reserve Program (CRP). Skeletonweed should be controlled on all land that is coming out of CRP before going back into production.

Contact Us 509-545-3847

fcwb@franklincountywa.gov

Franklin County 2025 Noxious Weed List





Scotch thistle is a Class B noxious weed in Franklin County and a serious problem in rangeland. Single plants can be dug by hand but be sure to destroy the entire root. Large infestations will require herbicide to achieve control.



Dense stands of this spiny plant reduce forage production, prohibiting land utilization for grazing or recreation.

Weeds found in Franklin County are in gold font.

FCNWCB Noxious Weed List

Class A Weeds—Eradication required by law

| common crupina | Crupina vulgaris |
|---------------------------|------------------------------|
| cordgrass, common | Spartina anglica |
| cordgrass, dense-flowered | Spartina densiflora |
| cordgrass, saltmeadow | Spartina patens |
| cordgrass, smooth | Spartina alterniflora |
| dyer's woad | Isatis tinctoria |
| eggleaf spurge | Euphorbia oblongata |
| false-brome | Brachypodium sylvaticum |
| floating primrose-willow | Ludwigia peploides |
| flowering rush | Butomus umbellatus |
| French broom | Genista monspessulana |
| garlic mustard | Alliaria petiolata |
| giant hogweed | Heracleum mantegazzianum |
| goatsrue | Galega officinalis |
| hydrilla | Hydrilla verticillata |
| Johnsongrass | Sorghum halepense |
| knapweed, bighead | Centaurea macrocephala |
| knapweed, Vochin | Centaurea nigrescens |
| kudzu | Pueraria montana var. lobata |
| meadow clary | Salvia pratensis |
| oriental clematis | Clematis orientalis |

Class B Weeds—Continued

| perennial pepperweed | Lepidium latifolium |
|----------------------|------------------------|
| poison-hemlock | Conium maculatum |
| *policeman's helmet | Impatiens glandulifera |
| puncturevine | Tribulus terrestris |
| *Ravenna grass | Tripidium ravennae |
| *rough chervil | Chaerophyllum temulum |
| rush skeletonweed | Chondrilla juncea |
| *saltcedar | Tamarix ramosissima |
| *Scotch broom | Cytisus scoparius |
| *shiny geranium | Geranium lucidum |
| *spurge flax | Thymelaea passerina |
| *spurge laurel | Daphne laureola |
| *spurge, leafy | Euphorbia virgata |
| *spurge, myrtle | Euphorbia myrsinites |
| *sulfur cinquefoil | Potentilla recta |
| Tansy, common | Tanacelum vulgari |
| | |

Class A Weeds—Continued

| Palmer amaranth | Amaranthus pakmeri |
|----------------------------------|--|
| purple starthistle | Centaurea calcitrapa |
| reed sweetgrass | Glyceria maxima |
| ricefield bulrush | Schoenoplectus mucronatus |
| round leaf bittersweet | Celastrus orbiculatus |
| sage, clary | Salvia sclarea |
| sage, Mediterranean | Salvia aethiopis |
| silverleaf nightshade | Solanum elaeagnifolium |
| small-flowered jewelweed | Impatiens parviflora |
| South American | Limnobium laevigatum |
| spongeplant | |
| Spanish broom | Spartium junceum |
| Syrian beancaper | Zygophyllum fabago |
| Texas blueweed | Helianthus ciliaris |
| thistle, Italian | Carduus pycnocephalus |
| thistle, marsh | Cirsium palustre |
| thistle, milk | Silybum marianum |
| thistle, slenderflower | Carduus tenuiflorus |
| thistle, Turkish | Carduus cinereus |
| Variable-leaf milfoil and hybrid | Myriophyllum heterophyllum and M. heterophyllum x M. hippuroides |
| wild four-o'clock | Mirabilis nyctaginea |

Class B Weeds—Continued

| *tansy ragwort | Jacobaea vulgaris |
|-----------------------|------------------------|
| *thistle, musk | Carduus nutans |
| *thistle, plumeless | Carduus nutans |
| thistle, Scotch | Onopordum acanthium |
| *velvetleaf | Abutilon theophrasti |
| *water primrose | Ludwigia hexapetala |
| white bryony | Bryonia alba |
| *wild basil | Clinopodium vulgare |
| *wild chervil | Anthriscus sylvestris |
| *yellow archangel | Lamiastrum galeobdolon |
| *yellow floatingheart | Nymphoides peltata |
| yellow nutsedge | Cyperus esculentus |
| yellow starthistle | Centaurea solstitialis |

Class B Weeds—Control or containment required *Class B Designate Weeds—Control required

| *blueweed | Echium vulgare |
|--|---|
| *Brazilian elodea | Egeria densa |
| *bugloss, annual | Lycopsis arvensis |
| *bugloss, common | Anchusa officinalis |
| butterflybush | Buddleja davidii |
| *camelthorn | Alhagi maurorum |
| *common fennel (except | Foeniculum vulgare (except |
| bulbing fennel) | F. vulgare var. azoricum) |
| common reed (nonnative) | Phragmites australis |
| *Dalmatian toadflax | Linaria dalmatica ssp. |
| Eurasian watermilfoil | Myriophyllum spicatum |
| *European coltsfoot | Tussilago farfara |
| *fanwort | Cabomba caroliniana |
| *gorse | Ulex europaeus |
| *grass-leaved arrowhead | Sagittaria graminea |
| hairy willowherb | Epilobium hirsutum |
| *hawkweed oxtongue | Picris hieracioides |
| *hanging sedge | Carex pendula, C pendula subsp. Pendula and C pen- dula subsp. agastachys |
| *hawkweed, orange | Hieracium aurantiacum |
| *Hawkweeds: non-natives and hybrids of meadow subgenus | Hieracium, subgenus Pilosella |

Class C Weeds—Local Control

| babysbreath | Gypsophila paniculata |
|----------------------|-----------------------|
| buffalobur | Solanum rostratum |
| cereal rye | Secale cereale |
| common St. Johnswort | Hypericum perforatum |
| field bindweed | Convolvulus arvensis |
| hairy whitetop | Lepidium appelianum |
| hoary cress | Lepidium draba |
| jointed goatgrass | Aegilops cylindrica |

Class B Weeds—Continued

| *Hawkweeds: nonnatives and hybrids of wall subgenus | Hieracium subgenus Hieracium |
|---|------------------------------|
| *herb-Robert | Geranium robertianum |
| *hoary alyssum | Berteroa incana |
| *houndstongue | Cynoglossum officinale |
| indigobush | Amorpha fruticosa |
| *knapweed, black | Centaurea nigra |
| *knapweed, brown | Centaurea jacea |
| knapweed, diffuse | Centaurea diffusa |
| knapweed, meadow | Centaurea x gerstlaueri |
| knapweed, Russian | Rhaponticum repens |
| *knapweed, spotted | Centaurea stoebe |
| *knotweed, Bohemian | Fallopia x bohemica |
| *knotweed, giant | Fallopia sachalinensis |
| *knotweed, Himalayan | Persicaria wallichii |
| *knotweed, Japanese | Fallopia japonica |
| *kochia (weed in transition) | Bassia scoparia |
| *lesser celandine | Ficaria verna |
| *loosestrife, garden | Lysimachia vulgaris |
| loosestrife, purple | Lythrum salicaria |
| loosestrife, wand | Lythrum virgatum |
| Malta starthistle | Centaurea melitensis |
| *parrotfeather | Myriophyllum aquaticum |

Class C Weeds—Local Control

| longspine sandbur | Cenchrus longispinus |
|---------------------------|----------------------------|
| Medusahead | Taeniatherum caput-medusae |
| reed canarygrass | Phalaris arundinacea |
| smoothseed alfalfa dodder | Cuscuta approximata |
| spikeweed | Centromadia pungens |
| thistle, bull | Cirsium vulgare |
| thistle, Canada | Cirsium arvense |
| yellowflag iris | Iris pseudacorus |

Learn more about noxious weeds at fcweedboard.com