

# QUEEN GARNET

## ADVANCED FRUIT AND VEGETABLES

### Prunus salicina Fruit Extract

Delivering "true to nature" hydrophilic plant profiles. Cellular Extraction harnesses the species' water-soluble blueprint of phyto-compounds and their derivatives (the entourage), in their synergistic relationships, as they exist in the cell. Extracted rapidly under high pressure, we maintain the molecules' natural integrity. Committed to conscious manufacturing, Cellular Extracts are created using purified, deionised, rainharvested water and less plant material than past processes to produce a natural extract, that is eco-sustainable, traceable to source, transparent in composition – delivering the brilliance of nature.

The Queen Garnet Plum is a patented non-native Australian species, originally endemic to Japan, but supercharged its profile as a result of adapting to the harsh conditions of the pristine Australian environment. Globally recognised and highly prized as a unique species and source of anthocyanins, this species offers exciting new possibilities for benefits in Cosmetic, Nutraceutical, natural Pharmaceutical, Food and Beverage industry sectors.

Introducing the world first NE Queen Garnet Cellular Extract offering a powerhouse of antioxidants and an interesting profile of anthocyanins, chlorogenic acid, rutin, epicatechin, quercetin and more.

In pursuit to learn more about this species profile, we have listed publicly available, peer reviewed research on these compounds below. This is for educational purposes and is not based on clinical trials of the Cellular Extract. Extracts deliver natural molecules and connects your brand to nature.

## WATER-SOLUBLE PHYTO-

### COMPOUNDS

#### ANTHOCYANINS

- **Skin lightening / Depigmentation potential** Human tissue *in-vitro* studies show inhibition of Tyrosinase activity, a key enzyme in melanogenesis process leading to pigmentation<sup>1</sup>  
Studies show decreased melanin formation topically<sup>1</sup>  
Inhibit pigmentation format, blocks the process.
- **Photo-aging protection**  
Demonstrated to significantly reduce UVA<sup>-13,14</sup> and UVB-induced DNA damage, large variety of photo-ageing and, protein oxidation and lipid peroxidation within reconstituted human skin tissue, dermal fibroblasts and keratinocytes<sup>15,20</sup>  
Reduces redness and sensitivity by inhibiting overexpression and activity of matrix metalloproteinases (MMPs) e.g. collagenase, elastase, thereby protecting the cells from UV-induced apoptosis<sup>15-20</sup>, presumably through its UVB-dominant absorption capacity reported at approximately 46%<sup>21</sup>
- **Anti-aging / Conditioning / Inhibitor damage** Potential anti-aging benefits through protective anti-proliferative effects against damaged skin cells, and uncontrolled division thereof<sup>6,14</sup>  
Protect cells from premature death/apoptosis induced metastasis control<sup>3-6,14</sup>  
Protection against oxidative stress improving skin longevity and maintains healthy condition<sup>3-6,14</sup>
- **Maintains skin viability, collagen and elastin support** Protects collagen-producing cells (fibroblasts) against oxidative damage via inhibition of peroxy radicals production<sup>8</sup>  
Captures the environmental and internal heavy metals that induce aging<sup>8</sup>  
Reduces amounts of damaging free radicals, lipid breakage (peroxidation), and DNA damage to prolong good skin, improving flexibility and functionality<sup>9</sup>
- **Anti-wrinkle and skin flexibility – anti-glycation activity**  
Prevents glycation process and protein damage<sup>30</sup>  
Alleviate tissue inflammation, reduced activation of stress cascades to improve skin cell function<sup>32</sup>
- **Potent antioxidant properties – improved skin life cycle, function & quality**  
Supported significant decrease of induced DNA damage and intracellular ROS, along with an increase in cellular defence enzyme<sup>22,23</sup>  
Holds antioxidant capacity towards oxygen radicals<sup>24,25</sup> and oxidised ferric species<sup>26-28</sup>, scavenging activity towards superoxide<sup>29</sup> and peroxynitrite<sup>30</sup>, inhibition of human LDL oxidation<sup>31</sup> and lipid peroxidation<sup>30</sup>, pro-antioxidant activity in form of ability to bind heavy metals such as iron, zinc and copper<sup>32</sup> and induction of intracellular defence enzymes<sup>33,34</sup>
- **Anti-inflammatory activity – inflammation reduction**<sup>42</sup>  
Reported to inhibit the expression of cytokines (molecules being produced in response to stress and

imitation) such as cyclooxygenase-2 (COX-2) and inducible nitric oxide (iNOS) protein, involved in many inflammatory processes, by suppressing NF- $\kappa$ B through down regulation of MAPK-mediated pathways<sup>35,38</sup> with almost no change to the antioxidative status *in-vivo*<sup>39,40</sup>

#### QUERCETIN

- **Anti-inflammatory activity via radical scavenging, inflammatory messengers and signalling pathways' inhibition**  
Inhibitor of tumour necrosis factor-alpha (TNF- $\alpha$ ) production *in-vitro*, as well as of numerous inflammatory mediators release<sup>1,2-5</sup>  
Inhibits epidermal and liver cyclooxygenase (COX) and lipoxygenase (LOX) activities<sup>6,7</sup>  
Exerts broad scavenging activity against reactive oxygen and nitrogen species<sup>48,9</sup>
- **Maintains skin-tissue vitality both by moderating sun exposure impact and exerting anti-glycation activities**  
Elevates epidermal cells' (keratinocytes) intracellular antioxidant activity following UVA irradiation, hence suppresses UVA-induced cell death (apoptosis) in human<sup>20,11</sup> and animal<sup>12</sup> keratinocytes  
Significantly prevents UVB-induced glutathione (GSH) depletion and tissue degradation in animal model<sup>13</sup>  
Produce sun protection factor (SPF) values similar to those of the classical sunscreen substances benzophenone<sup>41</sup> and homosalate (a reference filter used to establish FDA standards<sup>15</sup>  
Deemed a desirable compound for incorporation into sunscreen formulations due to excellent UVA/UVB absorption capacity<sup>16</sup>  
Found to significantly reduce secretion of tissue-degrading enzymes upon UVA or UVB exposure, thereby maintaining skin integrity via topical application<sup>17</sup>  
Represents a useful stabilising additive for the formulation of effective broad-spectrum sunscreens containing widely used and readily available filters<sup>18</sup>
- **Skin appearance enhancer via anti-glycation and potential longevity improvement**  
Presents good penetration of outermost skin layer (*Stratum Corneum*) and retention in viable epidermis when incorporated into emulsions<sup>19,20</sup>  
Dose dependently increases production of hyaluronic acid in epidermis cell line, elevating skin moisture and improve facial wrinkles<sup>21</sup>  
Inhibits AGEs formation via chelating catalytic metal ions and entrapment of both glycation mediators and reactive oxygen species<sup>22,23</sup>  
Shown to suppress the levels of AGEs and fructosamines<sup>24</sup>  
Identified as proteasome activator with antioxidant properties that consequently influence cells' lifespan, survival and viability, while creating a rejuvenating effect over senescent human fibroblasts<sup>25</sup>  
Associated with under-eye area skin rejuvenation by



#### PRODUCT

NE Queen Garnet Cellular Extract  
Concentrate: ANE0616  
Standard 1:10: ANE0616-10

#### POTENTIAL APPLICATIONS

- Body clays, muds, scrubs
- Body moisturisers, milks, lotions
- Day, night creams, lotions
- Eye contour products, serums
- Face masks, scrubs
- Face strips, patches
- Generally, suits most cosmetic applications
- Hair, scalp, products, treatments
- Hand, feet creams
- Hand liquid soaps, towellettes, wipes
- Hyperpigmentation creams, serums
- Men, teenage products, serums
- Sun care, after sun products

#### POSSIBILITIES TO TRIAL

- Anti-aging care, antioxidant care
- Anti-glycation, barrier function
- Blemish care (acne prone skin)
- Collagen, elastin support
- Dark pigmented areas, exfoliation
- Fine lines and wrinkles
- Hair growth, maintenance, protection
- Hair maintenance, protection
- Inflammation reduction
- Reactive skin conditions (dermatitis, psoriasis, rosacea)
- Skin conditioning, firmness, hydration
- Skin radiance, luminosity
- Skin improvement, wound healing

Possibilities are based on compound information and not on the extract information. This information is to assist you in your R&D.

#### PRODUCTION

Water/Glycerin Extract  
Added at the end phase, <40°C  
Ethanol option available

#### PLANT PART: ORIGIN

Fruit : Australia

#### USAGE GUIDE

Concentrate: 1.0 - 2.0%  
Standard 1:10: 3.0 - 6.0% Wash on/off formulations  
24 months unopened from date of manufacture

#### COMPLIANCE

INCI: Prunus salicina Fruit Extract  
CAS: 1214268-59-3  
EC: Not allocated

Cellular Extraction delivers the natural water-soluble compounds of a species. To assist your ingredient research we identified publicly available information on these compounds from other sources and their potential uses. Information is not based on specific clinical trials of the Cellular Extract.

Want to work with natural molecules as they exist in the cell?

Copyright © NATIVE EXTRACTS Pty Ltd 2017. All rights reserved. DISCLAIMER: The above information is provided for general educational information purposes only and should not be relied upon or used as the sole basis for making decisions without consulting primary, more accurate, more complete or more timely sources of information. The information contained has been gathered from publicly available current and reliable source material and is intended to provide general information to readers. NATIVE EXTRACTS Pty Ltd is not responsible for any reliance readers place on the material in this document and readers rely on the information in this document at their own risk and we recommend efficacy testing to be done on the finished product formulation to determine if it meets your specific target performance. However, this information is not intended to constitute an "authoritative statement" under the National Industrial Chemical Notification and Assessment Scheme Australia and New Zealand rules and regulations. The range of compounds delivered in NATIVE EXTRACTS Pty Ltd products is only a guide as there will be fluctuations in the range available per batch as this is a natural product and reflects the nature of natural differences from harvest to harvest, source to source, batch to batch etc. NATIVE EXTRACTS Pty Ltd provides the information contained herein in good faith but makes no representation as to its comprehensiveness or accuracy. Individuals receiving this information must exercise their independent judgment in determining its appropriateness for a particular purpose. As the ordinary or otherwise use(s) of this product is outside the control of NATIVE EXTRACTS Pty Ltd, no representation or warranty, expressed or implied is made as to the effect(s) of such use(s), (including damage or injury), or the results obtained. NATIVE EXTRACTS Pty Ltd expressly disclaims responsibility as to the ordinary or otherwise use(s). Furthermore, nothing herein should be considered as a recommendation by NATIVE EXTRACTS Pty Ltd as to the fitness for any use. The liability of NATIVE EXTRACTS Pty Ltd is limited to the value of the goods and does not include any consequential loss. NATIVE EXTRACTS Pty Ltd shall not be liable for any errors or delays in the content, or for any actions taken in reliance thereon. NATIVE EXTRACTS Pty Ltd shall not be responsible for any damage resulting from use of or reliance upon this information. The user of the product is solely responsible for compliance with all laws and regulations applying to the use of the products, including intellectual property rights of third parties.



# QUEEN GARNET

ADVANCED FRUIT AND  
VEGETABLES

Prunus salicina Fruit Extract

taking a role in improving dark circles and coarse wrinkles<sup>26</sup>

- **Prevents hair loss and promotes hair fibre proliferation**

Demonstrated to provide effective treatment for alopecia areata (AA) as well as reduce spontaneous onset of AA in preclinical trials<sup>27</sup>  
Shows protective effects against induced toxicity and hair cell damage *in-vivo*<sup>28</sup>

Effectively and positively affects human scalp in clinical trials, presenting 214% hair growth compared to control and 94% growth compared to commercially available treatment (Minoxidil), in combination with EGCg<sup>29</sup>

- **Potentially holds skin whitening effect**

Reducing levels of the melanogenic tyrosinase *in-vitro*, associated with, but not limited to the activation of proteasome<sup>25</sup>

## RUIN

- **Multifunctional activity as antioxidant, anti-inflammatory and organo-protective agent – maintaining vascular integrity and well being<sup>4</sup>**

- **Holds skin whitening effects**

Anti-melanogenic activity via inhibition of tyrosinase and related proteins<sup>57</sup>

- **Completely inhibits hair follicle cell death and halts regression**

When applied in combination with accompanying compounds<sup>8</sup>

- **Improves skin elasticity and reduces wrinkle formation through anti-glycation activities**

With high antioxidant activity, it effectively inhibited the generation of pre-AGE products and exhibited inhibitory effects on the formation of other intermediates and AGEs<sup>9</sup>

Evidently penetrates outermost skin layer (Stratum Corneum)<sup>10</sup> and inhibits glycation of type I collagen *in-vitro*<sup>11</sup>

Increased elasticity and decreased the length, area and number of wrinkles<sup>12</sup>

Holds strong affinity to suppress the formation of both initial and advanced stages of the glycation process as well as collagen crosslinking<sup>13-15</sup>

- **Effective and safe compound to be used in multifunctional sunscreens**

Robustly elevated clinical solar protection factor (SPF) by about 70% when combined into a formulation containing UV filters (sunscreens agents)<sup>16</sup>  
Produce SPF values similar to that of homosalate (a reference UV filter used to establish FDA standards)<sup>17</sup>

## EPICATECHIN

- **Protects from skin aging due to sun exposure**

Outstanding antioxidants<sup>11,16</sup>

Demonstrated to protect skin cells against UVB-induced damage by modulating cellular defence enzymes' activity<sup>3</sup>

Shown to inhibit skin proteins' and tissue degradation caused by damaging UV exposure<sup>4</sup>  
Prevents collagen breakdown by inhibiting collagenase by up to 70%<sup>5</sup>

- **Maintains skin flexibility and prevents wrinkle formation**

Exerts significant inhibitory effect on skin structural proteins glycation<sup>6</sup>, acting indirectly on the crosslinking process<sup>17</sup>

- **Exerts significant wound healing properties and alleviates swelling**

Reduces swelling and exudates, accelerates wound closure and enhances amounts of fibroblasts and collagen fibres in affected region<sup>4,20</sup>

Inhibits swelling (edema) and associated pain<sup>5,8</sup>

- **Counteracts acne via broad spectrum anti-inflammatory properties**

Suppresses inflammation by inhibiting expression and secretion of inflammatory messengers<sup>9,15,16,21-23</sup>  
Reduces sebum secretion in topical treatment of acne<sup>10</sup>

## FLAVONOID GLYCOSIDES

- **Large group of natural substances with various phenolic structures derived from the plant world, well known for their beneficial countless effects on health<sup>12</sup>**

Considered indispensable component in a variety of nutraceutical, pharmaceutical, medicinal and cosmetic applications

Possess anti-oxidative, anti-inflammatory, anti-microbial and anti-aging properties coupled with their capacity to modulate key cellular enzyme function

Hold various protective roles against numerous human diseases

## PHENOLICS

- **Key class of large and diverse dietary compounds, natural antioxidants<sup>12</sup>**

- **Exhibit a variety of functions, including: Antioxidant activity – neutralise or capture toxic and mutagenic species constantly forming in the skin as by product of metabolism<sup>16</sup>**

**Anti-inflammatory properties** – Responding to the exposure to external factors which may cause various types of damage, internal injuries, irritation or allergies<sup>79</sup>

**Antimicrobial action** – possess potent antifungal, antiviral and antibacterial activity, as an alternative to broad spectrum antibiotics which may lead to negative influence on skin microflora and resistance of many bacterial strains<sup>10,14</sup>

- **Anti-aging benefits:**

**Skin cell renewal** – can slow down or even reverse the decreasing capacity of epidermis to replace and detoxify dead cells, which rapidly declines with age<sup>15,17</sup>  
**Stimulation of collagen and elastin synthesis** – facilitate maintenance of proper skin structure and integrity of skin tissue, thereby stalling fine line and wrinkle formation<sup>18,23</sup>

**Attenuation of pigment (melanin) production in epidermis** – excessive production of melanin caused predominantly by UV exposure, as well as drugs, chemicals and particular disease states, may lead to dermal disorders<sup>24-30</sup>

**Protection from sun damage (UV radiation)** – Restores the antioxidant defence properties of the skin and attenuates the accumulation of DNA alterations<sup>31,36</sup>

**Preventing or attenuating the progression of certain skin disorders** – such as, but not limited to,

psoriasis<sup>37,38</sup>, rosacea<sup>39</sup>, a acne<sup>40-42</sup>, allergies<sup>43,44</sup>

dermatitis<sup>45</sup> and dermatophytosis<sup>46,47</sup>  
**Enhanced healing of incised and chronic wounds and burns** – by accelerating the following cascade of regenerative processes<sup>48-51</sup>

## CHLOROGENIC ACID (CGA)

- **Potent antioxidant and chelating agent stable under UV radiation**

Can serve as topical agent due to not degrading under UVA or UVB irradiation<sup>1</sup>

Significantly attenuated the accumulation of aluminium-based biotoxicity and associated oxidative stress<sup>2</sup>  
Provides effective defence against hydrogen-peroxide oxidative stress to protect tissues from adverse effects<sup>3</sup>

- **Long term skin lightening potential**

Natural suppressor of pigment formation<sup>4</sup>

- **Accelerates wound healing**



## PRODUCT

NE Queen Garnet Cellular Extract  
Concentrate: ANE0616  
Standard 1:10: ANE0616-10

## POTENTIAL APPLICATIONS

- Body clays, muds, scrubs
- Body moisturisers, milks, lotions
- Day, night creams, lotions
- Eye contour products, serums
- Face masks, scrubs
- Face strips, patches
- Generally, suits most cosmetic applications
- Hair, scalp, products, treatments
- Hand, foot creams
- Hand liquid soaps, towellettes, wipes
- Hyperpigmentation creams, serums
- Men, teenage products, serums
- Sun care, after-sun products

## POSSIBILITIES TO TRIAL

- Anti-aging care, antioxidant care
- Anti-glycation, barrier function
- Blemish care (acne prone skin)
- Collagen, elastin support
- Dark pigmented areas, exfoliation
- Fine lines and wrinkles
- Hair growth, maintenance, protection
- Hair maintenance, protection
- Inflammation reduction
- Reactive skin conditions (dermatitis, psoriasis, rosacea)
- Skin conditioning, firmness, hydration
- Skin radiance, luminosity
- Skin improvement, wound healing

Possibilities are based on compound information and not on the extract information. This information is to assist your R&D team

## PRODUCTION

Water/Glycerin Extract  
Added at the end phase, <40°C  
Ethanol option available

## PLANT PART: ORIGIN

Fruit : Australia

## USAGE GUIDE

Concentrate: 1.0 - 2.0%  
Standard 1:10: 3.0 - 6.0% Wash on/off formulations  
24 months unopened from date of manufacture

## COMPLIANCE

INCI: Prunus salicina Fruit Extract  
CAS: 1214268-59-3  
EC: Not allocated

Cellular Extraction delivers natural water-soluble compounds from a species. To assist your ingredient research we identified publicly available information on these compounds from other sources and their potential uses. Information is not based on specific clinical trials of the Cellular Extract.

Reduce your carbon footprint. Extracted using rain-harvested water

Copyright ©NATIVE EXTRACTS Pty Ltd 2017. All rights reserved. DISCLAIMER: The above information is provided for general educational information purposes only and should not be relied upon or used as the sole basis for making decisions without consulting primary, more accurate, more complete or more timely sources of information. The information contained has been gathered from publicly available current and reliable source material and is intended to provide general information to readers. NATIVE EXTRACTS Pty Ltd is not responsible for any reliance readers place on the material in this document and readers rely on the information in this document at their own risk and we recommend efficacy testing to be done on the finished product formulation to determine if it meets your specific target performance. However, this information is not intended to constitute an 'authoritative statement' under the National Industrial Chemical Notification and Assessment Scheme Australia and New Zealand rules and regulations. The range of compounds delivered in NATIVE EXTRACTS Pty Ltd products is only a guide as there will be fluctuations in the range available per batch as this is a natural product and reflects the nature of natural differences from harvest to harvest, source to source, batch to batch etc. NATIVE EXTRACTS Pty Ltd provides the information contained herein in good faith but makes no representation as to its comprehensiveness or accuracy. Individuals receiving this information must exercise their independent judgment in determining its appropriateness for a particular purpose. As the ordinary or otherwise use(s) of this product is outside the control of NATIVE EXTRACTS Pty Ltd, no representation or warranty, expressed or implied, is made as to the effect(s) of such use(s), including damage or injury, or the results obtained. NATIVE EXTRACTS Pty Ltd expressly disclaims responsibility as to the ordinary or otherwise use(s). Furthermore, nothing herein should be construed as a recommendation by NATIVE EXTRACTS Pty Ltd as to the fitness for any use. The liability of NATIVE EXTRACTS Pty Ltd is limited to the value of the goods and does not include any consequential loss. NATIVE EXTRACTS Pty Ltd shall not be liable for any errors or delays in the content, or for any actions taken in reliance thereon. NATIVE EXTRACTS Pty Ltd shall not be responsible for any damage resulting from use of or reliance upon this information. The user of the product is solely responsible for compliance with all laws and regulations applying to the use of the products, including intellectual property rights of third parties.



# QUEEN GARNET

ADVANCED FRUIT AND  
VEGETABLES

Prunus salicina Fruit Extract



Supports wound healing in a dose-dependent manner by enhancing antioxidant defence, fibroblast migration to affected area and angiogenesis (blood vessel formation)<sup>57</sup>

### • Possesses significant anti-inflammatory activities

Suppresses signalling pathways involved in inflammatory response and cell death (apoptosis) by decreasing secretion of pro-inflammatory cytokines<sup>89</sup>  
Reversed inflammatory cytokine (IL-1 $\beta$ )-mediated downregulation of collagen II<sup>10</sup>

### • Exerts anti-wrinkle activity and maintains skin firmness and flexibility

Inhibits dermal glycation, thus preventing formation of altered rigid protein-sugar crosslinked structures<sup>11,12</sup>  
Stimulates up to 2-fold increase in synthesis of collagen, elastin and associated polysaccharides<sup>13</sup>  
Decreases levels of matrix degrading enzymes (MMPs) and increases collagen and hyaluronic acid contents<sup>14</sup>

### • Alleviates skin irritation and redness

Prevents UV-induced Erythema (skin reddening)<sup>15</sup>

### ORGANIC ACIDS

• Immense group of multifunctional ingredients encompassing phenolic, carboxylic, phenol-carboxylic and sulfonic acids, as well as weaker acids containing enol, alcohol and thiol residues.

• In addition to their antimicrobial and anti-acne<sup>1</sup> properties, organic acids can also provide skin discoloration benefits<sup>1</sup>, exfoliating<sup>2</sup>, emollient<sup>2</sup>, moisturising<sup>3</sup> and conditioning properties.

• Internally, organic acids are known for its antioxidant and scavenging capacities against naturally occurring radicals, suppression of inflammatory agents<sup>5</sup> and positive effects on microbiome<sup>6</sup>.

### SUGARS

#### • Useful humectants

Hydrophilic, water soluble and in most cases hygroscopic (absorbs water vapour from the air) because their hydroxyl groups form hydrogen bonds with water molecules, resulting in their hydration and dissolution into the hosting skin<sup>1</sup>

### AMINO ACIDS

• Amino Acids (AAs) are the building blocks of proteins and peptides (proteins too small to make up tertiary structures). More than 300 AAs have been described, however only 20 AAs take part in protein synthesis.

There are 11 non-essential AAs which our body is able to synthesise, and 9 essential AAs which we must supplement from our diet<sup>1</sup>

• AAs are generally known to regulate gene expression<sup>2</sup> and protein metabolism e.g. skin collagen synthesis rates<sup>3</sup>, enhance wound healing<sup>4,5</sup>, exert antioxidant activities<sup>5</sup>, strengthen the immune system<sup>7</sup>, improve skin barrier function<sup>8</sup>, maintain skin vitality as anti-aging

agents<sup>9</sup>, contribute to hair viability by constituting the building blocks for hair strand synthesis alongside multiple aspects of hair maintenance and protection<sup>10,15</sup>, and play a pivotal role in disease prevention<sup>16,18</sup>

### REFERENCE

Peaks identified and verified by an independent Australian Government body, Southern Cross University, SDS and Technical Specification available on request.

#### ANTHOXYANINS

1. Jhan J K et al., Int J Cosmet Sci., 2006; 38(3): 319-24
2. Bowen-Forbes CS et al., Horticulture Biodiversity Nutr., 2010; 23(6): 554-60
3. Bunea A et al., Phytochemistry, 2013; 95: 436-44
4. Diaconeasa Z et al., Int J Mol Sci., 2015; 16(2): 2326-5
5. Wang E et al., Food Nutr Res., 2017; 61(1): 1325308
6. Afaq F et al., Int J Cancer., 2005; 113(3): 423-33
7. Lu J N et al., Int J Oncol., 2014; 44(5): 1756-66
8. Solomon A et al., J Agric Food Chem., 2010; 58(11): 6660-5
9. Giampieri F et al., Molecules., 2014; 19(6): 7738-816
10. Thilavach T et al., Int J Biol Macromol., 2015; 75: 515-20
11. Rojo LE et al., Food Chem., 2012; 131(2): 387-96
12. Valenza A et al., Biomed Res Int., 2018; 2018: 6413172

13. Giampieri F et al., J Agric Food Chem., 2012; 60(9): 2322-7
14. Tarozzi A et al., Photochem Photobiol., 2005; 81(3): 629
15. Afaq F et al., Exp Dermatol., 2009; 18(6): 553-61
16. Afaq F et al., J Invest Dermatol., 2007; 127(1): 222-32
17. Afaq F et al., Photochem Photobiol., 2010; 86(6): 1318-26
18. Bae J Y et al., Mol Nutr Food Res., 2009; 53(6): 726-38
19. Cimino F et al., J Agric Food Chem., 2006; 54(11): 4041-7
20. Tsoyi K et al., J Agric Food Chem., 2008; 56(22): 10600-5
21. Chan C F et al., J Cosmet Sci., 2010; 61(5): 333-41
22. Schantz M et al., J Berry Res., 2010; 1(1): 25-33
23. Ghosh D et al., J Sci Food Agric., 2008; 86(5): 678-86
24. Zheng W et al., J Agric Food Chem., 2008; 51(2): 502-509
25. Steed L E et al., J Food Sci., 2008; 73(5): S215-21
26. Moyer R A et al., J Agric Food Chem., 2002; 50(3): 519-25
27. Lee J et al., J Agric Food Chem., 2004; 52(23): 7039-44
28. Koca I et al., Sci Hort., 2009; 121(4): 447-50
29. Constantino L et al., Planta Med., 1992; 58(4): 342-4
30. Muselik J et al., Int J Mol Sci., 2007; 8(8): 797-809
31. Heinonen I M et al., J Agric Food Chem., 1998; 46(10): 4107-12
32. Havsteen B., Biochem Pharmacol., 1983; 32(7): 1141-8
33. Fiander H et al., Cancer Lett., 2000; 156(2): 117-24
34. Turner M. K., MSc thesis., 2009, Athens, Georgia USA
35. Wang H et al., J Nat Prod., 1999; 62(2): 294-6
36. Hou D X et al., Biochem Pharmacol., 2005; 70(3): 417-25
37. Pergola C et al., Nitric Oxide., 2006; 15(1): 30-9
38. Wang L S et al., Cancer Lett., 2008; 269(2): 281-90
39. Xia X et al., J Nutr., 2005; 136(8): 2220-5
40. Wang Q et al., Asia Pac J Clin Nutr., 2007; 1: 2531
41. Chawla A et al., Nat Med., 2001; 7(1): 48-52
42. Walker D et al., Arterioscler Thromb Vasc Biol., 2006; 26(5): 1022-8
43. Giampieri F et al., Molecules., 2014; 19(6): 7738-816
44. Thilavach T et al., Int J Biol Macromol., 2015; 75: 515-20
45. Rojo LE et al., Food Chem., 2012; 131(2): 387-96
46. Valenza A et al., Biomed Res Int., 2018; 2018: 6413172
47. Giampieri F et al., J Agric Food Chem., 2012; 60(9): 2322-7
48. Tarozzi A et al., Photochem Photobiol., 2005; 81(3): 629
49. Afaq F et al., Exp Dermatol., 2009; 18(6): 553-61
50. Afaq F et al., J Invest Dermatol., 2007; 127(1): 222-32
51. Afaq F et al., Photochem Photobiol., 2010; 86(6): 1318-26
52. Bae J Y et al., Mol Nutr Food Res., 2009; 53(6): 726-38
53. Cimino F et al., J Agric Food Chem., 2006; 54(11): 4041-7
54. Tsoyi K et al., J Agric Food Chem., 2008; 56(22): 10600-5
55. Chan C F et al., J Cosmet Sci., 2010; 61(5): 333-41
56. Schantz M et al., J Berry Res., 2010; 1(1): 25-33
57. Ghosh D et al., J Sci Food Agric., 2008; 86(5): 678-86
58. Zheng W et al., J Agric Food Chem., 2008; 51(2): 502-509
59. Steed L E et al., J Food Sci., 2008; 73(5): S215-21
60. Moyer R A et al., J Agric Food Chem., 2002; 50(3): 519-25
61. Lee J et al., J Agric Food Chem., 2004; 52(23): 7039-44
62. Koca I et al., Sci Hort., 2009; 121(4): 447-50
63. Constantino L et al., Planta Med., 1992; 58(4): 342-4
64. Muselik J et al., Int J Mol Sci., 2007; 8(8): 797-809
65. Heinonen I M et al., J Agric Food Chem., 1998; 46(10): 4107-12
66. Havsteen B., Biochem Pharmacol., 1983; 32(7): 1141-8
67. Fiander H et al., Cancer Lett., 2000; 156(2): 117-24
68. Turner M. K., MSc thesis., 2009, Athens, Georgia USA
69. Wang H et al., J Nat Prod., 1999; 62(2): 294-6
70. Hou D X et al., Biochem Pharmacol., 2005; 70(3): 417-25
71. Pergola C et al., Nitric Oxide., 2006; 15(1): 30-9
72. Wang L S et al., Cancer Lett., 2008; 269(2): 281-90
73. Xia X et al., J Nutr., 2005; 136(8): 2220-5
74. Wang Q et al., Asia Pac J Clin Nutr., 2007; 1: 2531
75. Chawla A et al., Nat Med., 2001; 7(1): 48-52
76. Walker D et al., Arterioscler Thromb Vasc Biol., 2006; 26(5): 1022-8

#### QUERCININ

1. Myhrstad M C et al., Free Radic Biol Med., 2002; 32(5): 386-93
2. Endale M et al., Immunobiology., 2013; 218(12): 1452-67
3. Bureau G et al., J Neurosci Res., 2008; 86(2): 403-10
4. Manjeet K R and Ghosh B., Int J Immunopharmacol., 1999; 21(7): 435-43
5. Kempuraj D et al., Br J Pharmacol., 2005; 145(7): 934-44
6. Kim H P et al., Prostaglandins Leukot Essent Fatty Acids., 1998; 58(1): 17-24
7. Lee K M et al., J Agric Food Chem., 2010; 58(9): 5815-20
8. Lopez-Lopez G et al., Mol Pharmacol., 2004; 65(4): 851-9
9. Ertug P U et al., Fundam Clin Pharmacol., 2010; 24(2): 223-32
10. Kimura S et al., Biochem Biophys Res Commun., 2009; 387(1): 109-14
11. Vincertini F T et al., J Dermatol Sci., 2011; 61(3): 162-8
12. Park J M et al., J Korean Soc Appl Biol Chem., 2012; 55(3): 443-6
13. Vincertini F T et al., Eur J Pharm Biopharm., 2008; 69(3): 948-57
14. Kostyuk V et al., Rejuvenation Res., 2018; 21(2): 91-101
15. Choquet B et al., J Nat Prod., 2008; 71: 1117-1118
16. Gonçalves M D C et al., Quim Nova., 2019; 42(4): 365-370
17. Fahlman B M., PhD Thesis, College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, Canada, 2010
18. Scalia S and Mezzena M., Photochem Photobiol., 2010; 86(2): 273-278
19. Zillikh OV et al., Int J Cosmet Sci., 2013; 35(5): 491-501
20. Dal Belo SE et al., Skin Pharmacol Physiol., 2009; 22(6): 299-304
21. Kim S H et al., J Soc Cosmet Scientists Korea., 2005; 31(1): 97-102
22. Bhuiyan, M.N.I. et al., Biosci. Biotechnol. Biochem., 2017; 81(5): 882-890
23. Li, X. et al., J. Agric. Food Chem., 2014; 62(50): 12152-12158
24. Morimitsu, Y. et al., Biosci. Biotechnol. Biochem., 1995; 59(11): 2018-2021
25. Chondrogian N et al., Exp Gerontol., 2010; 45(10): 763-71
26. Colvan L E et al., J Cosmet Dermatol., 2019; 18(2): 530-8
27. Wikramanayake T C et al., Cell Stress Chaperons., 2012; 17(2): 2574
28. Lee S K et al., Hum Exp Toxicol., 2015; 34(11): 1043-52
29. Lefevre F et al., J Cosmet Sci., 2014; 10: 1-3

#### RUBIN

1. Milde J et al., Phytomedicine., 2004; 11(2-3): 105-13
2. Hattemariam S et al., Mini Rev Med Chem., 2015; 15(7): 524-8
3. Lee W et al., Food Chem Toxicol., 2012; 50(9): 3048-55
4. Selloum L et al., Exp Toxicol Pathol., 2003; 54(4): 303-18



### PRODUCT

NE Queen Garnet Cellular Extract  
Concentrate: ANE0616  
Standard 1:10: ANE0616-10

### POTENTIAL APPLICATIONS

- Body clays, muds, scrubs
- Body moisturisers, milks, lotions
- Day, night creams, lotions
- Eye contour products, serums
- Face masks, scrubs
- Face strips, patches
- Generally, suits most cosmetic applications
- Hair, scalp, products, treatments
- Hand, feet creams
- Hand liquid soaps, towellettes, wipes
- Hyperpigmentation creams, serums
- Men, tennage products, serums
- Sun care, after-sun products

### POSSIBILITIES TO TRIAL

- Anti-aging care, antioxidant care
- Anti-glycation, barrier function
- Blemish care (acne prone skin)
- Collagen, elastin support
- Dark pigmented areas, exfoliation
- Fine lines and wrinkles
- Hair growth, maintenance, protection
- Hair maintenance, protection
- Inflammation reduction
- Reactive skin conditions (dermatitis, psoriasis, rosacea)
- Skin conditioning, firmness, hydration
- Skin radiance, luminosity
- Skin improvement, wound healing

Possibilities are based on compound information and not on the extract information. This information is to assist you in your R&D.

### PRODUCTION

Water/Glycerin Extract  
Added at the end phase, <40°C  
Ethanol option available

### PLANT PART: ORIGIN

Fruit : Australia

### USAGE GUIDE

Concentrate: 1.0 - 2.0%  
Standard 1:10: 3.0 - 6.0% Wash on/off formulations  
24 months unopened from date of manufacture

### COMPLIANCE

INCI: Prunus salicina Fruit Extract  
CAS: 1214268-59-3  
EC: Not allocated

Cellular Extraction delivers the natural water-soluble compounds of a species. To assist your ingredient research we identified publicly available information on these compounds from other sources and their potential uses. Information is not based on specific clinical trials of the Cellular Extract.

CLEAN & FUNCTIONAL – delivering the brilliance of natures design

Copyright © NATIVE EXTRACTS Pty Ltd 2017. All rights reserved. DISCLAIMER: The above information is provided for general educational information purposes only and should not be relied upon or used as the sole basis for making decisions without consulting primary, more accurate, more complete or more timely sources of information. The information contained has been gathered from publicly available current and reliable source material and is intended to provide general information to readers. NATIVE EXTRACTS Pty Ltd is not responsible for any reliance readers place on the material in this document and readers rely on the information in this document at their own risk and we recommend efficacy testing to be done on the finished product formulation to determine if it meets your specific target performance. However, this information is not intended to constitute an 'authoritative statement' under the National Industrial Chemical Notification and Assessment Scheme Australia and New Zealand rules and regulations. The range of compounds delivered in NATIVE EXTRACTS Pty Ltd products is only a guide as there will be fluctuations in the range available per batch as this is a natural product and reflects the nature of natural differences from harvest to harvest, source to source, batch to batch etc. NATIVE EXTRACTS Pty Ltd provides the information contained herein in good faith but makes no representation as to its comprehensiveness or accuracy. Individuals receiving this information must exercise their independent judgment in determining its appropriateness for a particular purpose. As the ordinary or otherwise use(s) of this product is outside the control of NATIVE EXTRACTS Pty Ltd, no representation or warranty, expressed or implied is made as to the effect(s) of such use(s), including damage or injury, or the results obtained. NATIVE EXTRACTS Pty Ltd expressly disclaims responsibility as to the ordinary or otherwise use(s). Furthermore, nothing herein should be considered as a recommendation by NATIVE EXTRACTS Pty Ltd as to the fitness for any use. The liability of NATIVE EXTRACTS Pty Ltd is limited to the value of the goods and does not include any consequential loss. NATIVE EXTRACTS Pty Ltd shall not be liable for any errors or delays in the content, or for any actions taken in reliance thereon. NATIVE EXTRACTS Pty Ltd shall not be responsible for any damage resulting from use of or reliance upon this information. The user of the product is solely responsible for compliance with all laws and regulations applying to the use of the products, including intellectual property rights of third parties.



# QUEEN GARNET

ADVANCED FRUIT AND VEGETABLES

Prunus salicina Fruit Extract

5. Taira J et al., Asian Pac J Trop Biomed., 2015; 5(10): 819-23

6. Chang LW et al., Food Chem Toxicol., 2011; 49(4):785-90

7. Kim SS et al., Asian Pac J Trop Biomed., 2013; 3(8):617-22

8. Carzillo et al., J Pharm Biomed Anal., 2015; 112:335-341

9. Estroff et al., J Pharm Biomed Anal., 2015; 112:335-341

10. Choi SJ et al., Int J Mol Med., 2016; 38(1): 357-63

11. Nagasawa T et al., Mol. Cell. Biochem., 2003; 249(1-2): 3-10

12. Nagasawa T et al., Mol. Cell. Biochem., 2003; 252(1-2): 141-7

13. Odetti PR et al., Diabetes, 1990; 39(7): 796-801

14. Tamazelli LC et al., Int J Pharm., 2018; 552(1-2): 401-6

15. Choquenot B et al., J Nat Prod., 2008; 71: 1117-8

16. Caporaso et al., J Food Compos Anal., 2008; 21(8): 589-98

17. Dobashi Y et al., J Photochem Photobiol Chem., 197(2-3):141-8

18. Jeon SE et al., Photodermatol Photomed Photobiol., 2003; 19(5): 254

19. Vayalil PK et al., J Invest Dermatol., 2004; 122(6): 1480-7

20. Madhan B et al., Int J Biol Macromol., 2007; 41(1): 16-22

21. Wu Q et al., Nutrients, 2014; 6(8): 3230-44

22. Rutter K et al., Int J Vitam Nutr Res., 2003; 73(6): 453-60

23. Afzar T et al., BMC Complement Altern Med., 2015; 15: 136

24. Jung MK et al., Arch Dermatol Res., 2012; 304(8): 655-63

25. Saric S et al., Antioxidants (Basel), 2017; 6(1): 2

26. Grzesik, M. et al., Food Chemistry., 2018; 241:480-492

27. Andres A. et al., Toxicology in Vitro., 2019; 54:1-9

28. He, J., et al., Medical science monitor: international medical journal of experimental and clinical research., 2018; 24: 8198-8206

29. Wojciech Koch, et al., Journal of AOAC INTERNATIONAL., 2017; 100(6): 1694-1699

30. Cosarca, A. et al., ABM., 2019, 2(1): 21-29

31. Bernal-Mercado, AT. et al., Molecules., 2018; 23(11):2813

32. Yang, W., et al., Korean Journal of Medicinal Crop Science., 2019; 27(2):143-150

33. Wang, L. et al., Environmental Pollution., 2019; 252:1318-1324

34. Wu, Q., et al., Cutaneous and Ocular Toxicology., 2021, 100.10.1007/s556957.2021.1879112

35. Chaniad, P. et al., PLOS ONE., 2020; 15(12)

36. Cheng, AW. et al., PLOS ONE., 2019; 14(5)

37. Fechtner, S., et al., Toxicology and Applied Pharmacology., 2017; 329:112-120

38. Isemura, M., Molecules., 2019; 24(3):528

**FLAVONOIDS**

1. Panche AN et al., J Nutr Sci., 2016; 5: e47

2. Kumar Sand Pandey AK, Scientific World Journal., 2013; 2013: 162750

**PHENOLICS**

1. Dziabl M et al., Int J Mol Sci., 2016; 17(2): 102

2. Soto ML et al., Cosmetics., 2015; 2(3): 2956

3. Samoylerko A et al., Antiox Redox Signal., 2013; 19(17): 2157-96

4. Liaudanskas M et al., ScientificWorldJournal., 2014; 2014: 306217

5. Alov P et al., Curr Top Med Chem., 2015; 15(2): 85-104

6. Andjelkovic M et al., Food Chem., 2005; 98(1): 23-31

7. Pastore S et al., Ann N Y Acad Sci., 2009; 1171: 305-13

8. Rhein LS and Fluhr JW, Aging Skin: Current and Future Therapeutic Strategies, Allured Business Media: Carol Stream, IL, USA, 2010; pp. 182- 184, 225-240

9. Arct J et al., J Appl Cosmetol., 2006; 21: 117-27

10. Czempik M et al., Sci Microb Pathog Commun Curr Res Technol Adv., 2011; 76: 39-47

11. Pinho E et al., Biomed Res Int., 2014; 2014: 814590

12. Zuk M et al., BMC Biotechnol., 2014; 14: 70

13. Anani K et al., J Ethnopharmacol., 2015; 164: 16-21

14. Vora J et al., Informatics in Medicine Unlocked., 2018; 13: 128-32

15. Koster ML, Ann N Y Acad Sci., 2009; 1170: 7-10

16. Dudonne S et al., Phytother Res., 2011; 25(5): 686-93

17. Dudonne S et al., J Agric Food Chem., 2011; 59(9): 4527-36

18. Wittenauer J et al., Fitoterapia., 2015; 101: 179-87

19. Fujii T et al., J Ethnopharmacol., 2008; 119(1): 53-7

20. Chowdhury A et al., Cell Biochem Funct., 2015; 33(7): 463-77

21. Quan T et al., J Invest Dermatol Symp Proc., 2009; 14(1): 20-4

22. Pluemsamran T et al., Photochem Photobiol., 2002; 88(4): 961-8

23. Kim HH et al., Arch Pharm Res., 2015; 38(1): 117

24. Liu WS et al., Mar Drugs., 2013; 11(6): 1899-908

25. Parvez S et al., Phytother Res., 2005; 20(11): 921-34

26. Brenner M and Hearing VJ, Photochem Photobiol., 2008; 84(3): 539-49

27. Mitari K et al., Planta Med., 2013; 79(1): 37-44

28. Rangkadilok N et al., Food Chem Toxicol., 2007; 45(2): 328-36

29. Zucca P et al., BMC Res Notes., 2015; 8: 562

30. Jeong JY et al., Molecules., 2015; 20(5): 8730-41

31. Almeida IF et al., J Photochem Photobiol B., 2015; 144: 28-34

32. Godic A et al., Oxid Med Cell Longev., 2014; 2014: 860479

33. Calvo-Castro L et al., Photochem Photobiol., 2013; 89(5): 1199-207

34. Svobodova A et al., J Dermatol Sci., 2009; 56(3): 196-204

35. Vostalova J et al., J Photochem Photobiol B., 2013; 128: 1-11

36. Svobodova A et al., Biomed Papers., 2003; 147(2): 137-45

37. Rahman M et al., Endocr Metab Immune Disord Drug Targets., 2012; 12(3): 287-302

38. Newman DJ and Cragg GM, J Nat Prod., 2007; 70(3): 461-77

39. Rigopoulos D et al., J Eur Acad Dermatol Venerol., 2005; 19(5): 564-8

40. Vijayalakshmi A et al., Int J ChemTech Res., 2011; 3: 320-7

41. Pothirath W et al., Pharm Biol., 2010; 48(2): 182-6

42. Huang WC et al., Food Funct., 2015; 6(8): 2550-60

43. Tanaka T., Curr Pharm Des., 2014; 20(6): 879-85

44. Kawai M et al., Allergol Int., 56(2): 113-23

45. Tanaka T et al., J Physiol Anthropol Appl Hum Sci., 2001; 20: 358-46

46. Park B J et al., Yonsei Med J., 2011; 52(3): 535-38

47. Sharma B et al., Indian J Microbiol., 2011; 51(2): 217-22

48. Furumoto T et al., Phytomedicine., 2014; 21(3): 247-53

49. Albaayit SF et al., Drug Des Dev Ther., 2015; 9: 3507-18

50. Nyanhongo GS et al., Eur J Pharm Biopharm., 2013; 83(3): 336-404

51. Bahramzamani R et al., Arch Dermatol Res., 2014; 306(7): 601-17

**CHLOROGENIC ACID (CGA)**

1. Rivelli DP et al., Photochem Photobiol., 2010; 86(5): 1005-7

2. Cheng D et al., J Inorg Biochem., 2019; 190: 113-20

3. Song J et al., Int J Mol Med., 2018; 41(2): 765-72

4. Li HR et al., Molecules., 2014; 19(9): 12940-8

5. Monghadam SE et al., Molecules., 2017; 22(9): 1501

6. Bagdas D et al., Naunyn Schmiedeberg Arch Pharmacol., 2014; 387(11): 1101-16

7. Bagdas D et al., Food Chem Toxicol., 2015; 81: 54-61

8. Vukelic E et al., Food Chem Toxicol., 2018; 121: 140-50

9. Liang N et al., Nutrients, 2016; 8(1)

10. Liu CC et al., Mol Med Rep., 2017; 16(2): 1369-75

11. Gasser P et al., Int J Cosmet Sci., 2011; 33(4): 366-70

12. Kim J et al., Arch Pharm Res., 2011; 34(3): 495-500

13. Martinez-Saez M et al., Food Chemistry., 2014; 150: 227-34

14. Alves GAD et al., PLoS One., 2019; 14(5): e0216301

15. Kitagawa S et al., Chem Pharm Bull., 2011; 59(6): 793-6

**ORGANIC ACIDS**

1. Kapuscinska A and Nowak I, Postepy Hig Med Dosw., 2015; 69: 3483

2. Tang SC and Yang JH, Molecules., 2018; 23(4): 863

3. Hon KL et al., Drugs Context., 2018; 7: 212530

4. Purnamawati S et al., Clin Med Res., 2017; 15(3-4): 75-87

5. Shahidi Fand Yeo J., Int J Mol Sci., 2018; 19(6): 1573

6. Ditto DK et al., Front Vet Sci., 2018; 5: 216

**SUGARS**

1. BeMiller JN., Carbohydrate Chemistry for Food Scientists 3rd Ed., 2019; AACC International, Chapter 1, 1-23

**AMINO ACID**

1. Akram M et al., J Med Plant Res., 2011; 5(17): 3997-4000

2. Scot R and Leonard S, Am J Clin Nut., 83(2): 500-7

3. Murakami H et al., Amino Acids., 2012; 42(6): 2894

4. Badiu DL et al., Protein J., 2010; 29(2): 81-92

5. Corsetti G et al., Acta Histochem., 2010; 112(5): 4937-6

6. Marcure R., Nature., 1900; 186: 886-7

7. Li P et al., Br J Nutr., 2007; 98(2): 237-52

8. Hashizume E., Clin Cosmet Investig Dermatol., 2013; 6: 499

9. Farris PK et al., J Drugs Dermatol., 2014; 13(1): 41-6

10. Oshimura E et al., J Cosmet Sci., 2017; 58(4): 347-57

11. Oshimura E and Ino M., J Cosmet Sci., 2015; suppl: S85-70

12. Leng N et al., Therapy., 2007; 4(1): 59-65

13. Hertel H et al., Hautarzt., 1999; 40(8): 490-5

14. Wood JM et al., FASEB J., 2003; 23(7): 2065-75

15. Rushton DH, Clin Exp Dermatol., 2002; 27(5): 336-404

16. Liu Y et al., Amino Acids., 2017; 49(8): 1277-91

17. Zhang ZY et al., Hypertension., 2018; 72(5): 1012-22

18. Dong Z et al., Am N Y Acad Sci., 2018; 1418(1): 44-55



## PRODUCT

NE Queen Garnet Cellular Extract Concentrate: ANE0616 Standard 1:10: ANE0616-10

## POTENTIAL APPLICATIONS

- Body clays, muds, scrubs
- Body moisturisers, milks, lotions
- Day, night creams, lotions
- Eye contour products, serums
- Face masks, scrubs
- Face strips, patches
- Generally, suits most cosmetic applications
- Hair, scalp, products, treatments
- Hand, feet creams
- Hand liquid soaps, towellettes, wipes
- Hyperpigmentation creams, serums
- Men, tennage products, serums
- Sun care, aftersun products

## POSSIBILITIES TO TRIAL

- Anti-aging care, antioxidant care
- Anti-glycation, barrier function
- Blemish care (acne prone skin)
- Collagen, elastin support
- Dark pigmented areas, exfoliation
- Fine lines and wrinkles
- Hair growth, maintenance, protection
- Hair maintenance, protection
- Inflammation reduction
- Reactive skin conditions (dermatitis, psoriasis, rosacea)
- Skin conditioning, firmness, hydration
- Skin radiance, luminosity
- Skin improvement, wound healing

Possibilities are based on compound information and not on the extract information. This information is for informational purposes only.

## PRODUCTION

Water/Glycerin Extract

Add at the end phase, <40°C

Ethanol option available

## PLANT PART: ORIGIN

Fruit : Australia

## USAGE GUIDE

Concentrate: 1.0 - 2.0%

Standard 1:10: 3.0 - 6.0% Wash

on/off formulations

24 months unopened from date of manufacture


## COMPLIANCE

INCI: Prunus salicina Fruit Extract

CAS: 1214268-59-3

EC: Not allocated

Cellular Extraction delivers natural water-soluble compounds from a species. To assist your ingredient research we identified publicly available information on these compounds from other sources and their potential uses. Information is not based on specific clinical trials of the Cellular Extract.



Forging the future of A-Beauty : POWERED BY NATURE

Copyright © NATIVE EXTRACTS Pty Ltd 2017. All rights reserved. DISCLAIMER: The above information is provided for general educational information purposes only and should not be relied upon or used as the sole basis for making decisions without consulting primary, more accurate, more complete or more timely sources of information. The information contained has been gathered from publicly available current and reliable source material and is intended to provide general information to readers. NATIVE EXTRACTS Pty Ltd is not responsible for any reliance readers place on the material in this document and readers rely on the information in this document at their own risk and we recommend efficacy testing to be done on the finished product formulation to determine if it meets your specific target performance. However, this information is not intended to constitute an "authentic statement" under the National Industrial Chemical Notification and Assessment Scheme Australia and New Zealand rules and regulations. The range of compounds delivered in NATIVE EXTRACTS Pty Ltd products is only a guide as there will be fluctuations in the range available per batch as this is a natural product and reflects the nature of natural differences from harvest to harvest, source to source, batch to batch etc.

NATIVE EXTRACTS Pty Ltd provides the information contained herein in good faith but makes no representation as to its comprehensiveness or accuracy. Individuals receiving this information must exercise their independent judgment in determining its appropriateness for a particular purpose. As the ordinary or otherwise use(s) of this product is outside the control of NATIVE EXTRACTS Pty Ltd, no representation or warranty, expressed or implied is made as to the effect(s) of such use(s), including damage or injury, or the results obtained. NATIVE EXTRACTS Pty Ltd expressly disclaims responsibility as to the ordinary or otherwise use(s). Furthermore, nothing herein should be considered as a recommendation by NATIVE EXTRACTS Pty Ltd as to the fitness for any use. The liability of NATIVE EXTRACTS Pty Ltd is limited to the value of the goods and does not include any consequential loss. NATIVE EXTRACTS Pty Ltd shall not be liable for any errors or delays in the content, or for any actions taken in reliance thereon.

NATIVE EXTRACTS Pty Ltd shall not be responsible for any damage resulting from use of or reliance upon this information. The user of the product is solely responsible for compliance with all laws and regulations applying to the use of the products, including intellectual property rights of third parties.