

CATARACT SURGERY STEPS/DATES - Dr's Crichton and Ford

Consent Video available on our website: www.glaucomacalgary.com

Questions: Call our office surgery Line at 403 245-3730, ext 2

1. ASCAN (1 to 12 months prior to Surgery):

This is a measurement of your eye needed to determine the type and power of the artificial lens that will be placed in your eye at surgery.

WHERE: Dr. Crichton and Ford's office: #102, 49 Richard Way SW, Calgary, AB

WHEN: MMM-DD-YYYY @

takes about 45 min, no dilation, a family member is welcome to join you to listen to discussion/options

1 WEEK PRIOR TO ASCAN: begin instilling artificial tears (eg. Refresh, Systane, Theoloz) 4 times per day in both eyes (continue any other drops already taking); please do not instill any drops 1 hour before exam.

- ◆ Artificial tears are available over the counter or alternatively, you can purchase a surgical bag from our office which contains preservative-free Theoloz-Duo drops as well as surgical tape and sunglasses.

- ◆ If you wear CONTACT LENSES, soft contacts must be removed 1 week prior, Rigid Gas Permeable contacts must be removed 2-3 weeks prior to the ASCAN.

MEDICAL FORM: This form is for YOU to complete IN FULL and return at AScan; please ensure you have seen your family doctor within the year prior to surgery. If you are having any health changes within the 3 months prior to your surgery date or a glaucoma surgery is added to your cataract surgery, please call our office to discuss as a more detailed medical exam may be needed by your family doctor.

- ◆ Please bring a translator if needed.

2. EYE DROPS (2 days before surgery):

see instruction sheet in package; the prescription with refills will be faxed to your pharmacy. These drops are not covered under seniors blue cross and cost ~\$130/eye without other health insurance. Bottles can be open for ~4-5 weeks. (Continue any other drops you already use for glaucoma, dry eye, etc)

3. SURGERY (time is tentative and may need to change):

. **EYE:** MMM-DD-YYYY Arrival Time:

. **EYE:** MMM-DD-YYYY Arrival Time:

WHERE: Holy Cross Surgical Services: Room 101, 2210 2nd ST SW, Calgary, Alberta

(see enclosed street and building maps), pay parking is available

- ◆ Nothing to eat after midnight the night prior (or a minimum of 8 hours), clear fluids permitted up to 3-6 hours prior.

- ◆ HOW LONG? You will be at the surgery center for approximately 1.5 to 2.5 hours

- ◆ **BRING current picture ID and your Alberta health care card**

FEES? IF you chose a specialized lens or do not have Alberta HealthCare, you must prepay part of the lens fee at our office and the balance at the surgery center on the day of surgery; you may pay with

- ◆ Debit/VISA/Mastercard although there is a small percentage added at the surgery center for credit cards (AHC pays for surgery with standard lens)

- ◆ **Somebody else must drive you home from surgery; taxi is acceptable if no other option and no sedation is given, you are not legal to drive for a minimum of 24 hours**

4. AFTER SURGERY Follow up appointments:

Will be booked automatically at our office closer to your surgery date, although you may choose to arrange with your local optometrist if preferred.

1 DAY after surgery:

@

~2-3weeks after surgery:

@

ASCAN/CATARACT DISCUSSION HIGHLIGHTS

PLEASE LISTEN TO THE VIDEOS ON OUR WEBSITE IF POSSIBLE

www.glaucomacalgary.com

CATARACT SURGERY:

A cataract is a clouding of the natural lens of your eye. Cataract surgery breaks up your natural lens (cataract) using a blade, ultrasound and fluid (not laser). This lens is then replaced with an implanted, artificial lens (IOL). The type/power of the IOL depends on the A-Scan measurements and your preferred outcome.

RISKS OF SURGERY:

There is approximately a 1% risk factor associated with cataract surgery, with the majority of the potential risks being treatable. Risks include but are not limited to; unexpected visual outcome requiring full-time use of glasses, inflammation, corneal irritation, increased light sensitivity, infection, retinal detachment, additional surgery needed and blindness. Some risks occur several weeks after surgery so please discuss with staff if you have any travel plans within 6 weeks after surgery. Advise staff if you have ever taken Flomax/Hytrin/Avodart or Risperdal/Invega medications OR if you have ever had eye surgery before.

LENS (IOL) RANGE TARGETS:

A. Distance Target:

Aims to eliminate distance glasses. Glasses will be required for any near activities within arm's length (ie. reading, looking at watch or in mirror, cutting fingernails), over the counter reading glasses are often helpful

B. Near Target:

Aims to eliminate reading glasses. Glasses will be required for any activities beyond arm's length (ie. driving, tv). This scenario is typically for people who currently remove distance glasses to read (and wish to keep this scenario)

C. Intermediate Target:

helpful for reading larger print at arm's length, computer monitors, etc but will likely need glasses for sharp distance vision (driving, reading guide on TV) as well as for detailed near work (ie. reading, looking at watch)

LENS (IOL) OPTIONS AVAILABLE:

(ANY FEES TO BE PAID PRIOR TO SURGERY, divided between office and surgery center)

1. "Standard"/Monofocal IOL:

This lens is paid by Alberta Health Care, it corrects one range only (see options above) and does not correct for astigmatism (part of your prescription). If you have significant astigmatism, you will still require glasses at all ranges to correct this.

2. Toric IOL: cost ~\$1,200.00 per eye

This lens aims to correct one range only (see options above) but also corrects astigmatism.

3. Extended Vision or Trifocal IOL: cost: ~\$2,300.00-\$2,500.00 per eye.

These lenses are newer technology and are designed to limit the need for glasses. A trifocal lens is likely to provide better uncorrected near vision but does have an increased risk of decreased quality of vision, particularly at night, with halos/glare around lights. An extended range lens does not provide as good near vision (will likely still need reading glasses for finer print) but does not have the increased risks of glare and halos.

NOTE: Individual healing responses vary, and we cannot guarantee the exact outcome of your surgery or requirement for glasses regardless of your lens choice

Dr. Crichton and Ford - Surgery Eye Drop Schedule

You will be prescribed the following eye drops for your surgery:

- Vigamox (Moxifloxacin):** 3x/day, start 2 days PRIOR to surgery, stop 1 week after surgery
Proensa (Bromfenac) 1x/day, Start 2 days PRIOR to surgery, stop 4 weeks after surgery
Durezol (Difluprednate): 2x/day, start the day AFTER surgery, stop 2 weeks after surgery
(Note: if Durezol is unavailable it will be substituted with Maxidex 4x/day)

** A Prescription will be faxed to your pharmacy, you do not need to pick up your eye drops until closer to your surgery date**

Note: once per day just needs to be consistent with time; twice per day ideally means 12 hours apart; three times per day means morning, mid-afternoon and bedtime; four times per day means morning, lunch, supper and bedtime.

Wait five (5) minutes between types of drops. Glaucoma drops (if you use any) carry on as per normal. If you have questions, call our office 403-245-3730 ext 2.

Start 2 Days BEFORE Surgery

	Day1	Day 2
VIGAMOX	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
PROLENSA	<input type="checkbox"/>	<input type="checkbox"/>
Notes:		

Day of Surgery

VIGAMOX	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
PROLENSA	<input type="checkbox"/>
*instill 1st doses before surgery, 2nd and 3rd doses not needed if have eye patch on	

1st Week After Surgery

	Day1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
VIGAMOX	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
PROLENSA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DUREZOL	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>
Notes:							

2nd Week After Surgery

	Day1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
PROLENSA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DUREZOL	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>
Notes:							


3rd Week After Surgery

	Day1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
PROLENSA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Notes:							

4th Week After Surgery

	Day1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
PROLENSA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Notes:							

HOLY CROSS SURGICAL CENTER




Holy Cross Surgical Services is located at:

Room 101-2210 2nd ST SW,
Calgary, Alberta
T2S 3C3

Paid parking is available.

Overall Site Plan

Our address:
2219 2nd Street SouthWest
Suite: 101



Holy Cross Surgical Services Patient Parking

Patient parking is marked by the grass

1. Enter off of 2nd Street by the crosswalk signals
2. Park in vacant stall
3. Parking fees apply; payable at any of four meters onsite
4. Enter at the Main entrance (Marked by the red "1") located to the right of the pharmacy.
5. The entrance is to the right after the full entrance vestibule.
6. Use the intercom to identify yourself with reception. Once identified, reception will open the door for you.



Frequently Asked Questions - Please Read Carefully

1. Do I take all my general health and eye medications before surgery?

A. Yes, take anything as you normally would, including any glaucoma and surgery eye drops, with the following exception:

- any medications recommended to take with food, such as Diamox or diabetic medications

2. Do I continue to take my blood thinner?

A. Continue any bloodthinners as you normally would (ensure staff are) with the following exceptions:

- If you take Coumadin/Warfarin, you should have your INR checked approx 1-2 weeks prior to surgery.

- NO BLOOD THINNERS ARE PERMITTED WITH GATT SURGERY. Discuss with staff if you are on bloodthinners and booked for a GATT surgery.

3. Can I eat and drink before surgery?

A. It is recommended to drink 3 cups of fluid (NO ALCOHOL) the evening prior. DO NOT EAT anything after midnight the night before surgery (Including gum/candies)

- You can drink up to 1 cup of CLEAR fluid (water, apple juice, sports drink, BLACK tea or coffee without sweeteners) up to 4-6 hours then nothing (6 hours for regional anesthetic, ie. needle or sedation).

4. What if I get sick before surgery?

A. You should contact the office (403-245-3730) immediately if you have an uncontrollable or new cough, fever, vomiting, possible eye infection, or sudden health changes.

5. Can I drive after surgery?

A. You are not allowed to drive yourself for 24 hours after surgery. It is preferred that a friend/family member drive you home after surgery.

- a taxi is permitted pending the type of freezing used and if no sedation is taken (ask staff for clarification.

- Public transit or walking by yourself is NOT permitted.

- You may ask a staff member at your follow up appointment if you are legal to drive or if you need to update your drivers license.

6. What can I expect on the day of surgery?

Ensure to wear loose-fitting clothing, nothing tight or restricting. Do not wear make-up or jewelry. Contact lenses will need to be removed

1. Check-in at Surgery Center Reception:

- you will need your current PICTURE ID and HEALTHCARE CARD You may need to sign a consent form for surgery

- Your driver may wait in the waiting room or return to pick up after surgery (~2hrs)

2. Anesthetist:

- The anesthetist will offer a mild sedative if you prefer and ONLY if you have someone who can be with you for the remainder to the day

- He/She will discuss your health, monitor your blood pressure and instill drops to "freeze" and dilate your eye.

- Rarely a needle is required to freeze your eye, this will be discussed with you prior to surgery if needed.

3. Surgery Room:

- You will need to lie down on the surgery table. The nurses will make you as comfortable as possible with know or neck support

- A drape will be placed over you leaving only your surgery eye exposed

- you will need to listen to instructions regarding where to look. If needed, a translator may accompany you in the surgery suite.

- the surgery will take about 15 minutes and is generally painless.

- a clear shield will be given to you to tape over the eye before sleeping for the first 1-2 weeks.

4. Recovery Room:

The nurse will take you back to the waiting room and ensure you are stable and comfortable prior to allowing you to go home.

7. Are there restrictions after surgery?

Immediately after surgery:

- You should rest quietly the day of surgery.
- If you leave the surgery center with a patch over your surgery eye, you may remove the shield (discard the gauze) the morning after surgery. You should keep the shield and purchase some medical tape to tape it back over your eye without gauze whenever sleeping for the first 1-2 weeks. If your skin is sensitive to tape, you may use elastic to make a "pirate patch" instead but this is not typically as secure.

1-2 weeks after surgery:

- For the first TWO WEEKS after surgery you should avoid rubbing your eye and any pools/hot tubs and dirty/dusty environments.
- You should avoid bending, lifting over 10lbs, straining or potentially strenuous activities including any sports.
- Try to avoid squeezing or closing eyes tightly.
- You may wash your hair after a few days but try to keep your head back for the first week when showering to minimize water/soap getting in eyes; do not squeeze your eyes shut.
- Sunglasses are recommended outside
- You should avoid wearing any eye makeup or having creams/lotions near eyes for one week after surgery.

8. What should I expect after surgery?

A. Your eye may be somewhat irritated the first 24 hours after surgery, however, extreme pain is not expected.

- Instilling artificial tears may help any mild irritation/itching.

Vision is often functional for the targeted range 24-48 hours after surgery (unless you also have a glaucoma surgery) however do not be alarmed if it is not as there are many factors involved in healing.

- Although your glasses will no longer be correct, you can continue to wear them if they help or you can have your optician remove the lens from in front of your surgery eye (this may not work well if you still have a high prescription in the other eye)
- Over the counter reading glasses may help with near work initially; it is recommended to wait about 5-6 weeks after surgery before updating glasses.

9. Can I travel after surgery?

A. Extended/international travel is not advised for the first six weeks after surgery (discuss with staff if you do have travel plans, travel insurance may be affected by surgery)

10. Are there any tips for instilling drops?

A. It is generally best to pull your lower lid down to form a "pouch", look up and instill one drop. Some people find it easier to lie down and look up.

- Be careful not to touch the tip of the bottle to your lid/lashes.
- ALWAYS WAIT 5 MINUTES BETWEEN DROPS
- Drops that are to be used multiple times a day need to be spaced out throughout the day.
- If you were using glaucoma drops before surgery, these should be continued.
- It is common for drops to sting for ~1 minute after instilling
- There are refill on your prescription for surgery drops

11. What if I still have questions?

A. You may call our office at 403-245-3730 ext 2 or 0, we will do our best to return any messages within 24 hours.

****Call IMMEDIATELY if there is a sudden drop in vision, severe pain or discharge or if you think you are noticing flashes of light or a large amount of floaters.****

If not during office hours, you may call Healthlink at 811, see your optometrist or go to the Rockyview Hospital Emergency