

FULL CORNEAL TRANSPLANT (PKP)

POST OPERATIVE

EYE DROP SCHEDULE WEEKS 1 & 2



Prednisolone/Pred Forte
(pink or white top)
One (1) drop every two (2) hours




Moxifloxacin/Vigamox
(beige/tan top)
One (1) drop every four (4) hours



Preservative Free Artificial Tears
(single-use vials)
One (1) drop every four (4) hours

8:00 am		
	9:00 am	
10:00 am		
		11:00am
12:00pm		
	1:00 pm	
2:00 pm		
		3:00 pm
4:00 pm		
	5:00 pm	
6:00pm		
		7:00 pm
8:00 pm		
	9:00 pm	



Muro-128 Ointment (sodium chloride)
(tube)
Quarter inch (1/4") each night at bedtime

See Reverse Side for Activity Limitations

ACTIVITY LIMITATIONS

1. Strict bedrest with face up positioning for 2 days after surgery. You may take 5-minute breaks every hour to move around, use the bathroom and eat meals. No watching TV or using computer unless it is bolted to the ceiling.
 2. Keep the eye covered at all times, except when placing drops. Wear either safety glasses or an eye shield during the daytime and wear an eye shield at night when sleeping.
 3. Do not lift anything heavier than 5 pounds.
 4. No getting the eye wet, keep it closed when showering.
 5. **DO NOT RUB THE EYE.**
 6. No swimming, no bending over, no strenuous exercise. Light activity only.
-

It is important to attend ALL your post-operative appointments.
Please call with any questions or concerns.

Remember the signs to be reported:

- R-Redness
- S-Sensitivity to Light
- V-Vision Decrease
- P-Pain