

REFER OR RELAX: RETINA

Steven Ferrucci, OD, FAAO
Chief, Optometry, Sepulveda VA
Professor, MBKU/SCCO

1

Lattice Degeneration

- 5-10% in General Population
- Found in 30% of all RD cases
- But, less than 1% of all lattice results in RD!!
—Byer NE. OPTH 1989. 0.7% over 10 years

2

Lattice Degeneration

- Circumferential oval lesions often with thin white blood vessels
- Pigment can vary
- Vitreous adhesion at borders
- Syneretic vitreous overlying the lesion itself
- Can have atrophic round holes without operculum typically towards end of lesions
— Occur up to 30% of the time

3

Lattice Degeneration

- Most common in Superior and inferior retina
— 2/3 cases from 5-7 or 11-1 o'clock
- Typical lesion size
— ½ to 2.5 DD in width
— 1-4 DD in length
- Average numbers of lesion per eye: 2
— Range: 1-19
- Bilateral in >>50% of cases

4

Lattice degeneration

- Most new cases discovered from 10-20 years of age
- May have hereditary component
- No apparent gender or race bias

5

Risk Factors

- Myopia > 3D, especially if < 30.
- Myopia > 6 D at any age
- Fellow eye has RD
- Family history of RD
- Symptoms
- Presence of traction
- High risk behavior

6

Follow up

- Lattice as only sign/symptom
 - Scleral depression
 - Pt ed.
 - RTC 1 year
- Lattice with symptoms of flashes/floaters
 - Reexamine q 1-2 mos
 - Or REFER if not comfortable
 - Repeat DFE/scleral depression
 - Pt ed

7

Follow up

- Lattice with holes but no risk factors
 - Scleral depression
 - Pt ed
 - Rtc 6 mos
 - Sooner if young myope, myope > 5 D, inferior holes, or adhesion
- Lattice with risk factors for RD
 - Consider retinal consult
- Lattice with breaks at margin of lesion
 - Consider retinal consult

8

Retinal Breaks

- Occur in 3 to 7% of adult population
- *Usually* asymptomatic
- 30-50% of *symptomatic* HST progress to detachment if untreated
- $\approx 5\%$ of *asymptomatic* HST progress to detachment if untreated
- Risk factors include lattice degeneration, high myopia, atrophic holes, aphakia/pseudophakia, and trauma

9

Horseshoe tears

- | | |
|---|--|
| <ul style="list-style-type: none"> • Common locations <ul style="list-style-type: none"> – Near lattice – Near pigment clumps – Near chorioretinal scars | <ul style="list-style-type: none"> • Worst locations <ul style="list-style-type: none"> – Superior – Near equator – Close to posterior pole |
|---|--|

10

Treatment

- **prompt treatment for *symptomatic* HST advised**
 - Reduces risk of RD to <5%
- Treatment of *asymptomatic* HST advised unless clear signs of chronicity
 - Even still most retinal MDs recommend treatment

11

Treatment

- Laser treatment is used to seal the break by creating adhesion between the retinal tissue and underlying RPE
- Provides barrier to continued enlargement from vitreo-retinal traction and prevents accumulation of subretinal fluid
- Adhesion present 24 hours after surgery, and strengthens over several days
 - Provides barrier to continued enlargement from vitreo-retinal traction and prevents accumulation of subretinal fluid
 - Adhesion present 24 hours after surgery, and strengthens over several days

12

Procedure

- Typically Topical or rarely retrobulbar anesthesia
- Entire lesion should be enclosed by at least 3 rows in a honeycomb pattern

13

Follow-up

RTC 1-2 weeks after treatment

Then 4-6 weeks

Then 3-6 mos

Then annually

If untreated, must be followed closely with proper pt education

AACD - Posterior vitreous detachment, retinal breaks and lattice degeneration Preferred Practice Pattern, 2019

14

Complications

- Few complications
 - inadequate burn intensity, causing ineffective adhesion
 - possible CNVM
 - intraretinal hemorrhage
 - vitreous hemorrhage
 - ERM formation
- More theoretical than actual
- Between 5-14 % of pts with initial break will develop additional breaks in future
 - Cataract surgery is risk factor for new breaks

15

Operculated holes

- Round, red hole with overlying free operculum attached to vitreous
 - Operculum often appears smaller than hole
- Minimal risk as no traction
- Treatment sometimes
 - High myopia
 - Aphakia
 - h/o RD in the fellow eye
 - Other factors

16

Atrophic Retinal Holes

- Small round, red hole w/o operculum
 - May have surrounding pigment
 - Occasional edema
- 2-3% of general population
- Most often in vitreous base
- Found in atrophic retina, perhaps 2° to vascular insufficiency

17

Atrophic Retinal Holes

- | | |
|--|--|
| <ul style="list-style-type: none"> • No traction <ul style="list-style-type: none"> – Minimal risk of detachment • Asymptomatic holes <ul style="list-style-type: none"> – Yearly – Pt ed • Asymptomatic with surrounding edema <ul style="list-style-type: none"> – Follow more closely | <ul style="list-style-type: none"> • Symptomatic <ul style="list-style-type: none"> – Consider consult • Other associated issues <ul style="list-style-type: none"> • As warranted • Rarely treated |
|--|--|

18

Treatment of Asymptomatic Lesions

Lesion	Phakic	High Myopia	RD etc in other eye
Atrophic hole	No	No	Rarely
Operculated hole	No	Rarely	Rarely
Lattice with or without hole	No	Rarely	Sometimes
Flap tear	Sometimes	Sometimes	Usually

19

Treatment of Symptomatic Lesions

Lesion	Treat
• Horseshoe tears	• Yes
• Operculated holes	• Rarely
• Atrophic holes	• No
• Lattice w/o holes	• No
• Lattice with holes	• Sometimes

20

Retinal Detachments

- Rule-of-thumb:
 - For macula off RD, want to get it repaired in same amount of time it has been off
 - So if off for 4 days, best to try repair within 4 days!
- Macula on RD is emergency!
 - Same day referral to retinal specialist
 - **Remind pt NPO until sees specialist in case same-day surgery**

21

Retinal Detachments

- Rhegmatogenous RD occur when liquefied vitreous fluid enters the sub-retinal space through a full-thickness retinal break.
- Occurs in 1/100,000 per yr
- Treatment options include scleral buckle, pars planar vitrectomy, and pneumatic retinopexy

22

Retinal Detachments

- Many factors go into selecting which procedure is best for patient
 - Phakic/pseudophakic
 - Location of tear
 - Size of tear
- Experience of retinal surgeon is essential!
 - Do your homework!

23

Retinoschisis

- Defined as splitting of the neurosensory retinal layers
 - Typically in outer plexiform layer
- variability: large, small, bullous, flat, bilateral, unilateral, progressive, non-progressive
- Two major types
 - Acquired
 - x-linked (XLR)

24

Acquired retinoschisis

- Usually benign and non-progressive
 - Myopic pts tend to be more progressive than hyperopic pts
- Asymptomatic, found on routine DFE, but may cause VF defect
- Incidence:
 - 3.9% in pts 60-80
- Most commonly affects inferotemporal retina
- Bilateral 33-82% of time

25

Schisis vs RD

- Schisis:
 - More translucent with visible vasculature
 - Less flexible
 - Well demarcated borders
 - Overall smoother appearance
 - Should have absolute VF defect vs relative with RD
- B scan/OCT can be helpful
 - OCT often difficult to image due to location

26

Retinoschisis

- Can have outer or inner wall breaks
 - Outer:
 - larger, often have ring of pigment
 - 11-24% of time
 - Inner:
 - smaller
 - Look like atrophic holes
 - Either associated with increased risk for detachment, so retinal consultation advised
 - Inner and outer together very dangerous
- if no holes, generally benign and can be monitored

27

Retinoschisis

- Very rare to have detachment into macula area
- Prophylactic Laser treatment has not been shown effective in most studies to halt progression
- Cataract surgery and PVD do not seem to have adverse effect
- If progresses to detachment, retinal surgery indicated
 - Only about 0.05% to 2.2% of cases
 - Typically respond poorly to surgery
- Most are benign and can be monitored yearly unless holes, enlargement, or symptoms

28

PVD

- Really no consensus
- Symptomatic PVD without retinal break
 - AOA: 1-2 weeks
 - **AAO: depending on symptoms, risk factors and clinical findings:**
 - 1-6 weeks
 - Then 6 mos to 1 year
 - Cleveland Clinic: 4-6 Weeks
 - Others: if no heme or other issues, very low risk so no need to see to back

29

PVD

- Floaters are typically most common symptom
 - Cobwebs
 - Flies
 - Hairs
- Flashes
 - Indicative of traction on retina, but not necessarily a tear or break

30

The Vitreous Humor

- Vitreous attached most firmly at
 - Macula
 - VMT
 - Vitreous base
 - Around optic nerve head
 - Weiss' Ring
 - Also, some traction on blood vessels
 - Vit heme

31

Incidence of PVD

Age	Incidence
>30	RARE
30-59	10%
60-69	27%
>70	63%
>80	75%

- 65%>65 HAVE A PVD

32

Incidence of PVD

- Incidence may be accelerated by
 - Myopia
 - Trauma
 - Prior vitreoretinal disease
 - Surgery
 - Inflammation
- Symmetrical 90% of the time
- Happens to second eye with 1-2 years

33

PVDs

- Good News:
 - Retinal Tears/Breaks *Relatively* uncommon
 - One study: only 7-15% of symptomatic PVDs have a retinal break
- Bad news:
 - 7-15% have a retinal break

34

CHRPE

- Lesions are almost always stable in size, but color may change.
 - Very rare instances of enlargement with time
- Typically asymptomatic, and found on routine exam, but large lesions have been shown to have VF defects

35

CHRPE

- Can also appear as multifocal CHRPE
 - From 3 to 30 lesions, 0.1 to 3.0 mm in size
- Benign, stationary and unilateral in 85% of the cases
- Often called bear tracks

36

Gardner's Syndrome

- Multifocal CHRPE have been associated with Gardner's Syndrome
 - AKA FAP: familial adenomatous polyposis
 - Familial condition of colonic polyps that may be precursor to colon cancer
 - However, these lesions are bilateral, have more irregular borders, and are often scattered throughout the fundus

37

CHRPE

- Differential includes nevi and choroidal melanoma
 - Nevi: nevi are rarely jet black and tend to have more indistinct borders
 - Melanomas tend to be greater than 2mm in thickness, where CHRPE are flat
- B-scan, serial photos and frequent monitoring of assistance

38

Nevus

- Common, benign tumor of the posterior fundus
- Typically slate –gray or brown in color, with somewhat indistinct borders
 - Often have overlying drusen, which signify chronicity of lesion
- Vary in size from 1/3 DD to as much as 7 DD
 - Flat or minimally elevated, < 2mm

39

Nevus

- Very common, with prevalence ranging from 0.2% up to 32% of patients
- More common in Caucasian population
- Asymptomatic, and usually found on routine exams
- Management consists of serial photography and frequent follow-up, with ultrasound if needed for more suspicious lesions

40

Nevus

- TFSOM: To Find Small Ocular Melanomas (1995)
 - T: Thickness: lesions > 2 mm
 - F: Fluid: any subretinal fluid suggestive of RD
 - S: Symptoms of photopsia or vision loss
 - O: Orange pigment overlying the lesion
 - M: Margin touching the optic nerve head
 - No factor= 3% risk of converting to melanoma in 5 yrs
 - 1 factor=8% risk
 - 2 or more factors =50% risk

41

Update 2019

- Incorporates imaging and re-evaluates risk factors
- TFSOM-DIM
 - To Find Small Ocular Melanomas Doing Imaging
 - T: Thickness > 2mm (US)
 - F: Fluid, subretinal (OCT)
 - S: Symptoms of vision loss (VA)
 - O: Orange pigment (FAF)
 - M: Melanoma Hollowness (US)
 - DIM: diameter > 5 mm (photos)

42

Update 2019

- Risk of converting to melanoma over 5 years
 - 0 factors: 1 % risk
 - 1 factor: 11%
 - 2 factors: 22 %
 - 3 factors: 34%
 - 4 factors: 51%
 - 5 factors: 55%
 - 6 factors: who knows?
- Bottom line: Increasing number of risk factors imparts greater risk for transformation

43

Update 2019

- M: Tumor Margin replaced with ultrasound
- S: Vision loss (VA < 20/50) rather than flashes/floaters
- Most important:
 - Thickness, Fluid, orange Pigment, Hollowness
- Least important:
 - Symptoms, Diameter

INSERT POLL QUESTION 7 and 8

44

Case 7

- Sent from Urgent care
- “61 y/o smoker newly diagnosed stage 4 lung CA with 5-6 week h/o loss of vision right eye”
- Back story: pt diagnosed ≈10 mos ago with stage 4 metastatic lung CA. refused all treatment and additional testing at that time
 - Saw herbalist. Drinking herbal tea and other homeopathic treatments

45

Choroidal Metastasis

- Most common site of metastasis to eye is choroid ≈ 88%
 - Iris 9%
 - Ciliary body 2%
- Most common primary sites
 - Men:
 - Lung 40%
 - GI 9%
 - Kidney 8%
 - Women
 - Breast 68%
 - Lung 12%

46

Choroidal Metastasis

- Typically yellow in color
- Often associated with subretinal fluid
- Solitary and unilateral or multiple and bilateral
- On ultrasound, have high internal reflectivity vs melanoma which has low internal reflectivity

47

Choroidal Metastasis

- 90% present with visual symptoms
 - Blurred vision 70%
 - Flashes /floaters 12%
 - Pain 7%
- Asymptomatic metastasis often detected in fellow eye
- Not uncommon to be asymptomatic

48

Choroidal Metastasis

• Differential diagnosis

- Choroidal amelanotic melanoma
- Choroidal amelanotic nevus
- Posterior scleritis
- Choroidal Hemangioma
- Choroidal Granuloma
- Choroidal osteoma

• Differential diagnosis

- Posterior Uveal Effusion syndrome
- VKH
- Central Serous Retinopathy
- Infectious lesions
- Organized subretinal hemorrhage
- Solitary idiopathic choroiditis
- Idiopathic Sclerochoroidal Calcification

49

Treatment

- Collaborative effort with oncologist/retinal specialist
 - Most pts have known primary cancer
 - If no known metastatic disease, PET CT needed to look for other metastasis as well as source
- If other metastasis, chemo alone can be effective
- Plaque radiotherapy for solitary metastases
- External beam radiation for bilateral and multifocal disease
- PDT for small lesions

50