



## PROTOCOL

### RECOMMENDED METHOD TO RESTART A STUCK FERMENTATION USING UVAFERM 43 RESTART


#### Prepare the Stuck Wine




#### Prepare the “Pied-de-cuve” (starter)



#### Yeast Rehydration



#### Incorporation of the “Pied-de-cuve”



For 1000 gals of stuck wine

#### Prepare the Stuck Wine

- Depending on analysis, address any potential spoilage organisms with SO<sub>2</sub> and/or lysozyme additions
- Add 1.5kg (3.3lb) Reskue and mix tank.
- Allow the tank to settle for 48 hours then rack off the settled lees.
- Adjust the temperature of the Reskue treated wine to 20–25°C (68–77°F).

#### Prepare the “Pied-de-cuve” (starter)

- Prepare the following:
  - 40 gallons water
  - 50 gallons post Reskue treated wine
  - 0.3kg (2/3 lb) Fermaid O
- Adjust Brix to 5°.

#### Yeast Rehydration

- Add 2kgs (4.4 lbs) of Go Ferm Protect Evolution in 10 gallons of water at 43°C (110°F).
- Cool solution to 40°C (104°F) and add 1.5kgs (3.3 lbs) of Uvaferm 43 Restart.
- Wait 20 minutes and slowly add rehydrated yeast to the “Pied-de-cuve.”
- Maintain temperature of 20–25°C (68–77°F).

#### Incorporation of the “Pied-de-cuve”

- Allow “Pied-de-cuve” to drop to 0° Brix and transfer immediately to the full volume of Reskue treated wine.
- Add 1.5kg (3.3lbs) of Fermaid O.
- Mix tank to homogenize.