Pain Relief You Can Feel. **Evidence You Can Trust.**

Clinically proven for perineal trauma, haemorrhoids, and post-surgical wound recovery.



- Punasundari, S., 2008. Perineal cold gel versus oral analgesic to relieve episiotomy wound pain: Part of Evidence-Based Nursing to Improve Patient Care.

- Punasundari, S., 2008. Perineal cold gel versus oral analgesic to relieve episiotomy wound pain: Part of Evidence-Based Nursing to Improve Patient Care.
 El-Saidy, T.M.K., Aboushady, R.M.-N. and Soliman, H.F.A., 2018. Effect of applying crushed ice gel pads on episiotomy pain and wound healing among postpartum primiparous women. International Journal of Nursing Didactics, 8(07), pp.19–29. DOI: 10.15520/ijnd.v8i07.2223
 Elmizadeh, K., Khezri, M.B., Pakniat, H., Pandamouz, V., Azh, N. and Molaverdikhani, S., 2024. The effect of uterine cooling during cesarean delivery on blood loss in patients undergoing cesarean section: A randomized controlled trial. Journal of Obstetrics, Gynecology and Cancer Research, 9(5), pp.487–491. DOI: 10.30699/jogcr.9.5.487
 East CE, Dorward ED, Whale RE, Liu J. Local cooling for relieving pain from perineal trauma sustained during childbirth. Cochrane Database Syst Rev. 2020 Oct 9;10(10):CD006304. doi: 10.1002/14651858.CD006304.pub4. PMID: 33034900; PMCID: PMC8094618.
 O J Traynor, A E Carter, Cryotherapy for advanced haemorrhoids: A prospective evaluation with 2-year follow-up, British Journal of Surgery, Volume 71, Issue 4, April 1984, Pages 287–289, https://doi.org/10.1002/bjs.1800710411



Disclaimer: For external use only. Discontinue use if irritation occurs and consult a healthcare professional. This product is not a substitute for medical advice or treatment. Always consult a healthcare provider and use under their guidance.

