
MAR / JUN 2003 No. 1-2

LEPIDOPTERA NEWS



Published by the Association for Tropical Lepidoptera, Inc.

P. O. Box 141210

Gainesville, FL 32614-1210, USA

Tel: (352) 373-5630 FAX: (352) 373-3249

www.troplep.org

Frequency: Quarterly (March, June, September, December) e-mail: troplep@aol.com

ISSN: 1062-6581

The Association for Tropical Lepidoptera, Inc., is a non-profit corporation for the study and conservation of tropical and subtropical Lepidoptera of world. Contributions are tax-deductible. Advertising is accepted.

ASSOCIATION FOR TROPICAL LEPIDOPTERA

BOARD OF DIRECTORS

Vitor O. Becker (Brazil) Don R. Davis (USA)

John B. Heppner (USA), Chair Gerardo Lamas (Peru)

Boyce A. Drummond, III (USA)

Olaf H. H. Mielke (Brazil) Eugene G. Munroe (Canada))

Peter J. Eliazar (USA Thomas C. Emmel (USA)

Jon D. Turner (USA)

Executive Director: J. B. Heppner

ADVISORY COUNCIL

Andrés O. Angulo (Chile)

Yutaka Arita (Japan)

George T. Austin (USA)

Manuel A. Balcázar L. (Mexico) Zsolt Bálint (Hungary)

Henry S. Barlow (Malaysia)

Dubi Benyamini (Israel)

Ronald Boender (USA)

Keith S. Brown Jr. (Brazil)

José A. Clavijo A. (Venezuela)

Charles V. Covell Jr. (USA)

U. Dall'Asta (Belgium)

Philip J. DeVries (USA)

Edward W. Diehl (Indonesia)

Julian P. Donahue (USA)

Ulf Eitschberger (Germany)

Eric Garraway (Jamaica)

Dale H. Habeck (USA)

Christoph Häuser (Germany) Lowell N. Harris (USA)

Hiroshi Inoue (Japan)

Daniel H. Janzen (USA)

Kurt Johnson (USA)

R. L. Kitching (Australia)

George O. Krizek (USA)

Tosio Kumata (Japan)

Jean-Francois Landry (Canada)

Torben B. Larsen (England)

Claude Lemaire (France)

Jorge Llorente B. (Mexico)

Martin Lödl (Austria)

Wolfram Mey (Germany) Kauri Mikkola (Finland)

Scott E. Miller (USA)

Joël Minet (France)

W. M. Neukirchen (Germany)

K. T. Park (South Korea)

Rod E. Parrott (Canada)

Amnuay Pinratana (Thailand)

Dalibor Povolný (Czech Rep.)

Jozef Razowski (Poland)

M. Alma Solis (USA)

Dieter Stüning (Germany)

Gerhard Tarmann (Austria)

Paul Thiaucourt (France) Jürgen H. R. Thiele (Germany)

Antonio Vives M. (Spain)

Hsiau-Yue Wang (Taiwan)

Per O. Wickman (Sweden)

Allen M. Young (USA)

MAR / JUN 2003

No. 1-2

CONTENTS

- HEPPNER: Tribal Classification of the Lepidoptera
- HEPPNER: Distribution Notes on Taiwan Moths, 2. Mimicia pseudolibatrix (Pyralidae: Pyralinae)
- 25 3. Pyrausta mystica (Pyralidae: Pyraustinae)
- 27 4. Genera Lixa and Tamraca (Pyralidae: Pyralinae)
- 29 5. Udea stigmatalis (Pyralidae: Pyraustinae)
- HEPPNER: Primitive Sedge Moths from New Zealand and Tasmania: Transfer of Proditrix and Relatives to Orthoteliinae (Glyphipterigidae)
- LAMAS: Additions to the Bibliography of Butterflies, in the Atlas of Neotropical Lepidoptera. VI. Comprising mostly Works Published in 2001
- HEPPNER: Neurobathra leafminer records on Jatropha new to Florida (Gracillariidae)

COVERS.- FRONT COVER: Argema mittrei (Guerín-Méneville), after Moucha (1962) (© 2005 ATL).

The Association for Tropical Lepidoptera, Inc., is a non-profit organization for the support of research on the biology, systematics, and conservation of tropical and subtropical Lepidoptera of the world. Funding for the Association helps to support research projects, field studies, and publications on tropical and subtropical Lepidoptera. The Association was organized in 1989 in Florida as a tax-exempt corporation under Section 501(c)3 of the IRS Code and is a publicly supported organization as defined in Sections 170(b)(1)(vi) and 509(a). Contributions are tax deductible in the United States.

LEPIDOPTERA NEWS (ISSN 1062-6581) is published quarterly (March, June, September and December) by the Association for Tropical Lepidoptera, Inc. ATL membership is \$90 per year, including the journals Lepidoptera News, Holarctic Lepidoptera, and Tropical Lepidoptera, plus newsletter. Membership is open to all persons interested in Lepidoptera. Membership applications and other information can be obtained from the ATL website (www.troplep.org), or by mail from the Executive Director, Dr. John B. Heppner, Association for Tropical Lepidoptera, P. O. Box 141210, Gainesville, FL 32614-1210. Tel: (352) 373-5630. FAX: (352) 373-3249.

LEPIDOPTERA NEWS, 2003 (1-2): 1-22 (2005)

TRIBAL CLASSIFICATION OF THE LEPIDOPTERA

J. B. HEPPNER 1

Florida State Collection of Arthropods, DPI, FDACS, P. O. Box 147100, Gainesville, Florida 32614, USA

ABSTRACT.—The worldwide biodiversity of the Lepidoptera is discussed in terms of current classification from higher categories down to the tribal and subtribal level. All 125 known families are treated, with 294 subfamilies listed.

KEY WORDS: biodiversity, cladistics, families, fauna, groups, subfamilies, subtribes, superfamilies, systematics, taxonomy, tribes, world.

The family classification of the Lepidoptera has been noted in several recent publications (see Heppner, 1992, 1996, 1998a, 1998b; Kristensen et al., 1999; Scoble, 1992), but the classification down to tribal level has not been uniformly proposed for many decades (Zerny and Beier, 1936-38). One reason for this is the great confusion of tribal names, their irregular usage — either as tribes, subtribes, or elevated to subfamilies and families - based on the varying ideas of classification of various specialists, and the vast number of genera yet to be adequately studied worldwide. In terms of Lepidoptera, one can point out the dozens of odd genera that were found to be in completely wrong families when the editors of the generic catalog project checked every genus, to the extent possible, to determine their correct family placement (Fletcher and Nye, 1975-91); and still the family placement of many genera remains uncertain. One can also note below the great number of tribal names, particularly in Geometridae and Noctuidae, many of which may prove to be only subtribes once further studied in relation to the world fauna of the groups (likewise, see the large number of subtribes listed for the tribe Geometrini).

We do not yet have enough comparative knowledge of all groups, on a world basis, to know if the name proliferation in some families represents real biodiversity, or only that some enthusiastic taxonomic splitters have been diligently naming every odd genus as another subtribe (which eventually may be relegated to some level below the subtribe). The difficulty for Lepidoptera studies is that few taxonomists work on a group worldwide, particularly for such huge families as Noctuidae, and, thus, cannot realistically judge tribal limits based only on their study of a small group in one region of the world. Thus, we have a large number of tribal names for European Geometridae and Noctuidae, for example, but with no adequate study to see which of these names may apply to related groups in tropical Africa or Asia. Many researchers fail (or refuse) to comprehend that to elevate tribes to subfamily level, or subfamilies to family level, without comparative study of all Lepidoptera, brings only chaos to the parameters of such groups: one cannot have a "family" for one group be equal to only a tribal level in some other family if one is trying to still have a logical and reasonable classification. The correllary to this is also that one cannot obtain a realistic classification based on current "political correctness" or general "global consensus" views, as some authors like to state, implying that the question is already solved. During the Middle Ages the general view was that the earth was flat: that was also a "global consensus" for that time, yet this did not represent reality. Consensus does not bring us a classification that resembles the reality in nature that we have, only facts and logic can do that.

The classification presented herein is the most up-to-date available, based on as uniform an adherence to parameters at each level (family, subfamily, and tribe) as possible at this time. It already varies somewhat from earlier reports at the subfamily level (see Heppner,

classification has been used only sporadically so far, but these names have been added herein to elucidate the classification, since many of these names may be found listed as tribes in other works. On the other hand, some families are not even subdivided into subfamilies, let alone tribal divisions (of course, there are also a few families that have only one known species: e.g., Heterobathmiidae and Carthaeidae). Thus far, 294 valid subfamilies are considered herein for the 125 known families of Lepidoptera in the world.

1992, 1996), since so much is still being discovered. A subtribe

ANALYSIS

Most specialists are not involved in worldwide studies in Lepidoptera, instead working only on regional faunas or particular families, and usually species only of one faunal region. A few exceptions exist: e.g., the work of Minet (1986, 1991), who has been doing extensive worldwide studies of different family groups and odd genera, although he splits off many more monobasic groups than may be useful. Likewise, there are my studies over the years (Heppner, 1991, 1996, 1998a,b, 2004), as also some global reviews like those of Common (1970, 1975) and the later revision of that (Nielsen and Common, 1991), plus summaries like those of Ehrlich (1960), Kristensen (1999), Munroe (1982), and Scoble (1992). Other studies, like those of Kuznetsov and Stekolnikov in a series of papers since 1976 (see Kuznetsov and Stekolnikov, 1978, for a basic summary), while in depth for most families, are nonetheless flawed because their studies involved only a single segment of adult morphology: i.e., exclusively the musculature of the male genitalia (as if this somehow would provide the answer to lepidopteran phylogeny better than other characters or what one finds in the larvae and pupae, etc.).

Among various studies, however, the tribal names found in many regional works are often the result of what one might call a "provincial" view (with no detriment intended), inasmuch as what are used as tribes in one region may actually only be equivalent to subtribes when compared with the world biodiversity of Lepidoptera. And, all this varies from specialist to specialist, as well. This is not to say that the classification published herein is the final word either: there is so much remaining to be discovered among many groups that we are a long way yet from a reasonably complete and stable classification down to tribal level. Even so, considerable work has been done in the last 30 years to elucidate the true placement of many odd genera that were left of uncertain status or else were completely misplaced (see Heppner, 1977; Nye et al., 1975-91), although much remains to be done. Families are now fairly stable, however, and even subfamilies in most families, but beyond that there is still much confusion in many of the larger families, particularly Geometridae and Noctuidae. At the subfamily level, one source of confusion is the continued elevation of odd tribes as subfamilies, then later perhaps again considered a tribe after more study, sometimes as part of a different subfamily than in the past, and so forth.

^{1.} Contribution No. 895, Entomology Section, Bur. Ent. Nema. Plant Path., Div. Plant Industry, Florida Dept. Agric. & Consumer Serv.

Family-level changes at this time mainly involve the different views among specialists as to what constitutes a family: some split every odd group as a new family (the motto seems to be: better a new family than to try to work at it more and alter the parameters of an existing family), while others are at the opposite extreme and combine all kinds of groups together as larger families (remember that Linnaeus had only 3 "families" for all Lepidoptera, or at most 10, depending on how you view his groupings). There is also a major divergence of opinion on higher categories (families, subfamilies, tribes), between those who study only butterflies and those who only study moths. For example, some butterfly specialists once had almost every butterfly family as a superfamily in itself, rather than being all under the superfamily Papilionoidea: this is contrary to what the superfamily level is considered to be in the remainder of the Lepidoptera.

Since evolution is not always at the level where, when all the Lepidoptera are compared among each other, such that one has all category levels (from species on up) with distinct boundaries, we thus have intergrades as well: e.g., sibling-species for species that are just evolving into distinct species (and these are usually retained as a single species in our classifications at this time), so also it is reasonable to expect there to be "sibling" subfamilies and "sibling" families. This is part of what I view as the problem behind many disagreements as to what level various groups are: e.g., the controversy between Pyralidae and Crambidae, or more recently between such odd groups as Oenosandrinae in Notodontidae, etc. While I alter family definitions to incorporate odd tropical groups into existing families, if feasible, others tend to immediately split off such groups as separate entities, as already noted above. What may also be involved are what was just noted, a kind of "sibling" status. To engage this problem in a classification, I believe, the use of the "group" category below the family level is a useful option. Thus, we have the groups Crambinina and Pyralinina, since I do not consider the single main character dividing these two groups sufficient to warrant reverting to their split into two families, as was already done 100 years ago. Likewise, we have the case of the "families" Oenosandridae and Thaumetopoeidae, split off from Notodontidae: this case has been an on-going controversy (other than the newly segregated Oenosandridae) for decades. Yet again, I believe, this is a case of what could be termed "sibling" families, and thus, in the classification below these are listed as group names: Oenosandrinina, Thaumetopoeinina, and Notodontinina. Group names in a classification demonstrate that there is some level of difference between the groups greater than evident as subfamilies, yet not enough to warrant separate families. Of course, many cladists do not recognize this and elevate every odd group as a new higher taxon. Group names are also used in Arctiidae, Noctuidae, and Nymphalidae, and possibly more such arrangements may be needed in other families that are very diverse and may have "sibling" groupings; e.g., the Oecophoridae, where some authors have again elevated all the subfamilies as separate families (Kristensen et al., 1999). Other candidates for the possible addition of "group" classifications may be Gracillariidae, Tineidae, and Zygaenidae, all with odd subfamilies that have sometimes been considered separate families (viz. Phyllocnistidae, for what now is subfamily Phyllocnistinae in Gracillariidae, and Hieroxestidae, for what now is subfamily Hieroxestinae in Tineidae). However, as with subgenera and subtribes, part of the problem therein is that many specialists simply refuse to use such relatively non-traditional categories, preferring instead to elevate all unusual taxons to the next higher level (yet among other insect orders subgroup usage is normal).

Until most specialists can agree on what is a family (subfamily, tribe, etc.) there will remain controversies as to what level any particular group should be classified as. For example, a recent work created a large number of "families" in Gelechioidea for odd groups that previously were maintained in Oecophoridae (see Hodges, 1999), this being a reversion back to a similar classification before

these groups were placed within Oecophoridae. So, back and forth from generation to generation, every 30 years or so, the family levels change when a new group of workers looks at the same problems and thinks of doing it differently again — vive la différence!, as it were — although in this particular case it was the result of an extreme interpretation of a cladistic analysis of the Gelechioidea that skewed (or rather, skewered) the results. Likewise, there are a larger number of monobasic superfamilies that have been proposed, simply because some specialists split off each odd family rather than redefine groupings to accomodate the odd, often tropical, species being studied. Remember that our classification was originally based mainly on European genera, and has since 1758 been repeatedly altered as more and more of the tropical diversity was discovered.

Current so-called "phylogenetic" cladistics (or Hennigian cladistics) is not any different or "better" than other recent taxonomic methodologies (e.g., actual phylogenetic taxonomy, as from Mayr and Simpson), being as inherently biased as past methodologies, although many specialists using cladistics generally do not acknowledge any subjectivity: their results, however, are as dependent on the characters chosen (and the computer program used when a computer analysis is involved) as older conclusions, and these results must also be interpreted (see the note above about Gelechioidea). Current methodologies are superior, however, at least in the one aspect that more and more specialists are using a wider array of more fundamental characters in family-level classification and their studies are more precise in assessments of synapomorphies, which is an improvement over what was done earlier in the past century; the only problem is that these same specialists may include wildly varying characters as family-level characters in trying to come to their conclusions (e.g., using various genitalic characters for family-level studies of inter-relationships is foregone to result in possibly erroneous conclusions simply because genitalic characters are mainly species specific, or generic-level, and will only rarely offer some characteristic that can be validly applied to family-level studies). In many families we also have varying classifications because one specialist uses only adult characters, while another makes use of larval and pupal characters, which often produces very different results: rather, all these characters should be integrated into an anlysis, to thus provide a more solid framework for our classification. And finally, many of the cladists adhere to strict monophyly in groups (which actually is the most ideal), yet will then subjectively determine where to draw the line on what the family level is, or where the tribal level is, and so forth.

Another problem in achieving a stable classification is the lack of characters of the immature stages of a great many groups, even entire families. If we do not have the array of characters offered by larval and pupal morphology, then relationships can often be very inconclusive. Here is an area of study that students should make note of, since searching for the immature stages and biologies of families (and other unique groups), where these remain unknown, can bring great advances in our classification. For example, in Noctuidae the members of the highly unusual Papuan subfamily Cocytiinae are completely unknown biologically, yet the larvae and pupae could offer many characters to show where their true affinities lie, since the adults are strange day-flying moths (only 2 species are known) that appear more like small sphingids than owlet moths. There are many other cases like this: entire families such as Neopseustidae, Eriocottidae, and Ratardidae, among others, are completely unknown biologically (see Heppner (1998) for a listing of other biologically unknown families and subgroups).

Thus, to reiterate, the classification below is based on a worldwide study of the groups involved, not just a regional understanding or bias. Various works have been consulted (see Heppner (1998) for listings of references), as well as personal studies over the last 30 years. While monophyly is the standard applied to groupings, differential rates of evolution have also split groups that could

otherwise be combined (e.g., for Sesiidae, and particularly for Zygaenoidea and Cossoidea), thus, a reasoned approach has been used overall.

The classification below is most tentative at the tribal level in Geometridae and Noctuidae. As noted previously, regional workers have often elevated what are subtribes to tribal level, such that more names have been proposed than for the biodiversity that is involved. Even herein, the tribal listings of some families may be too extensive (e.g., in Geometridae), and many of the names may be dropped to subtribe level in the future, following more extensive study and comparative review of each group. However, the large biodiversity of some families may require large numbers of tribes. And, if some genus is unique, it may require a separate tribe (or other category) even if monobasic (witness Carthaeidae, with a single unique species that requires its own family). Many families are so vast, however, that progress is slow and will require the work of many specialists willing to take on the task. Phylogenetic arrangements (from primitive to advanced taxa) also vary regionally and by specialist in some cases: as an example, most subfamily and tribal arrangements in the family Geometridae vary between works and specialists of the New World fauna (e.g., Ennominae as a more primitive group in the family, thus, near the beginning of the listing) and the Old World fauna (e.g., Ennominae at the end, or as the most advanced geometrids). As noted previously, many worldwide studies are still needed but cannot be realistically advantageous until the immature stages are known for all families, thus giving those crucial character sets their due consideration and possibly resolving the conflicts of classifications as we see in families like the Geometridae

TRIBAL NAMES

Synonyms are not included in the classification listing below, but can be found in the index (1,984 names), with notation as to what the correct name is for each taxon. Some of the synonym names have been used by others simply due to the erroneous view that once an older generic name is added to a group then the higher level taxon must adopt the name of the oldest included genus: the new Zoological Code (ICZN, 1999) specifically advises against this kind of instability, thus the more stable names in the literature are used. One of the best examples of this is the continued erroneous usage of Roeslerstammiidae instead of the long-used name Amphitheridae (in this case merely because an older European genus was added to the formerly tropical family and the reviser, not being familiar enough with the Code, simply thought the family name had to then be changed; and oddly, other specialists who know the Code better continue to perpetuate this error). The name Roeslerstammiidae is also invalid, since the original usage did not even include any of the species currently in this group, thus the senior name is Amphitheridae, as well as being the only valid name (Heppner, 2005b).

Another erroneous usage is Acentropinae, instead of Nymphulinae, in Pyralidae; again because an older generic name is involved. In this case the ICZN has even made a formal ruling for the use of Acentropinae, but many specialists will ignore this since a strange ICZN commission vote was involved (the commissioners actually voted "against" Acentropinae but due to bureaucratic rules of the ICZN their lack of a quorum of voters automatically favored the ruling for Acentropinae!). Should one question the sanity of ICZN rules that leave it to some bureaucratic solution, rather than the logical choice that even Commission members voted for based on the Code and the facts? One also has the recent use of Macariini instead of Semiothisini in Geometridae, simply because the genus Macaria is now added to the Semiothisa group as a valid, but older genus (again erroneous usage). Likewise, there is the recent unconscionable use of Strepsimaninae (Kristensen et al., 1999) instead of the well-known Hypenodinae in Noctuidae (in this case, because the older genus Strepsimanes has now been added to

Hypenodinae; again erroneous usage). Even more egregious is the new use of Erebiinae instead of the well-known Catocalinae by some researchers, again for supposed use of the oldest name. Contrarily, the use of Nepticulidae instead of Stigmellidae, is precisely the same type of case (Stigmella is the senior generic name over Nepticula), yet various researchers strangely in this case retain the use of the supposed junior name, Nepticulidae. If workers fail to read the Code and continue to erroneously change higher taxon names whenever an older genus is added to a group, or when it suits them, then we will continue to have a lot of name changes and cause continued instability and confusion.

DISCUSSIONS

Primitive Moths

Among those families considered the most primitive of lepidopterans, there is little controversy that Micropterigidae are the most basal group remaining among living Lepidoptera. Recent studies by Davis (1999), plus those of Minet (1986, 1991) and others, have clarified the basal divisions of the order Lepidoptera such that, barring any discoveries of other primitive taxa in remote tropical regions, we probably have a clear picture of the evolutionary development from Zeugloptera to Ditrysia, the remainder of lepidopteran families. The only recent addition is of a surprising new monotrysian family added to Nepticulina, the Andesianidae (Davis, 2003), for three Valdivian species previously misplaced among Cossidae.

Incurvarioidea

This group of families is still in need of further study, since some specialists continue elevating almost all the subfamilies as separate families, thus we have some listing Crinopterygidae, Lamproniidae, and Prodoxidae as separate families (Davis, 1999). As noted in the introduction, some specialists always make new families rather than incorporate new character states within the framework of an existing family, thus multiplying families right and left. As listed herein, the 5 families show cohesive characters with their included subfamilies.

Tineoidea

Among Tineoidea, the series category is used to demarcate Tineiformes and Gracillariiformes; some specialists, however, make these into two separate superfamilies. The basal area of the superfamily now has the Acrolophidae segregated as a separate family from the remaining Tineidae, the acrolophids being considered the most primitive group of the superfamily.

In Tineidae, there remains considerable study yet needed on a worldwide basis to clarify all the levels and relationships among the subfamilies and tribes listed, many of which have also been thought of as separate families in the past. Also, the presumed most advanced group, the Hieroxestinae, have recently again been presented as a separate family by some (Davis and Robinson, 1999).

Psychidae likewise need study for their subfamily classification, with some specialists still using family names for Lypusidae, so the relationships among all the psychid taxa remain uncertain.

The remaining families, comprising the Gracillariiformes, are all small except for the main leafminer group, the Gracillariidae. Amphitheridae is still unjustifiably called Roeslerstammiidae by many specialists, mainly just following each other without further question from a beginning point where someone incorrectly changed the family name.

In Gracillariidae, the Phyllocnistinae have often been viewed as a separate family, but cohesive characters do not warrant that. As in other such cases, to elevate this group as a separate family would require every subfamily of Lepidoptera to also be elevated to family-level, if one tried to treat all groups with the same arguments as used for having Phyllocnistidae as a separate family, thus giving us some 300 "families" worldwide.

Gelechioidea

As already noted in the introduction, this superfamily has reverted from few families to many families, back and forth, about once every generation of specialists. Currently, we have 11 families, but some specialists now want again to elevate practically every subfamily, especially among the Oecophoridae, as a distinct family (Hodges, 1999): this can easily be done, but the definition we would have of such "families" would require the elevation of all other subfamilies as well, as already mentioned above. The most recent modification of this view is by Kaila (2004), with at least 16 "families" indicated in his cladology, and others would push the total to over 20 families. Note again, that to elevate all these groups in Gelechioidea to family-level would require equal usage for the rest of the order, or about 300 "families" in total.

Is it useful to classify the Lepidoptera with 300 families or with 125 families? Should every odd group simply by called a "family" now? In Coleoptera, various specialists are also now engaged in a widespread "splitting" campaign, and thus have about 10 families of weevils for what previously was considered one family, for example. If one wishes, one can continue on such a path to the point where we have a "family" for every odd group, or as in birds where one has a different genus for almost every other species: obviously every species is different but we need some common sense in where to draw the lines for higher taxon groupings. The denegration of what a "genus" is has also been accelerating in recent decades as much as has the family concept: one can look at bird classification, where many "genera" now contain only a few or a single species. Some taxonomists have actually enunciated a belief that a genus should automatically not have more than 5 species, as if this were a fact of nature! Of course, these persons have never studied the genus Coleophora with over 1,000 species worldwide, among other such large genera! As noted earlier, the reduction in parameters for genera and families has been progressing since the time of Linnaeus, but in the last 100 years or so we have made a critical analysis of what a family should be and what a genus should be, notwithstanding taxonomically insane trends among higher animals, like bird classification.

In Oecophoridae, notwithstanding the above comments, there still remains a need for considerable study. The group is so vast worldwide that it has been difficult to handle by any one person, and most of the past classifications have mainly involved the European fauna instead of dealing with the vast tropical fauna. Ian Common (formerly of CSIRO, Canberra, Australia) is one of the few now trying to make some sense of the many genera of oecophorids from Australia, where most of the biodiversity appears to reside. The current subfamilies have been listed in Oecophoridae for some time and only recently has there again been an effort to split them all into separate families (Hodges, 1999). New developments have shown, however, that some odd groups, like the so-called Symmocidae and Holcopogonidae, previously either separate families or placed in Blastobasidae, may well best go in Oecophoridae, along with Glyphidocerini, in Autostichinae. The Oecophoridae are extremely diverse, yet have cohesive character traits, thus they are a very large family that has become very biodiverse, yet compared to other family characters across all Lepidoptera, are rightfully together as one family. Unfortunately, some researchers cannot comprehend (or refuse to do so) a "family" that is diverse and large (yet cohesive nonetheless), much as was noted above for the splitting of genera based on a maximum size of 5 species, as if this were not some arbitrary choice but some law of nature.

Lecithoceridae were long thought to belong in Gelechiidae, but are now considered more primitive and related more to Oecophoridae and Elachistidae.

Elachistidae are considered herein as a smaller family than some other researchers view it. A recent classification has relegated almost all Oecophoridae to an enlarged concept of Elachistidae (Minet, 1990), or even as a separate superfamily (Kuznetsov and

Stekolnikov, 1984), but this elevates all other groups, such that Gelechioidea would have to be a "suborder" of Lepidoptera, as Kaila (1999) also noted): clearly there is considerable elevation going on for different groups in the minds of some specialists, or splitting, as already alluded to in the introduction. The true Elachistidae (sensu stricto) have recently been studied on a worldwide basis by Kaila (1999), but he did not arrange the genera into a higher classification other than what his consensus tree showed: based on his studies, the classification used herein makes use of appropriate subfamily and tribal names but more study is needed.

Copromorphoidea

This small superfamily has only 4 included families: Copromorphidae, Alucitidae, Carposinidae, and Epermeniidae. The cohesive characters of this grouping have already been well established (Heppner, 1977). Recent attempts to continue splitting up this group into separate superfamilies for Epermeniidae and Alucitidae are based on erroneous conclusions of Minet (1986, 1991) as to what should be a superfamily, as well as continuing the taxonomic fiction that Alucitidae are somehow related to Pterophoridae (as was done even 150 years ago because both have split wings) (Dugdale *et al.*, 1999b).

Yponomeutoidea

This superfamily comprises 9 families at present: Ochsenheimeriidae, Glyphipterigidae, Plutellidae, Attevidae, Yponomeutidae, Argyresthiidae, Lyonetiidae, Acrolepiidae, and Heliodinidae. My studies had first considered Glyphipterigidae to belong in Copromorphoidea (Heppner, 1977), based especially on larval characters, but the family evidently best represents basal elements within Yponomeutoidea, as does Ochsenheimeriidae (earlier transferred from Tineoidea). Recent study has shown some primitive genera from New Zealand and Australia, thought to be Plutellidae, to actually belong to basal groups within Glyphipterigidae, subfamily Orthoteliinae (Heppner, 2005a). While splitters run rampant in other superfamilies, they oddly continue maintaining Attevidae as a subfamily in Yponomeutidae, yet segregate Ypsolophinae as a separate family from Plutellidae (Dugdale *et al.*, 1999a).

Immoidea

This is a monobasic superfamily, with only the family Immidae (Heppner, 1977, 1982). The family is pantropical, but mostly from tropical Asia. A recent addition is a subfamily classification due to the addition of Prodidactinae (new status), currently a *nomen nudum* published in a paper by Epstein and Brown (2003) that is invalid by nomenclatural rules of the zoological code (Art. 8.6 and 9.7): the name is listed herein in case the authors validate the name. Prodidactinae were defined as a family (Prodidactidae) but the characters indicate that the single genus is only of subfamily status in Immidae (another case where authors would rather have a "new family" than appropriately modify an existing family to accomodate a strange new tropical member).

Pyraloidea

The superfamily includes three families: Hyblaeidae, Thyrididae, and Pyralidae. Cladist splitters have split this into no less than three separate "superfamilies" (two subfamilies of Thyrididae are even split off by some as separate "families"). Larval characters in particular show superfamily cohesion for all the included groups.

One large controversy is the recent splitting of Pyralidae by elevating to family level its two groupings, as Crambidae and Pyralidae, whereby almost all genera commonly viewed as "pyralids" are now in Crambidae, leaving Pyralidae with only the minor subfamilies and the phycitines (Munroe and Solis, 1999). Minet (1991) even considered all the major subgroups as separate "families", thus with 5 families (including also "Acentropidae," Pyraustidae, and Phycitidae). There is nothing new in all this: all

has already been done over 100 years ago when another splitting craze in taxonomy was loose on the planet. Again, the cohesiveness of Pyralidae, with all the groupings included as subfamilies, has been viewed as logical for the family already for many decades, until recently. A recent paper has refined characters of the subfamilies of "Crambidae" (Solis and Maes, 2002), although it is based only on adult characters.

When one internal morphological character was discovered that did indeed segregate the pyralids from the crambines, Munroe (1972) chose the logical view to use a subgroup classification, as is often done in other insect orders where the family may be evolving but still retains enough other characters to remain as one family. Cladist purists cannot follow this (and others often refuse to use unconventional subgroups), so see only splitting as a solution, although if they simply lowered their "family" level cut on their cladology trees (where they arbitrarily have chosen a certain level of similarity to be called a subfamily or family), they would come to the conclusion also of having only one family Pyralidae. Using the "group" category below family level, of course, shows that there is a distinction between the two groups but also shows that this is not large enough a difference overall to segregate them as separate families.

Pterophoroidea

This superfamily includes the plume moths, Pterophoridae, plus two odd related tropical families, Tineodidae and Oxychirotidae (Heppner, 1997, 1998c). Some researchers now include Oxychirotidae within Tineodidae (Dugdale *et al.*, 1999b), but wing venation and larval biologies differ.

Sesioidea

This superfamily has been defined with modern character studies already since 1981 (Heppner and Duckworth, 1981), now including 4 families: Brachodidae, Sesiidae, Urodidae, and Choreutidae. The cladist splitters do not agree and split off Urodidae and Choreutidae as separate superfamilies, mainly based on studies of only adult characters and idiosyncratic levels of cohesion of groups to exclude (Edwards *et al.*, 1999). There has been some recent restructuring of the tribes within Sesiidae, plus the addition of tribes for odd tropical genera.

Zygaenoidea

The old concept of this superfamily has been shown to be heterogeneous, mainly due to groupings thought to be similar due to some traits (e.g., larval shape as slug caterpillars) that now are known to only indicate convergent evolution (Heppner, 1998b, 2005c). While the main family, Zygaenidae, is clear enough, the odd smaller groups that are related have always caused confusion. Thus, the families that actually are cossoid, and not zygaenoid, were moved to Cossoidea (Heppner, 1998b). Researchers focused only on apparent similarities (e.g., the similar larvae), fail to agree with this and forget that convergent evolution can provide such results as similar larvae (which nonetheless show differences as well). Families remaining in Zygaenoidea include Heterogynidae, Zygaenidae, Himantopteridae, Lacturidae, Somabrachyidae, and Megalopygidae. Some researchers continue attempts to split Zygaenidae into several "families" and others include Somabrachyidae within Megalopygidae, or split out groups like Aididae (Epstein et al., 1999).

Cossoidea

This superfamily is actually the basal stem of the Cossina group of superfamilies that includes all the remaining Lepidoptera up to Noctuidae, being fundamentally evolved from different ancestors than where the Zygaenoidea ended up from. Cossoidea now include 6 families: Cossidae, Dudgeoneidae, Metarbelidae, Cyclotornidae, Epipyropidae, and Dalceridae. Again, the slug-like larvae of

Dalceridae have some researchers confused into believing they should be part of Zygaenoidea, while the family morphology shows an evolution derived from cossoid ancestors (Heppner, 1998b).

Castnioidea

This monobasic superfamily represents a basal, rather primitive family, the Castniidae, that appear to best be placed as a separate superfamily, although clearly also derived from cossoid-like ancestors.

Tortricoidea

We now place the Tortricidae in their own superfamily. The actual placement of the group in the overall classification varies among researchers, since some place the group near the Tineoidea (Horak, 1999), but in relation to all Lepidoptera they appear best placed in our linear format after the Castnioidea and at the end of what are termed the "microlepidoptera." The true layout of families in our classification would be better seen in three dimensions, whereby Tortricoidea would be seen to retain some primitive features like Tineoidea but otherwise advanced enough to place the group 3-dimensionally to the side in another lineage from tineoid-like ancestors; but, a paper layout requires placement linearly in only two dimensions (Heppner, 1998b).

Calliduloidea

This small Old World group includes only 3 families: Ratardidae, Pterothysanidae, and Callidulidae. All are day-fliers, and most callidulids even fly and perch like butterflies.

Uranioidea

This superfamily includes 5 families: Epicopeiidae, Apoprogonidae, Sematuridae, Uraniidae, and Epiplemidae. Most species are Old World, but at least Epiplemidae are found in all regions. Cladists have recently moved Epiplemidae into Uraniidae (Minet and Scoble, 1999), in a reversal from their usual splitting, but the groups are different enough to remain as separate families, particularly in terms of their larvae. The superfamily is rather closely related to Calliduloidea.

Geometroidea

The family Geometridae, being such a biodiverse family (21,150 spp. worldwide), still requires considerable study to conform all the higher category (subfamily and tribe) names by region, such that what a tribe is in Europe has the same meaning as a tribe in North America or Asia: currently, there is wide disagreement on this matter between North American specialists and European specialists. What we have now are numerous tribal names that remain of uncertain status: are they tribes or only subtribes? For example, most of the group names in the subfamily Geometrinae have been treated as tribes in the past, but recently Holloway (1996) probably correctly noted that there should be only two tribes and all remaining names ought to be at most subtribes. Likewise, to follow the use of subtribes in Geometrinae, many of the Ennominae tribes probably only warrant subtribal status as well, but which do and how they should be arranged remains unclear: e.g., one can identify about 6 groupings of tribes in Ennominae, so perhaps these should be tribes and the remaining names only subtribes. Holloway (1996) also now uses an enlarged concept of Boarmiini that would put many other groups as subtribes therein, even more than listed herein. Consequently, the listing of tribes in Geometridae herein is still tentative. The listed arrangement of the tribes within each subfamily is also tentative, since this also varies regionally: most classifications in Europe have Ourapterygini, for example, near the middle of Ennominae, while in North America it is towards the end of the subfamily. One needs to also determine which groups are considered the most advanced: most specialists now agree that such subfamilies as Archiearinae and Oenochrominae are more primitive geometrids,

but some place Ennominae as advanced while others place Larentiinae as advanced. The classification of Geometridae herein is an attempt to show the relationships of the various tribes and subfamilies, but it is far from final and mainly represents revision of conventional arrangements by specialists that often do not like to use subtribes (thus, all such names are tribes in their mind). Additionally, more odd tropical groups are discerned all the time, so much work remains to be done with this large family.

Hedylidae are retained in Geometroidea herein. While they may indeed represent the evolutionary development that gave us the butterflies, their base characters retain them more to Geometroidea than they do to Papilionoidea (Heppner, 1998b). Obviously, others disagree and put them as part of the butterflies, or else make the usual split as a separate superfamily, Hedyloidea (Minet and Scoble, 1999).

Papilionoidea

The butterfly workers have often had considerably differing classifications over the years, from the 7 families we now list to many more and even added superfamilies for the larger families of butterflies. Again, putting the classification in the perspective of world lepidopteran biodiversity, one finds that most butterflies are not that different from each other (e.g., nymphalids are all in one family and not in their own superfamily) in relation to how moths are classified. To maintain a uniform family level one cannot have more splitting among the butterflies. Some researchers now include Libytheidae within the Nymphalidae (Ackery et al., 1999), but this is unrealistic given their great morphological differences. Hesperiidae are split by some into their own superfamily, but this is not necessary when one uses the "series" subgrouping to show that their divergence from higher butterflies is not so great that a separate superfamily is needed. Riodinidae follows the new Neotropical catalog (Lamas et al., 2004).

Drepanoidea

A small superfamily, including 4 families: Axiidae, Thyatiridae, Cyclidiidae, and Drepanidae. Most species are Old World.

Bombycoidea

This well-known grouping of moths now includes 13 families: Carthaeidae, Eupterotidae, Apatelodidae, Bombycidae, Mimallonidae, Anthelidae, and Lasiocampidae in the series Bombyciformes, and Endromidae, Lemoniidae, Brahmaeidae, Oxytenidae, Cercophanidae, and Saturniidae in the series Saturniiformes (Heppner, 1998b). Sphingidae are generally excluded to their own superfamily, but some workers keep them within Bombycoidea (Lemaire and Minet, 1999).

Sphingoidea

The single family Sphingidae comprises this monobasic superfamily of well-known moths, the hawk moths (or sphinx moths). Some alterations of the tribes have been made in recent years (Lemaire and Minet, 1999).

Noctuoidea

Almost a third of all Lepidoptera belong to Noctuoidea, including 6 families: Notodontidae, Dioptidae, Thyretidae, Lymantriidae, Arctiidae, and Noctuidae (Heppner, 1991, 1998b). Various odd members of the families have more recently been segregated as several "families" of their own, again based mainly on cladist artificial rules on what level a researcher considers a family to be defined. In our classification we use subgroupings to show what differences there are, rather than do unnecessary splitting: this is especially needed in Noctuidae, where recent splitting has not only given us Aganaidae and Herminiidae, but now also include a complete splitting of the Noctuidae to include also the "families" Erebiidae (even the Catocalinae name is changed!), Nolidae,

Acronictidae, Pantheidae, etc. Tribes are noted in the listing below but subtribes are mostly not included, as has been done to a profuse extent for Europe (Beck, 1999b), since the rest of the world fauna has not had equivalent treatment and many of the European subtribal names may be simply excessive splitting (many of these European subtribes are monobasic, representing only a single genus) (Kitching and Rawlins, 1999).

The splitting trend in noctuoids has been progressing for the past 20 years or so (see Kitching, 1984; Speidel et al., 1996), what cladists consider a correct trend due to "rigorous application of cladistic principles" rather than past vague group definitions, as Speidel et al. (1996) put it: the problem with this is as noted already, the results are from whatever their computer program version and cladology analysis gives them, which is not necessarily correct but based only on these current parameters and their subjective analysis. This is no different than past work, just with more precise characters and the use of computers, but the result is still subjective as always and dependent on how experienced a taxonomist the compiler actually is — the problem is that some cladists fervently believe their methodology is akin to reassembling the phylogenetic lineage from God himself (or DNA) and the results cannot thus be questioned other than possibly with slight refinements using another computer program, all the while forgetting that it is they (not God) who decide where to the draw the line on what a family is or what a genus is.

Again, the noctuid workers cannot split odd subgroups away in Noctuidae as "families" without also having all other Lepidoptera subfamilies elevated to family-level in their view of what a family constitutes. There has to be concordance throughout the order for all higher groupings. Of course, if one can agree with a classification where a "family" is defined one way among micros and another way among butterflies and another way among noctuoids, then perhaps the mad hatters tea party can resume and all logic for an overall classification of all the Lepidoptera can be ignored. Is it rational to have three different definitions of what a "family" is within the classification of one order of insects? Clearly, each level of a classification must be maintained in as rational a cohesive form throughout an order as possible, given all our human subjectivities in any case and based on what the evolution of the order shows us, not cladist fantasies based on a narrow view of only one group. In Noctuidae, for example, various attempts have continued for the past 100 years to resurrect "Nolidae" about every generation as new taxonomists studied the group: they otherwise conform well to the overall concept of the family Noctuidae. The Nolinae (or Nolidae) may represent an offshoot, also supported by larval characters, but this shows more a development from lithosiine Arctiidae than from Noctuidae (Beck, 1999a), however what may be involved here is again a basal element remaining in the family, much as with some of the oecophorids that are herein retained within Oecophoridae. Holloway (2003) has expanded this segregation to include other subfamilies within an enlarged Nolidae, based on his views of the group for the Asian tropics. Likewise, the use of non-conventional subgroups also retains such odd tropical groups as Aganainae within the Noctuidae, so Nolinae may have to be handled similarly if the evidence mounts.

CONCLUSIONS

Since few researchers study all Lepidoptera from all regions of the world, the specialist studying a single group, perhaps only for one faunal region, quickly can find any odd tropical group to require a "new" family. To classify our Lepidoptera one can only resort to uniformity of what each level of the classification is to mean, at least insofar as is humanly possible. The pure cladist view is to simply draw a line on their computer trees for each level, but this methodology, while seemingly simple and clear, is not so when we are dealing with nature: too many character variables negate this from being perfectly applicable to each group for the entire order.

Likewise, the methodology of making all characters in a computer analysis equal (i.e., some hair on a leg being equal in value to having 6 legs, for example) does not work in nature, since some characters are more fundamental than others (e.g., the 6 legs versus only 4 legs!). The true phylogenetic taxonomist, on the other hand, modifies each grouping to conform not only to an overall level within the classification but also to the evolution of the group involved; thus, there are sometimes shifts needed on the cladist trees on where the line is for different groups, yet in such a way that the definition of the level chosen is not compromised. What is meant here is that different "rates" of evolution need to be included in a valid classification, not simply just morphology, and this is something the cladologists do not fully acknowledge (thus, the errors, as for example, in not seeing convergent evolution in the case of some Zygaenoidea and Cossoidea, or to see that Brachodidae are related to Sesiidae even though the Sesiidae advanced very rapidly from their brachodid-like ancestors). To split families, like Noctuidae, into many "families" and yet be ignorant of what a family is among primitive moths, or among butterflies, will not give us a stable or logical classification: we need conformity of level throughout the Lepidoptera in order to arrive at a reasonable classification that approximates their evolution and phylogeny.

We can proceed with splitting and end up with 300 or so "families" in Lepidoptera - the beetle people are nearly their already, in their cladist frenzy of splitting (there are even 10 or more "families" just for weevils alone!) - but a rational classification will keep groups about equal throughout, from micros to noctuids. For the past 100 years or so the concept of what a "family" is in Lepidoptera has been relatively stable, until recently. If we want, we can have 300 families of Lepidoptera. Is this reasonable? Is this useful? These are questions we need to consider when continuing the efforts to split off odd groups from families that they can otherwise reside in. One can actually modify family definitions: this is something that some Lepidoptera researchers seem unaware of, thinking instead that somehow the original description of a family is sacred and that any group falling outside this first definition requires a "new" family to be named. We have refined family definitions since the time of Linnaeus, so modifying a family definition to include some odd group, if it really can be included, can be done. What we find is simply that many researchers refuse to change families and instead name a new family. This is more political than rational or scientific. There are more synonyms to family names than there are valid families, as many researchers have learned over the years. Thus, further erosion of what constitutes a "family" should be discouraged, otherwise to be rational in our classifications then all lepidopteran subfamilies would need to be elevated to family level: you cannot have "Nolidae" or "Danaidae" without also elevating all other groups currently at the subfamily level. You either have logic derived from what the evolution of the Lepidoptera shows us to be different families, their characters, or you have a mad hatters party.

Finally, what you see in the listing below of families, subfamilies, tribes, and in some cases subtribes, is not the final word on Lepidoptera classification. Only recently was a valid new family added to the list for an odd tropical group (Andesianidae), so as more remote areas of the world are further surveyed it is reasonable to assume that a few more odd groups may yet be discovered. Likewise, further knowledge and study of some obscure groups, as for example those families where we as yet do not know the larvae, may yet modify our classification in unforeseen ways. Nonetheless, the present classification, I believe, best represents a uniform and rational view of lepidopteran evolution and phylogeny as we currently know it to be, and with concordant values of higher groupings throughout the order Lepidoptera.

Trying to make a classification based not on fact, but on a "consensus" view of the facts is nonsensical, just like it was when the "global consensus" was that the earth was flat: the "global

consensus" can be wrong, as was discovered in 1492 (and already earlier by the Chinese). Many other erroneous "consensus" views abound even today, such as the current view that the Incas built the monuments in Cuzco, Peru (when they themselves told the early Spanish explorers that the buildings were already there and very old when their people first came into the valley of Cuzco and made use of them, yet the consensus archaeologists maintain the fiction that these are ruins built by the Incas); or, the view that the Egyptians somehow built the great pyramid and sphinx at Giza, Egypt (when we know that the early Egyptians did not have the mathematical or engineering skill to make such perfect buildings, while their own pyramid attempts were all inferior copies and the sphinx was even stated by them to be there from ancient days long before their people were even civilized, yet consensus egyptologists continue the fiction that these monuments were built by the Egyptians). Plate tectonics was also fully dismissed as complete fantasy 45 years ago by the then-prevalent "global consensus" of geologists, yet we all know that this is a proven fact and is now taught in every high school in the world as a basic tenent of modern geology. Even crustal displacement is gaining ground as a possibly valid theory of earth movements. Let the archaeologists and egyptologists continue with their heads in the sand, but we need not follow their woodenheaded examples in our work on Lepidoptera classification.

Classification is not a political game of consensus building, but a reasoned study of the species we are dealing with as they are in nature and as they evolved to the present day, whereby we then attempt to outline this evolution into some kind of order showing the relationships among the included groups. It also is not based on a single methodology, such as computer generated cladologies using the most recent cladistics computer program that uses all possible characters equally (no matter how nonsensical that may be), and which is then deemed the only methodology possible for classification. As noted earlier, no matter what classification is used, the classification outline should even be seen 3-dimensionally to better show relative evolution in different lineages, but we have to organize it 2-dimensionally to get it linearly on paper.

ACKNOWLEDGMENTS

Several specialists were consulted on the tribal listing, in particular M. J. Scoble (Natural History Museum, London) was most helpful for Geometridae. Any remaining errors, however, and the final arrangement in the list, are my own responsibility; in particular so, because no two specialists could agree on such a listing anyway. Common names for families are from Heppner (1998a), plus one change in the name for Lyonetiidae.

REFERENCES

Ackery, P. R., R. de Jong, and R. I. Vane-Wright

1999. The butterflies: Hedyloidea, Hesperioidea and Papilionoidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 263-300. Berlin: W. de Gruyter.

Beck, H.

1999a. Systematik allgemein (Überlegungen aufgrund der Erfarhrungen mit der Systematik der Noctuidae), larvale Charakterisierungen von Gruppen, Schlüssel. In Die Larven der Europäischen Noctuidae. Revision der Systematik der Noctuidae, 1:47-89. Marktleuthen: U. Eitschberger. 4 v. (Herbipoliana, 5).

1999b. Systematische Liste der Noctuidae Europas (Lepidoptera: Noctuidae). In Die Larven der Europäischen Noctuidae. Revision der Systematik der Noctuidae, 1:783-83. Marktleuthen: U. Eitschberger. 4 v. (Herbipoliana. 5).

Common, I. F. B.

1970. Lepidoptera (moths and butterflies). In The Insects of Australia. A Textbook for Students and Research Workers, 765-866. Melbourne: Melbourne Univ. Pr.

 Evolution and classification of the Lepidoptera. Ann. Rev. Ent. (Palo Alto), 20:183-203.

Davis, D. R.

1999. The monotrysian Heteroneura. In Kristensen, N. P. (ed.), Handbuch der

Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 65-90. Berlin: W. de Gruyter.

Davis, D. R., and G. S. Robinson

1999. The Tineoidea and Gracillarioidea. In Kristensen, N. P. (ed.), Hand-buch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 91-117. Berlin: W. de Gruyter.

Dugdale, J. S., N. P. Kristensen, G. S. Robinson, and M. J. Scoble

1999a. The Yponomeutoidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 119-130. Berlin: W. de Gruyter.

1999b. The smaller microlepidoptera-grade superfamilies. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 217-232. Berlin: W. de Gruyter.

Edwards, E. D., P. Gentili, M. Horak, N. P. Kristensen, and E. S. Nielsen 1999. The cossoid/sesioid assemblage. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 181-197. Berlin: W. de Gruyter.

Ehrlich, P. R.

8

 Lepidoptera. In McGraw-Hill Encyclopedia of Science and Technology, 459-473. New York: McGraw-Hill.

Epstein, M. E., and J. W. Brown

2003. Early stages of the enigmatic *Prodidactis mystica* (Meyrick), with comments on its new family assignment (Lepidoptera: Prodidactidae). www.mapress.com/zootaxa (*Zootaxa*, 247:1-16).

Epstein, M. E., H. Geertsema, C. M. Naumann, and G. M. Tarmann

1999. The Zygaenoidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie.
35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 159-180. Berlin: W. de Gruyter.

Fletcher, D. S., and I. W. B. Nye (eds.)

1975-91. The Generic Names of Moths of the World. London: Brit. Mus. (Nat. Hist.). 6 v. [alternating editor listing by volume]

Heppner, J. B.

1977. The status of the Glyphipterigidae and a reassessment of relationships in yponomeutoid families and ditrysian superfamilies. J. Lepid. Soc. (Los Angeles), 31:124-134.

1982. Review of the family Immidae with a world checklist (Lepidoptera: Immoidea). Entomography (Sacramento), 1:257-279.

 Faunal Regions and the Diversity of Lepidoptera. Gainesville: Assoc. Trop. Lepid. 85pp (Trop. Lepid., 2, Suppl. 1).

1992. Introduction. In Lepidoptera of Taiwan. Vol. 1. Part 2: Checklist, ixlix. Gainesville: Assoc. Trop. Lepid. 276pp.

1996. Keys to families of Lepidoptera. Gainesville: Assoc. Trop. Lepid. 28pp (Trop. Lepid., 4, Suppl. 3).

1997. Oxychirotidae. In Lepidopterorum Catalogus, (n.s.). Fasc. 62. Gainesville: Assoc. Trop. Lepid. 8pp.

1998a. Revised family list for Lepidoptera. Lepid. News (Gainesville), 1998 (3):57-62.

1998b. Classification of Lepidoptera. Part 1. Introduction. Gainesville: Assoc. Trop. Lepid. 148pp (Holarctic Lepid., 5, Suppl. 1).

1998c. Tineodidae. In *Lepidopterorum Catalogus*, (n.s.). Fasc. 61. Gainesville: Assoc. Trop. Lepid. 8pp.

2004. Butterflies and moths (Lepidoptera). In J. L. Capinera (ed.), Encyclopedia of Entomology, 387-428. Dordrecht: Kluwer Acad. Publ. 3 v.

2005a. Primitive sedge moths from New Zealand and Tasmania: transfer of Proditrix and relatives to Orthoteliinae (Lepidoptera: Glyphipterigidae). Lepid. News (Gainesville), 2003:31-42. (2005)

2005b. Review of Amphitheridae (Lepidoptera: Tineoidea). Tinea (Tokyo) (Moriuti memorial suppl.): [in press].

2005c. Classification of Lepidoptera. Part 2. Microlepidoptera: Micropterigoidea to Tortricoidea. Gainesville: Assoc. Trop. Lepid. Pp. 149-330. (Holarctic Lepid. (Gainesville), 12, Suppl. 1).

Hodges, R. W.

1999. The Gelechioidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 131-158. Berlin: W. de Gruyter.

Holloway, J. D.

 Family Nolidae. In *The Moths of Borneo*, 18. Kuala Lumpur: Southdene. 279pp, 104 + 10 pl.

Horak, M.

1999. The Tortricoidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 199-215. Berlin: W. de Gruyter.

International Commission on Zoological Nomenclature (ICZN)

1999. International Code of Zoological Nomenclature. (4th ed.). London: Internatl. Trust Zool. Nomencl. 306pp. Kaila, L.

1999. Phylogeny and classification of the Elachistidae s.s. (Lepidoptera: Gelechioidea). Syst. Ent. (London), 24:139-169.

2004. Phylogeny of the superfamily Gelechioidea (Lepidoptera: Ditrysia): an exemplar approach. *Cladistics* (London), 20:303-340.

Kitching, I. J.

1984. An historical review of the higher classification of the Noctuidae (Lepidoptera). *Brit. Mus. (Nat. Hist.), Ent.* (London), 49:154-234.

Kitching, I. J., and J. E. Rawlins

1999. The Noctuoidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 355-401. Berlin: W. de Gruyter.

Kristensen, N. P. (ed.)

1999. Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography. Berlin: W. de Gruyter. 487pp.

Kuznetsov, V. I., and A. A. Stekolnikov

1978. The system and evolution of infraorders in the Lepidoptera (Micropter-igomorpha-Papilionomorpha) treated on the basis of functional morphology of the genitalia. Ent. Obozr. (Moscow), 57:870-890. [in Russian (Engl. transl. 1979. Ent. Rev. (Washington), 57:601-615)]

Lamas, G. (ed.), C. J. Callaghan, M. M. Casagrande, O. H. H. Mielke, T. W. Pyrcz, R. K. Robbins, and A. L. Viloria

2004. Checklist: Part 4A. Hesperioidea - Papilionoidea. In J. B. Heppner (ed.), Atlas of Neotropical Lepidoptera. Gainesville: Assoc. Trop. Lepid. 439pp.

Lemaire, C., and J. Minet

1999. The Bombycoidea and their relatives. In Kristensen, N. P. (ed.), Hand-buch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 321-353. Berlin: W. de Gruyter.

Minet, J.

1986. Ébauche d'une classification moderne de l'ordre des Lépidoptères. Alexanor (Paris), 14:291-313.

1990. Remaniement partiel de la classification des Gelechioidea, essentiellement en fonction de caractères pré-imaginaux (Lepidoptera Ditrysia). Alexanor (Paris), 16:239-255 (1989).

1991. Tentative reconstruction of the ditrysian phylogeny (Lepidoptera: Glossata). Ent. Scand. (Copenhagen), 22:69-95.

1999. The Axioidea and Calliduloidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 256-261. Berlin: W. de Gruyter.

Minet, J., and M. J. Scoble

1999. The drepanoid/geometroid assemblage. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 301-320. Berlin: W. de Gruyter.

Munroe, E. G.

1982. Lepidoptera. In S. P. Parker (ed.), Synopsis and Classification of Living Organisms, 2:612-651. New York: McGraw-Hill. 2 v.

Munroe, E. G., and M. A. Solis

1999. The Pyraloidea. In Kristensen, N. P. (ed.), Handbuch der Zoologie. 35. Lepidoptera, Moths and Butterflies. Vol. 1: Evolution, Systematics, and Biogeography, 233-256. Berlin: W. de Gruyter.

Nielsen, E. S., and I. F. B. Common

1991. Lepidoptera (moths and butterflies). In The Insects of Australia. A Textbook for Students and Research Workers (2nd ed.), 817-915. Melbourne: Melbourne Univ. Pr.

Scoble, M. J.

1992. The Lepidoptera: Form, Function and Diversity. Oxford: Oxford Univ. Pr. 404pp.

Solis, M. A., and K. V. N. Maes

2002. Preliminary phylogenetic analysis of the subfamilies of Crambidae (Pyraloidea Lepidoptera). Belg. J. Ent. (Brussels) 4:53-95.

Speidel, , W., H. Fänger, and C. M. Naumann

1996. The phylogeny of the Noctuidae (Lepidoptera). *Syst. Ent.* (London), 21: 219-251.

Zerny, H., and M. Beier

1936-38. Ordnung der Pterygogenen: Lepidoptera = Schmetterlinge. *In* Kükenthal (ed.), *Handbuch der Zoologie*, 4(2). *Insecta*, 3:1554-1728. Berlin: W. de Gryter.

LEPIDOPTERA Suborder ZEUGLOPTERA

MICROPTERIGOIDEA

1. MICROPTERIGIDAE - Mandibulate Archaic Moths Micropteriginae Sabatincinae

Suborder AGLOSSATA

AGATHIPHAGOIDEA

2. AGATHIPHAGIDAE - Kauri Moths

Suborder HETEROBATHMIINA

HETEROBATHMIOIDEA

3. HETEROBATHMIIDAE - Valdivian Archaic Moths

Suborder GLOSSATA

Cohort DACNONYPHA Infraorder DACNONYPHA

ERIOCRANIOIDEA

4. ERIOCRANIIDAE - Sparkling Archaic Sun Moths

5. ACANTHOPTEROCTETIDAE - Archaic Sun Moths Acanthopteroctetinae Catapteriginae

Infraorder LOPHOCORONINA

LOPHOCORONOIDEA

6. LOPHOCORONIDAE - Australian Archaic Sun Moths

Cohort MYOGLOSSATA Subcohort MYOGLOSSATA

Infraorder NEOPSEUSTINA

NEOPSEUSTOIDEA

7. NEOPSEUSTIDAE - Archaic Bell Moths

Subcohort NEOLEPIDOPTERA

Infraorder EXOPORIA

MNESARCHAEOIDEA

8. MNESARCHAEIDAE - New Zealand Primitive Moths

HEPIALOIDEA

9. NEOTHEORIDAE - Amazonian Primitive Ghost Moths

10. ANOMOSETIDAE - Australian Primitive Ghost Moths

11. PROTOTHEORIDAE - African Primitive Ghost

Moths

12. HEPIALIDAE - Ghost Moths

13. PALAEOSETIDAE - Miniature Ghost Moths

Infraorder HETERONEURA

Division MONOTRYSIA

Section NEPTICULINA

ANDESIANOIDEA

14. ANDESIANIDAE - Valdivian Forest Moths

NEPTICULOIDEA

15. NEPTICULIDAE - Pygmy Moths

Pectinivalvinae

Nepticulinae

Nepticulini

Trifurculini

16. OPOSTEGIDAE - Eye-Cap Moths

TISCHERIOIDEA

17. TISCHERIIDAE - Trumpet Leafminer Moths

PALAEPHATOIDEA

18. PALAEPHATIDAE - Gondwanaland Moths

Section INCURVARIINA

INCURVARIOIDEA

19. INCURVARIIDAE - Leafcutter Moths

Incurvariinae

Crinopteryginae

20. CECIDOSIDAE - Gall Moths

21. PRODOXIDAE - Yucca Moths

Lamproniinae

Prodoxinae

22. ADELIDAE - Longhorned Fairy Moths

Nematopogoninae

Adelinae

23. HELIOZELIDAE - Shield Bearer Moths

Division DITRYSIA

Section TINEINA

Subsection TINEINA

TINEOIDEA

Series Tineiformes

24. ACROLOPHIDAE - Tube Moths

Amydriinae

Acrolophinae

25. TINEIDAE - Fungus Moths

Euplocaminae

Myrmecozelinae

Harmacloninae

Meessiinae

Infurcitineini

Meessiini

Dryadaulinae

Scardiinae

Nemapogoninae

Tineinae

Setomorphinae

Perissomasticinae

Hapsiferinae

Glyphidocerini

Oeciini Schistonoeini

Hieroxestinae Xyloryctinae Erechthiinae Stenomatinae Siloscinae Oecophorinae Stathmopolitinae Oecophorini Teichobiinae Oecophorina 26. ERIOCOTTIDAE - Old World Spiny-Winged Moths Herrichiina Eriocottinae Denisiina Compsocteninae Deoclonini 27. PSYCHIDAE - Bagworm Moths Philobotini Lypusinae Syringopaini Stathmopodini Naryciinae Lamprysticini Naryciini Dahlicini Pleurotini Hypertrophinae Taleporiinae Chimabachinae Eotaleporiini Carcinini Taleporiini Chimabachini Placodomini Deuterogoniinae Typhoniinae Penestoglossini Deuterogoniini Aeolanthini Dissoctenioidini 35. LECITHOCERIDAE - Tropical Longhorned Moths Typhoniini Ceuthomadarinae Scoriodytini Oditinae Psychinae Psychini Scythropiodini Oditini Peloponnesini Epichnopterygini Lecithocerinae Stichobasini Torodorinae Oiketicinae 36. ELACHISTIDAE - Grass Miner Moths Metasini Coelopoetinae Acanthopsychini Elachistinae Oreopsychini Mendesiini Phalacropterygini Perittiini Apteronini Stephensiini Oiketicini Elachistini 28. ARRHENOPHANIDAE - Tropical Lattice Moths 37. PTEROLONCHIDAE - Lance-Wing Moths 38. GELECHIIDAE - Twirler Moths Series Gracillariiformes Physoptilinae 29. AMPHITHERIDAE - Double-Eye Moths Gelechiinae 30. SCHRECKENSTEINIIDAE - Bristle-Legged Moths Apatetrini 31. DOUGLASIIDAE - Douglas Moths Anomologini 32. BUCCULATRICIDAE - Ribbed-Cocoon Maker Gelechiini Moths Gnorimoschemini 33. GRACILLARIIDAE - Leafminer Moths Pexicopiinae Gracillariinae Pexicopiini Lithocolletinae Anacampsini Phyllocnistinae Chelariini Dichomeridinae **GELECHIOIDEA** 39. BLASTOBASIDAE - Scavenger Moths 34. OECOPHORIDAE - Concealer Moths Holcocerinae Blastobasinae Depressariinae Blastobasini Depressariini Depressariina Pigritiini Epigraphiina 40. COLEOPHORIDAE - Casebearer Moths Telechrysidina Coleophorinae Amphisbatini Batrachedrinae Cryptolechiiini Epimarptini Cacochroina Batrachedrini Cryptolechiina 41. MOMPHIDAE - Mompha Moths Hypercalliini 42. AGONOXENIDAE - Palm Moths Ethmiinae Agonoxeninae Peleopodinae Blastodacninae Acriini Blastodacnini Peleopodini Parametriotini Autostichinae 43. COSMOPTERIGIDAE - Cosmet Moths Autostichini Antequerinae Oegoconiini Cosmopteriginae Symmocini

Chrysopeleiinae

44. SCYTHRIDIDAE - Flower Moths

COPROMORPHOIDEA

- 45. COPROMORPHIDAE Tropical Fruitworm Moths
- 46. ALUCITIDAE Many-Plumed Moths
- 47. CARPOSINIDAE Fruitworm Moths
- 48. EPERMENIIDAE Fringe-Tufted Moths

Epermeniinae Ochromolopinae

YPONOMEUTOIDEA

- 49. OCHSENHEIMERIIDAE Cereal Stem Moths
- 50. GLYPHIPTERIGIDAE Sedge Moths

Orthoteliinae

Glyphipteriginae

51. PLUTELLIDAE - Diamondback Moths

Ypsolophinae Plutellinae Scythropiinae

Praydinae

- 52. ATTEVIDAE Tropical Ermine Moths
- 53. YPONOMEUTIDAE Ermine Moths

Saridoscelinae

Yponomeutinae

Niphonymphini

Yponomeutini

- 54. ARGYRESTHIIDAE Shiny Head-Standing Moths
- 55. LYONETIIDAE Lyonet Moths

Cemiostominae

Lyonetiinae

Bedelliinae

- 56. ACROLEPIIDAE False Diamondback Moths
- 57. HELIODINIDAE Sun Moths

IMMOIDEA

58. IMMIDAE - Imma Moths

Prodidactinae

Imminae

PYRALOIDEA

59. HYBLAEIDAE - Teak Moths

60. THYRIDIDAE - Picture-Winged Leaf Moths

Simaethistinae

Whalleyaninae

Argyrotypinae

Thyridinae

Siculodinae

Siculodini

Rhodoneurini

Striglininae

61. PYRALIDAE - Snout Moths

Group Crambinina

Crambinae

Ancylolomiini

Diptychophorini

Argyriini

Haimbachiini

Chilonini

Myelobiini

Erupini

Crambini

Schoenobiinae

Cybalomiinae

Linostinae

Scopariinae

Scopariini

Eudoniini

Heliothelini

Musotiminae

Midilinae

Nymphulinae

Nymphulini

Argyractini

Noordinae

Odontiinae

Hercynini Odontiini

Eurrhypini

Wurthiinae

Evergestinae

Evergestini

Orenaiini

Glaphyriinae

Glaphyriini

Hydririni

Cathariini

Dichogamini

Pyraustinae

Pyraustini

Spilomelini

Group Pyralinina

Pyralinae

Pyralini

Endotrichini

Chrysauginae

Galleriinae

Galleriini

Megarthridini

Tirathabini Cacotherapiini

Epipaschiinae

Phycitinae

Cryptoblabini

Phycitini

Acrobasiina

Phycitina

Anerastiini

Peoriini

PTEROPHOROIDEA

- 62. TINEODIDAE False Plume Moths
- 63. OXYCHIROTIDAE Tropical Plume Moths
- 64. PTEROPHORIDAE Plume Moths

Macropiratinae

Agdistinae

Ochyroticinae

Deuterocopinae

Pterophorinae

Platyptiliini

Pterophorini

Subsection SESIINA

SESIOIDEA

65. BRACHODIDAE - Little Bear Moths

Pseudocossinae

Brachodinae

Phycodinae

66. SESIIDAE - Clearwing Moths

Tinthiinae

Tinthiini

Similipepsini

Pennisetiini

Paraglosseciini

Paranthreninae

Paranthrenini

Cissuvorini

Sesiinae

Osminiini

Melittiini

Sesiini

Synanthedonini

Bembeciini

67. URODIDAE - False Burnet Moths

Galacticinae

Urodinae

68. CHOREUTIDAE - Metalmark Moths

Millieriinae

Brenthiinae

Choreutinae

ZYGAENOIDEA

69. HETEROGYNIDAE - Mediterranean Burnet Moths

70 . ZYGAENIDAE - Burnet Moths

Zygaeninae

Phaudinae

Charideinae

Chalcosiinae

Agalopini

Chalcosiini

Anomoeotinae

Callizygaeninae

Procridinae

Procridini

Artonini

71. HIMANTOPTERIDAE - Long-Tailed Burnet Moths

72. LACTURIDAE - Tropical Burnet Moths

73. SOMABRACHYIDAE - Mediterranean Flannel Moths

74. MEGALOPYGIDAE - Flannel Moths

Aidinae

Megalopyginae

Trosiinae

Section COSSINA

Subsection COSSINA

COSSOIDEA

Series Cossiformes

75. COSSIDAE - Carpenterworm Moths

Chilecomadiinae

Hypoptinae

Cossulinae

Cossinae

Zeuzerinae

76. DUDGEONEIDAE - Dudgeon Carpenterworm Moths

77. METARBELIDAE - Tropical Carpenterworm Moths

Series Limacodiformes

78. CYCLOTORNIDAE - Australian Parasite Moths

79. EPIPYROPIDAE - Planthopper Parasite Moths

Epipyropinae

Heteropsychinae

80. DALCERIDAE - Tropical Slug Caterpillar Moths

Acraginae

Dalcerinae

81. LIMACODIDAE - Slug Caterpillar Moths

82. CHRYSOPOLOMIDAE - African Slug Caterpillar

Ectropinae

Moths

Chrysopolominae

CASTNIOIDEA

83. CASTNIIDAE - Giant Butterfly Moths

Synemoninae

Neocastniinae

Castniinae

Castniini

Gazerini

TORTRICOIDEA

84. TORTRICIDAE - Leafroller Moths

Tortricinae

Phricanthini

Schoenotenini

Cnephasiini

Cochylini

Euliini

Tortricini

Ceracini

Archipini

Epitimyrini

Atteriini

Niasomini

Sparganothini

Chlidanotinae

Polyorthini

Hilarographini

Chlidanotini

Olethreutinae

Gatesclarkeanini

Bactrini

Microcorsini

Olethreutini

Eucosmini

Grapholitini

Subsection BOMBYCINA

CALLIDULOIDEA

85. RATARDIDAE - Oriental Parnassian Moths

86. PTEROTHYSANIDAE - Parnassian Moths

Pterothysaninae

Hibrildinae

87. CALLIDULIDAE - Old World Butterfly Moths

Griveaudiinae

Callidulinae

URANIOIDEA

88. EPICOPEIIDAE - Oriental Swallowtail Moths

Schistomitrinae

Epicopeiinae

89. APOPROGONIDAE - African Skipper Moths

90. SEMATURIDAE - American Swallowtail Moths

91. URANIIDAE - Swallowtail Moths

Uraniinae

Microniinae

92. EPIPLEMIDAE - Crenulate Moths

Auzeinae

Epipleminae

GEOMETROIDEA

93. GEOMETRIDAE - Geometer Moths

Archiearinae

Oenochrominae

Alsophilini

Oenochromini

Diptychini Ametridini

Orthostixinae

Ennominae

Abraxini

Cassymini

Eutoeini

Semiothisini	Rhodometrini
Boarmiini	Larentiinae
Apochimina	Trichopterygini
Wilemaniina	Lythriini
Phaseliina	Cataclysmini
Gnophina	Xanthorhoini
Angeronina	Larentiini
Bupalina	Hydriomenini
Cystidiina	Stamnodini
Desertobiina	Cidariini
Cheimoptenina	Operophterini
Boarmiina	Asthenini
	Phileremini
Melanolophiini	Rheumapterini
Bistonini	Solitaneini
Bistonina	Melanthiini
Erranina	Perizomini
Milioniini	Eudulini
Obeidiini	Eupitheciini
	Chesiadini
Diptychini	
Anagogini	94. HEDYLIDAE - American Butterfly Moths
Scardamiini	
Baptini	DADII IONOIDEA
	PAPILIONOIDEA
Plutodini	Series Hesperiiformes
Sphacelodini	
Emplociini	95. HESPERIIDAE - Skipper Butterflies
Lithinini	Coeliadinae
	Pyrrhopyginae
Palyadini	Pyrrhopygini
Caberini	Passovini
Theriini	
Epionini	Oxynetrini
Colotoini	Zoniini
	Pyrginae
Onychorini	
Prosopolophini	Eudamini
Azelinini	Pyrgini
	Trapezitinae
Odontoperini	Heteropterinae
Thinopterygini	
Nacophorini	Megathyminae
Gonodontini	Aegialini
	Agathymini
Oxydiini	Megathymini
Campaeini	
Ennomini	Hesperiinae
Epirranthini	Series Papilioniformes
	96. PAPILIONIDAE - Swallowtail Butterflies
Ourapterygini	트 - 프라티 프로마인 (MEDINE) 프라마인 (MEDINE) (MEDINE) - 스트
Desmobathrinae	Baroniinae
Eumeleini	Parnassiinae
Desmobathrini	Parnassiini
	Zerynthiini
Geometrinae	
Dysphaniini	Papilioninae
Geometrini	Troidini
Pseudoterpnina	Battina
Geometrina	Troidina
	Leptocercini
Timandromorphina	
Neohipparchina	Graphiina
Aracimina	Leptocercina
	Teinopalpini
Nemoriina	Papilionini
Rhomboristina	
Lophochoristina	97. PIERIDAE - Yellow-White Butterflies
Comibaenina	Pseudopontiinae
	Dismorphiinae
Synchlorina	
Hemitheina	Leptideini
Microloxiiina	Dismorphiini
Hierochthoniina	Pierinae
	Pierini
Dichordophorina	
Sterrhinae	Anthocharidini
Sterrhini	Coliadinae
	Colotini
Scopulini	
Cyllopodini	Coliadini
Rhodostrophiini	Gonepterygini
Cosymbiini	
Timandrini	

Pronophilina

96. LICAENIDAE - Gossamer-Winged Butterflies	101. NYMPHALIDAE Brush-Footed Butterflies
Lipteninae Pentilini	Group Nymphalinina
Liptenini	Tellervinae
Poritiinae	Danainae
Liphyrinae	Danaini
Miletinae	Amaurina
Spalgini	Danaina
Lachnocnemini	Euploeini
Miletini	Euploeina
Tarakina	Ideina
Miletina	Lycoreini
Curetinae	Itunina
Lycaeninae	Lycoreina
Theclinae	Ithomiinae
Eumaeini	Tithoreini
Eumaeina	Melinaeini
Tomarina	Mechanitini
Deudorigina	Napeogenini Ithomiini
Luciini	Oleriini
Theclini	Dircennini
Ogyrina	Godyridini
Arhopalina	Acraeinae
Theclina	Acraeini
Zesiini	Pardopsini
Amblypodiini	Heliconiinae
Aphnaeini	Argynnini
Iolaini	Heliconiini
Iolaina	Nymphalinae
Remelanina	Coeini
Hypolycaenina	Nymphalini
Horagini	Kallimini
Oxylidina	Melitaeini
Hypotheclina	Euphydryina
Catapaecilmatina	Melitaeina
Loxurina	Phyciodina
Horagina	Neptini
Cheritrina	Pseudergolini
Polyommatinae	Euthaliini
Lycaenesthini	Catagrammini
Candalidini	Limenitidini
Polyommatini	Cyrestidini
Zizeerina	Biblidini
Lampidina	Biblidina
Everina	Eurytelina
Lycaenopsina	Epicaliina
Scolitantidina	Ageroniina
Polyommatina Niphandina	Epiphilina
99. RIODINIDAE - Metalmark Butterflies	Eubagina
Styginae Styginae	Callicorina
Hamearinae	Group Satyrinina
Euselasiinae	Calinaginae
Corrachiinae	Apaturinae
Riodininae	Apaturini
Mesosemiini	Charaxini
Mesosemiina	Anaeini
Napaeina	Zaretini
Eurybiini	Preponini
Riodinini	Prothoini Amathusiinae
Symmachiini	The state of the s
Charitini	Morphinae
Helicopini	Antirrheini
Emesini	Morphini
Nymphidiini	Brassolinae
Aricorina	Biini Proceedini
Lemoniadina	Brassolini
Nymphidiina	Naropini
Theopina	Satyrinae
Stalachtini	Lethini Saturini
100. LIBYTHEIDAE - Snout Butterflies	Satyrini Pronophilina
Dilout Dutterlifes	FIOHOPHINA

Maniolina Erebiina Hypocystina Euptychiina Coenonymphina Melanargiina Satyrina Hipparchiina Melanitini

Haeterini Elymniini Parargina Elymniina

Mycalesini Ypthimini

DREPANOIDEA

102. AXIIDAE - Gold Moths

103. THYATIRIDAE - False Owlet Moths

Thyatirinae

Thyatirini Habrosynini Macrothyatirini Tetheini Polyplocinae

Ceranemotini Polyplocini

104. CYCLIDIIDAE - Giant Hooktip Moths

105. DREPANIDAE - Hooktip Moths

Drepaninae Drepanini Cilicini

Oretinae Nidarinae

BOMBYCOIDEA

Series Bombyciformes

106. CARTHAEIDAE - Australian Silkworm Moths

107. EUPTEROTIDAE - Giant Lappet Moths Janinae

Eupterotinae Panacelinae

108. APATELODIDAE - American Silkworm Moths

Apatelodinae Epiinae

Phiditiinae

109. BOMBYCIDAE - Silkworm Moths

Bombycinae Prismostictinae

110. MIMALLONIDAE - Sackbearer Moths

111. ANTHELIDAE - Australian Lappet Moths

Munychryiinae Anthelinae

112. LASIOCAMPIDAE - Lappet Moths

Chondrosteginae Chionopsychinae Poecilocampinae Macromphaliinae Lasiocampinae

Gonometini Gastropachini Lasiocampini Malacosomatini

Series Saturniiformes

113. ENDROMIDAE - Glory Moths

Endrominae Mirininae

114. LEMONIIDAE - Autumn Silkworm Moths Lemoniinae

Sabaliinae

115. BRAHMAEIDAE - Brahmin Moths

Dactyloceratinae

Brahmaeinae

116. OXYTENIDAE - Tropical American Silkworm

Moths

117. CERCOPHANIDAE - Andean Moon Moths

Cercophaninae Janiodinae

118. SATURNIIDAE - Emperor Moths

Arsenurinae

Arsenurini Almeidaiini

Ceratocampinae

Hemileucinae

Hemileucini Polythysanini

Agliinae Ludiinae Salassinae

Saturniinae

Saturniini Attacini Urotini

SPHINGOIDEA

119. SPHINGIDAE - Hawk Moths

Sphinginae

Acherontiini Sphingini

Smerinthinae

Sphingulini Ambulycini

Smerinthini

Macroglossinae

Dilophonotini Philampelini Macroglossini

NOCTUOIDEA

120. NOTODONTIDAE - Prominent Moths

Group Oenosandrinina Oenosandrinae

Group Thaumetopoeinina Thaumetopoeinae

Group Notodontinina

Pygaerinae

Pygaerini Rosemini

Platychasmatinae

Notodontinae

Notodontini Dicranurini

Phalerinae Dudusinae

Dudusini Scranciini

Heterocampinae

Heterocampini Stauropini Hemiceratini

Hapigiini Hylaeorini Desmeocraerini

Ceirini Acrotenini

Nystaleinae Meiceratinae

121. DIOPTIDAE - American False Tiger Moths Doinae

Chloephorinae Westermanniini

Dioptinae	Eariadini
122. THYRETIDAE - African Maiden Moths	Chloephorini
123. LYMANTRIIDAE - Tussock Moths	Careini
Orgyiinae Orgyiinae	Camptolomini
Lymantriinae	Ariolicini
Lymantriini	Nolinae
Euproctidini	Plusiinae
124. ARCTIDAE - Tiger Moths	Abrostolini
	Argyrogrammatini
Group Pericopinina	Plusiini
Pericopinae	Autoplusiina
Group Arctiinina Lithosiinae	Euchalciina
	Plusiina
Phryganopterygini	Acontiinae
Acsalini	Cydosiini
Lithosiini	Eustrotiini
Eudesmiini	Eublemmini
Cisthenini	Acontiini
Nudariini	Bagisarini
Endrosini	Condicinae
Arctiinae	Condicini
Callimorphini	Leuconyctini
Callimorphina	Amphipyrinae
Nyctemerina	Amphipyrini
Utetheisini	Phosphilini
Arctiini	Stiriinae
Phaegopterini	Psaphidinae
Group Ctenuchinina	Psaphidini
Ctenuchinae	Nocloini
Syntominae	Triocnemidini
Syntomini	Feraliini
Euchromiini	Azeniini
125. NOCTUIDAE - Owlet Moths	Grotellini
Group Aganainina	
Aganainae	Agaristinae
Group Herminiinina	Agaristini
Herminiinae	Eucocytiini
Group Noctuinina	Cocytiinae
Hypenodinae	Heliothinae
Strepsimanini	Acronictinae Acronictini
Hypenodini	
Rivulinae	Raphiini Pantheinae
Hypeninae	Pantheini
Catocalinae	
Ophiderini	Dilobini
Pangraptina	Bryophilinae
Aventiina	Cuculliinae Cuculliini
Ophiderina	
Toxocampina	Xylenini Xylenina
Tytina	Antitypina
Polydesmina	Oncocnemidini
Aediina	Hadeninae
Catephiina	
Drasteriina	Apameini
Euclidiina	Eriopini
Achaeina	Elaphriini
Ophiusina	Hadenini
Thysaniina	Eriopygini
Gonopterini	Glottulini
Anomiini	Noctuinae
Catocalini	Agrotini
Euteliinae	Agrotina
Stictopterinae	Austrandesiina
Sarrothripinae	Aniclini
Eligmini	Noctuini
Risobini	Ufeini
Blenini	
Afridini	
Sarrothripini	
Collomenini	

Ariolicini (Noctuidae), 125

INDEX TO FAMILY-LEVEL NAMES

The index includes 1,984 names (including synonyms). Highlighted names are higher category names (above family level) and are capitalized (except for series names). Valid family names are also capitalized, but not highlighted. Synonyms have been added to the index as currently known; these are not noted in the classification listing but can be found in the index with notation to the current correct name (synonyms are all lower case in italics). All names are referred by family number; higher category names are referred to the next listed family number. A large number of subtribal names noted for European Noctuidae by Beck (1999b) are not included herein since the rest of the world is not defined equivalently to subtribal level in this family.

As in all zoological naming, the names are spelled according to specified rules as to endings, although there is irregularity in some of the higher category names: order names end in -tera; suborder, cohort, subcohort, infraorder names vary and end in -tera, -ata, or -ina; division names end in -trysia; section and subsection names end in -ina; superfamily names end in -oidea; series names end in -formes; family names end in -idae; group names end in -inia; subfamily names end in -inae; tribal names end in -ini; and subtribe names end in -ina (sometimes written with -idi or -iti endings). The actual taxon names are modified in some cases due to the spelling of the name-bearing generic-base name that a tribe, subfamily, or family is derived from: the spellings are corrected herein as currently known about (e.g., in Riodinidae, the misformed tribal name Lemoniini is corrected to Lemoniadini (as already noted in Hall and Heppner, 1999. Trop. Lepid., 10:30) (now Lemoniadina), since the generic-base name, Lemonias, is different from the name Lemonia, in the moth family Lemoniidae).

Abraxinae (= Abraxini), 93 Abraxini (Geometridae), 93 Abrostolini (Noctuidae), 125 Acanthopsychinae (= Oiketicinae), 27 Acanthopsychini (Psychidae), 27 ACANTHOPTEROCTETIDAE, 5 Acanthopteroctetinae (Acanthopteroctetidae), Acentriinae (= Nymphulinae), 61 Acentropidae (= Nymphulinae), 61 Acentropinae (= Nymphulinae), 61 Acentropodinae (= Nymphulinae), 61 Achaeini (= Achaeina), 125 Achaeina (Noctuidae), 125 Acherontiinae (= Sphinginae), 119 Achinagidae [sic] (= Oecophoridae), 34 Acidaliidae (= Sterrhinae), 93 Acidaliinae (= Sterrhinae), 93 Acidaliini unavail. (= Scopulini), 93 Acontiidae (= Acontiinae), 125 Acontiinae (Noctuidae), 125 Acontiini (Noctuidae), 125 Acraeidae (= Acraeinae), 101 Acraeinae (Nymphalidae), 101 Acraeini (Nymphalidae), 101 Acraginae (Dalceridae), 80 Acriinae (= Acriini), 34 Acriini (Oecophoridae), 34 Acrobasiina (Pyralidae), 61 ACROLEPIIDAE, 56 ACROLOPHIDAE, 24 Acrolophinae (Acrolophidae), 24 Acronictinae (Noctuidae), 125 Acronictini (Noctuidae), 125 Acronyctidae [sic] (= Acronictinae), 125 Acronyctinae [sic] (= Acronictinae), 125 Acrotenini (Notodontidae), 120 Acsalini (Arctiidae), 124 ADELIDAE, 22 Adelinae (Adelidae), 22 Adelocephalidae (= Ceratocampinae), 118 Adelocephalinae (= Ceratocampinae), 118 Adeloidea (= Incurvarioidea), 19 Adscitinae (= Procridinae), 70 Aechmiidae (= Glyphipterigidae), 50 Aecophoridae [sic] (= Oecophoridae), 34 Aediinae (=Aediina), 125 Aediini (= Aediina), 125 Aediina (Noctuidae), 125 Aegeriidae (= Sesiidae), 66 Aegeriinae (= Sesiinae), 66 Aegeriini (= Sesiini), 66 Aegerioidea (= Sesioidea), 65 Aegeroniina (Nymphalidae), 101 Aegeroniini (= Limenitidini), 101 Aegialini (Hesperiidae), 95 Aeolanthidae (= Aeolanthini), 34 Aeolanthinae (= Aeolanthini), 34 Aeolanthini (Oecophoridae), 34 Afridini (Noctuidae), 125 Agalopini (Zygaenidae), 70 Aganaidae (= Aganainae), 125 Aganainae (Noctuidae), 125 Aganainina (Noctuidae), 125 Agapetidae (= Cochylini), 84

Agapetinae (= Satyrini), 101

Agaristidae (= Agaristinae), 125 Agaristinae (Noctuidae), 125 Agaristini (Noctuidae), 125 AGATHIPHAGIDAE, 2 AGATHIPHAGOIDEA, 2 Agathymini (Hesperiidae), 95 Agdistinae (Pterophoridae), 64 Agdistopidae [sic] (= Macropiratinae), 46 Ageroniini (= Ageroniina), 101 Ageroniina (Nymphalidae), 101 Agliinae (Saturniidae), 118 AGLOSSATA, 2 Aglossinae (= Pyralini), 61 AGONOXENIDAE, 42 Agonoxeninae (Agonoxenidae), 42 Agrophilidae (= Acontiinae), 125 Agroterinae (= Pyraustinae), 61 Agrotidae (= Noctuinae), 125 Agrotini (Noctuidae), 125 Agrotina (Noctuidae), 125 Aidinae (Megalopygidae), 74 Aletinae (= Scopulini), 93 Aletini (= Scopulini), 93 Almeidaiini (Saturniidae), 118 Alsophilinae (= Alsophilini), 93 Alsophilini (Geometridae), 93 ALUCITIDAE, 46 Amathusiidae (= Amathusiinae), 101 Amathusiinae (Nymphalidae), 101 Amatidae (= Ctenuchinae), 124 Amaurina (Nymphalidae), 101 Amblypodiini (Lycaenidae), 98 Amblyptiliinae (= Pterophorinae), 64 Ambulicinae (= Smerinthini), 119 Ametridini (Geometridae), 93 Amphidasidae (= Bistonini), 93 Amphidasini (= Bistonina), 93 Amphipyridae (= Amphipyrinae), 125 Amphipyrinae (Noctuidae), 125 Amphipyrini (Noctuidae), 125 Amphisbatidae (= Amphisbatini), 34 Amphisbatini (Oecophoridae), 34 AMPHITHERIDAE, 29 Amydriinae (Acrolophidae), 24 Anacampsidae (= Anacampsini), 38 Anacampsinae (= Anacampsini), 38 Anacampsini (Gelechiidae), 38 Anaeini (Nymphalidae), 101 Anagogini (Geometridae), 93 Anaphinae (Thaumetopoeinae), 120 Anaphoridae (= Acrolophidae), 24 Anarsiidae (Gelechiidae), 38 Anarsiinae (= Chelariini), 38 Anchiniinae (= Hypercalliini), 34 Anchyloperidae (= Tortricinae), 84 Anchyloperinae (= Olethreutini), 84 Ancylolomiini (Pyralidae), 61 ANDESIANIDAE, 14 ANDESIANOIDEA, 14 Anerastiinae (= Anerastiini), 61 Anerastiini (Pyralidae), 61 Angeronini (= Angeronina), 93 Angeronina (Geometridae), 93 Aniclini (Noctuidae), 125 Animulidae (= Psychidae), 27

Anomiini (Noctuidae), 125

Anomoeotinae (Zygaenidae), 70 Anomologidae (= Anomologini), 38 Anomologinae (= Anomologini), 38 Anomologini (Gelechiidae), 38 ANOMOSETIDAE, 10 Antequerinae (Cosmopterigidae), 43 ANTHELIDAE, 111 Anthelinae (Anthelidae), 111 Antirrheini (Nymphalidae), 101 Anthocharinae (= Anthocharidini), 97 Anthocharidini (Pieridae), 97 Anthroceridae (= Zygaenidae), 70 Anthrocerinae (= Zygaeninae), 70 Anthroceroidea (= Zygaenoidea), 69 Antitypina (Noctuidae), 125 Anuini (= Ophiusina), 125 Anycylolomiidae (= Ancylolomiini), 61 Anycylolomiinae (= Ancylolomiini), 61 Apameidae (= Apameini), 125 Apameini (Noctuidae), 125 Apatelinae (= Acronictinae), 125 APATELODIDAE, 108 Apatelodinae (Apatelodidae), 108 Apatetrinae (= Apatetrini), 38 Apatetrini (Gelechiidae), 38 Apaturidae (= Apaturinae), 101 Apaturinae (Nymphalidae), 101 Apaturini (Nymphalidae), 101 Apeirini (= Ennomini), 93 Apheliidae unavail. (= Bactrini), 84 Apheliinae unavail. (= Bactrini), 84 Aphelosetiidae (= Elachistidae), 36 Aphnaeini (Lycaenidae), 98 Aplotinae (= Coleophorinae), 40 Apochimini (= Apochimina), 93 Apochimina (Geometridae), 93 Apodidae (= Limacodidae), 81 Apopestini (= Toxocampina), 125 APOPROGONIDAE, 89 Aporiinae (= Pierinae), 97 Apteroninae (= Oiketicinae), 27 Apteronini (Psychidae), 27 Aracimini (= Aracimina), 93 Aracimina (Geometridae), 93 Arbelidae (= Metarbelidae), 77 Archaeobalbina (= Pseudoterpnina), 93 Archaeopachinae (= Metarbelidae), 77 Archieariinae [sic] (= Archiearinae), 93 Archiearinae (Geometridae), 93 Archimeessiinae (= Dryadaulinae), 25 Archipini (Tortricidae), 84 ARCTIIDAE, 124 Arctiinae (Arctiidae), 124 Arctiini (Arctiidae), 124 Arctiinina (Arctiidae), 124 Argynnidae (= Argynnini), 101 Argynninae (= Argynnini), 101 Argynnini (Nymphalidae), 101 Argyractini (Pyralidae), 61 ARGYRESTHIIDAE, 54 Argyriini (Pyralidae), 61 Argyrogrammatini (Noctuidae), 125 Argyroplocini (= Olethreutini), 84 Argyrotypini (Thyrididae), 60

Arhopalina (Lycaenidae), 98

Aricorina (Riodinidae), 99

Aristoteliinae (= Anomologini), 38 ARRHENOPHANIDAE, 28 Arsenurinae (Saturniidae), 118 Arsenurini (Saturniidae), 118 Artonini (Zygaenidae), 70 Ascalaphini (= Thysaniina), 125 Ascotinae (= Boarmiini), 93 Ascotinae (= Boarmiina), 93 Ascotini (= Boarmiina), 93 Asemanophorae [sic] (= Sphinginae), 119 Asemanophorinae (= Sphinginae), 119 Ashinagidae (= Oecophoridae), 34 Asopiidae (= Pyralinae), 61 Asopiinae (= Pyralini), 61 Aspilatidae (= Angeronini), 93 Aspilatini [sic] (= Angeronina), 93 Aspitatini (= Angeronina), 93 Asteroscopina (= Psaphini), 125 Astheninae (= Asthenini), 93 Asthenini (Geometridae), 93 Attacidae (= Saturniidae), 118 Attacinae (= Saturniinae), 118 Attacini (Saturniidae), 118 Attacoidea (= Saturniiformes), 113 Atteridae [sic] (Atteriini), 84 Atteriini (Tortricidae), 84 ATTEVIDAE, 52 Atychiidae (= Brachodidae), 65 Augasmidae (= Coleophoridae), 40 Austrandesiina (Noctuidae), 125 Autographini (= Plusiini), 125 Automerinae (= Hemileucinae), 118 Autoplusiina (= Autoplusiina). 125 Autoplusiina (Noctuidae), 125 Autostichidae (= Autostichinae), 34 Autostichinae (Oecophoridae), 34 Autostichini (Oecophoridae), 34 Auzeinae (Epiplemidae), 92 Aventiidae (= Ophiderini), 125 Aventiini (Aventiina), 125 Aventiina (Noctuidae), 125 AXIIDAE, 102 Axioidea (= Axiidae), 102 Azelinini (Geometridae), 93 Azeniini (Noctuidae), 125 Azinidae (= Ethmiinae), 34 Bactrini (Tortricidae), 84 Bagisarinae (= Bagisarini), 125 Bagisarini (Noctuidae), 125 Baptini (Geometridae), 93 Baroniinae (Papilioniidae), 96 Batrachedridae (= Batrachedrinae). 40 Batrachedrinae (Coleophoridae), 40 Batrachedrini (Coleophoridae), 40 Battina (Papilioniidae), 96 Bedelliidae (= Lyonetiidae), 55 Bedelliinae (Lyonetiidae), 55 Bembeciini (= Synanthedonini), 66 Beninae (= Chloephorini), 125 Biblidina (Nymphalidae), 101 Biblidini (Nymphalidae), 101 Biblini [sic] (= Biblidini), 101 Biini (Nymphalidae), 101 Bijuginae (= Psychinae), 27 Biselachistini (= Elachistini), 36

18

Bistoninae (= Bistonini), 93 Bistonini (Geometridae), 93 Blabophaninae (= Tineidae), 25 Blabophaninae (= Tineinae), 25 BLASTOBASIDAE, 39 Blastobasinae (Blastobasidae), 39 Blastobasini (Blastobasidae), 39 Blastodacnidae (= Agonoxenidae), 42 Blastodacninae (Agonoxenidae), 42 Blastodacnini (Agonoxenidae), 42 Blenini (Noctuidae), 125 Boarmiidae (= Boarmiini), 93 Boarmiinae (= Boarmiini), 93 Boarmiini (Geometridae), 93 Boarmiina (Geometridae), 93 Boletobiidae (= Ophiderini), 125 Bolinidae (= Ophiderini), 125 Boloriini (= Argynnini), 101 BOMBYCIDAE, 109 Bombyciformes, 106 BOMBYCINA, 85 Bombycinae (Bombycidae), 109 BOMBYCOIDEA, 106 Botydae (= Pyraustinae), 61 Botynae (= Pyraustinae), 61 Braccini (= Boarmiina), 93 Brachmiinae (= Dichomeridinae), 38 BRACHODIDAE, 65 Brachodinae (Brachodidae), 65 Bradypodicolinae (= Chrysauginae), 61 BRAHMAEIDAE, 115 Brahmaeinae (Brahmaeidae), 115 Brassolidae (= Brassolinae), 101 Brassolinae (Nymphalidae), 101 Brassolini (Nymphalidae), 101 Brenthiinae (Choreutidae), 68 Bryoleucini (= Bryophilinae), 125 Bryophilidae (= Bryophilinae), 125 Bryophilinae (Noctuidae), 125 **BUCCULATRICIDAE**, 32 Bucculatrigidae [sic] (= Bucculatricidae), 32 Bupalini (= Bupalina), 93 Caberidae (= Caberini), 93 Caberini (Geometridae), 93 Cacochroini (= Cacochroina), 34 Cacochroina (Oecophoridae), 34 Cacoeciinae (= Tortricinae), 84 Cacoeciini (= Tortricini), 84 Cacotherapiini (Pyralidae), 61 Calamiina (= Apameini), 125 Calinaginae (Nymphalidae), 101 Calinaginae (Nymphalidae), 101 Callicorina (Nymphalidae), 101 Callidryinae (= Gonepterygini), 97 Callidryini (= Gonepterygini), 97 CALLIDULIDAE, 87 Callidulinae (Callidulidae), 87 CALLIDULOIDEA, 85 Callimorphidae (= Callimorphini), 124 Callimorphinae (= Callimorphini), 124 Callimorphini (Arctiidae), 124 Callimorphina (Arctiidae, 124) Callizygaeninae (Zygaenidae), 70 Callophryini (= Theclini), 98 Caloptiliidae (= Gracillariidae), 33 Caloptiliinae (= Gracillariinae), 33 Calothysanini (= Timandrini), 93 Calpidae (= Ophiderini), 125 Calpini (= Ophiderini), 125 Campaeini (Geometridae), 93 Camptolomini (Noctuidae), 125 Canephoridae (= Psychidae), 27 Canephorinae (= Oiketicinae), 27 Caradrinidae (= Hadeninae), 125 Carcinidae (= Carcinini), 34 Carcininae (= Carcinini), 34 Carcinini (Oecophoridae), 34 Careini (Noctuidae), 125 Carpocapsidae (= Grapholitini), 84 Carpocapsinae (= Grapholitini), 84 CARPOSINIDAE, 47 Carterocephalinae (= Heteropterinae), 95 CARTHAEIDAE, 106 Cassymini (Geometridae), 93 CASTNIIDAE, 83 Castniinae (Castniidae), 83 Castniini (Castniidae), 83 CASTNIOIDEA, 83 Cataclysmini (Geometridae), 93

Catagrammini (Nymphalidae), 101 Catapaecilmatina (Lycaenidae), 98 Catapterigidae (= Catapteriginae), 5 Catapteriginae (Acanthopteroctetidae), 5 Catephiidae (= Catephiina), 125 Catephiini (= Catephiina), 125 Catephiina (Noctuidae), 125 Cathariinae (= Odontiinae), 61 Cathariini (Pyralidae), 61 Catocalidae (= Catocalinae), 125 Catocalinae (Noctuidae), 125 Catocalini (Noctuidae), 125 Catopyrrhinae (= Caberini), 93 Catopyrrhinae (= Caberini), 93 Catopyrrhini (= Caberini), 93 Cecidosetidae [sic] (= Cecidosidae), 20 CECIDOSIDAE, 20 Cedestinae (Yponomeutidae), 53 Ceirini (Notodontidae), 120 Celastrinini (= Everina), 98 Cemiostomidae (= Lyonetiidae), 55 Cemiostominae (Lyonetiidae), 55 Ceracidae (= Ceracini), 84 Ceracini (Tortricidae), 84 Ceranemotini (Thyatiridae), 103 Ceratocampidae (= Ceratocampinae), 118 Ceratocampinae (Saturniidae), 118 CERCOPHANIDAE, 117 Cercophaninae (Cercophanidae), 117 Cerostominae (= Ypsolophinae), 51 Cerurinae (= Dicranurini), 120 Ceuthomadarinae (Lecithoceridae), 35 Chaerocampini [sic] (= Macroglossinae), 119 Chalcosiinae (Zygaenidae), 70 Chalcosiini (Zygaenidae), 70 Chaliinae (= Psychinae), 27 Chapmaniidae (= Eriocraniidae), 4 Charaxidae (= Charaxini), 101 Charaxidinae (Charaxini), 101 Charaxinae (Charaxini), 101 Charaxini (Nymphalidae), 101 Charideinae (Thyrididae), 60 Charitini (Riodinidae), 99 Chauliodidae (= Epermeniidae), 48 Cheimatobiini (= Theriini), 93 Cheimoptenini (= Cheimoptenina), 93 Chelariinae (= Chelariini), 38 Chelariini (Gelechiidae), 38 Cheloniidae (= Arctiidae), 124 Cheritrina (Lycaenidae), 98 Chesiadinae (= Chesiadini), 93 Chesiadini (Geometridae), 93 Chilecomadiinae (Cossidae), 75 Chilonidae (= Chilonini), 61 Chiloninae (= Chilonini), 61 Chilonini (Pyralidae), 61 Chimabacchidae [sic] (= Chimabachinae), 34 Chimabachidae (= Chimabachinae), 34 Chimabachinae (Oecophoridae), 34 Chimabachini (Oecophoridae), 34 Chimatophilidae (= Tortricini), 84 Chionopsychinae (Lasiocampidae), 112 Chlidanotinae (= Chlidanotini), 84 Chlidanotinae (Tortricidae), 84 Chlidanotini (Tortricidae), 84 Chloephorinae (Noctuidae), 125 Chloephorini (Noctuidae), 125 Chlorochromidae (= Geometrini), 93 Chlorochrominae (= Geometrinae), 93 Chlorochromini (= Hemitheina), 93 Choerocampinae (= Macroglossinae), 119 Chondrosteginae (Lasiocampidae), 112 CHOREUTIDAE, 68 Choreutinae (Choreutidae), 68 Choreutoidea (= Sesioidea), 65 Chrysaugidae (= Chrysauginae), 61 Chrysauginae (Pyralidae), 61 Chrysoesthiidae (Gelechiidae), 38 Chrysopeleiidae (= Chrysopeleiinae), 43 Chrysopeleiinae (Cosmopterigidae), 43 Chrysopeleioidea (= Gelechioidea), 34 Chrysophaninae (= Lycaeninae), 98 Chrysophanini (= Lycaeninae), 98 CHRYSOPOLOMIDAE, 82 Chrysopolominae (Chrysopolomidae), 82 Cidariidae (= Cidariini), 93 Cidariinae (= Cidariini), 93 Cidariini (Geometridae), 93

Cilicidae (= Cilicini), 105 Cilicini (Drepanidae), 105 Cimeliidae (= Axiidae), 102 Cingiliini (= Ennomini), 93 Cissuvorini (Sesiidae), 66 Cisthenini (Arctiidae), 124 Citheroniidae (= Ceratocampinae), 118 Citheroniinae (= Ceratocampinae), 118 Cledeobiinae (= Pyralinae), 61 Cleophanini unavail. (= Oncocnemidini), 125 Cleorini (= Boarmiina), 93 Cnephasiidae (= Cnephasiini), 84 Cnephasiini (Tortricidae), 84 Cnethocampinae (Thaumetopoeinae), 120 Cochlidiae (= Limacodidae), 81 Cochlidiidae (= Limacodidae), 81 Cochlidionidae (= Limacodidae), 81 Cochliopodi unavail. (= Limacodidae), 81 Cochylidae (= Cochylini), 84 Cochylini (Tortricidae), 84 Cocytiinae (Noctuidae), 125 Coeini (Nymphalidae), 101 Coeliadinae (Hesperiidae), 95 Coelopoetidae (= Elachistinae), 36 Coelopoetinae (Elachistidae), 36 Coenonymphinae (= Satyrini), 101 Coenonymphini (= Coenonymphina), 101 Coenonymphina (Nymphalidae), 101 COLEOPHORIDAE, 40 Coleophorinae, 40 Coleophoroidea (= Gelechioidea), 34 Coliadidae (= Coliadinae), 97 Coliadinae (Pieridae), 97 Coliadini (Pieridae), 97 Collomenini (Noctuidae), 125 Coloburini (= Coeini), 101 Colotini (Pieridae), 97 Colotoini (Geometridae), 93 Comibaenini (= Comibaenina), 93 Comibaenina (Geometridae), 93 Commophilidae (= Cochylini), 84 Comostolini (= Hemitheina), 93 Compsoctenidae (= Compsocteninae), 26 Compsocteninae (Eriocottidae), 26 Compsopterini (= Prosopolophini), 93 Conchylidae [sic] (= Cochylini), 84 Condicinae (Noctuidae), 125 Condicini (Noctuidae), 125 COPROMORPHIDAE, 45 COPROMORPHOIDEA, 45 Corrachiinae (Riodinidae), 99 Cosmiidae (= Hadeninae), 125 Cosmiinae (= Hadeninae), 125 Cosmiotini (= Elachistini), 36\ Cosmodesmidi (= Leptocircini), 96 COSMOPTERIGIDAE, 43 Cosmopteriginae (Cosmopterigidae), 43 Cosmopterigoidea (= Gelechioidea), 34 Cosmopterygidae [sic] (= Cosmopterigidae), COSSIDAE, 75 Cossiformes, 75 COSSINA, 75 Cossinae (Cossidae), 75 COSSOIDEA, 75 Cossulinae (Cossidae), 75 Cosymbiinae (= Cosymbiini), 93 Cosymbiini (Geometridae), 93 Crambinae (Pyralidae), 61 Crambini (Pyralidae), 61 Crambinina (Pyralidae), 61 Crinopterygidae (= Crynopteryginae), 19 Crinopteryginae (Incurvariidae), 19 Crocallini (= Ennomini), 93 Cryptoblabini (Pyralidae), 61 Cryptolechiidae (= Cryptolechiini), 34 Cryptolechiiini (Oecophoridae), 34 Cryptolechiini (= Cryptolechiina), 34 Cryptolechiina (Oecophoridae), 34 Cryptophasidae (= Oecophoridae), 34 Cryptopyrrhini [sic] (= Caberini), 93 Ctenuchidae (=Ctenuchinae), 124 Ctenuchinae (Arctiidae), 124 Ctenuchinina (Arctiidae), 124 Cuculliidae (= Cuculliinae), 125 Cuculliinae (Noctuidae), 125 Cuculliini (Noctuidae), 125 Curetinae (Lycaenidae), 98

Cyanirini (= Polyommatina). 98 Cybalomiinae (Pyralidae), 61 CYCLIDIIDAE, 104 Cyclopedinae [sic] (Heteropterinae), 95 Cyclophoridae (= Cosymbini), 93 Cyclophorini (= Cosymbiini), 93 Cyclopidinae nom. oblit. (Heteropterinae), 95 CYCLOTORNIDAE, 78 Cycnodiidae (= Elachistidae). 36 Cycnodioidea (= Gelechioidea), 34 Cydosiini (Noctuidae), 125 Cygnodiidae [sic] (= Elachistidae), 36 Cyllopodidae (= Cyllopodini). 93 Cyllopodini (Geometridae). 93 Cymatophoridae (= Thyatiridae), 103 Cyrestidini (Nymphalidae), 101 Cyrestini [sic] (= Cyrestidini), 101 Cystidiini (= Cystodiina), 93 Cystidiina (Geometridae), 93 DACNONYPHA, 4 Dactyloceratinae (Brahmaeidae), 115 Dahlicini (Psychidae), 27 DALCERIDAE, 80 Dalcerinae (Dalceridae), 80 Daliminae (= Boarmiini), 93 Dalimini (= Boarmiina), 93 Danaidae (= Danainae), 101 Danainae (Nymphalidae), 101 Danaini (Nymphalidae), 101 Danaina (Nymphalidae), 101 Danaoidea (= Danainae), 101 Dasyceridae (= Oecophoridae), 34 Dasycerinae (= Oecophorinae), 34 Dasychiridae (= Lymantriidae), 123 Dasydiini (= Gnophina), 93 Deiliniinae (= Caberini), 93 Deiliniini (= Caberini), 93 Dendroneurinae (= Hieroxestinae), 25 Dendroneurinae (= Tineidae), 25 Denisiina (Oecophoridae), 34 Denisiini (= Denisiina), 34 Deoclonidae (= Deoclonini), 34 Deoclonini (Oecophoridae), 34 Depressariidae (= Depressariinae), 34 Depressariinae (Oecophoridae), 34 Depressariini (Oecophoridae), 34 Depressariina (Oecophoridae), 34 Desertobiini (= Desertobiina), 93 Desertobiina (Geometridae), 93 Desmeocraerini (Notodontidae), 120 Desmobathrinae (Geometridae), 93 Desmobathrini (Geometridae), 93 Deudorigina (Lycaenidae). 98 Deuterocopinae (Pterophoridae), 64 Deuterogoniinae (Oecophoridae), 34 Deuterogoniini (Oecophoridae), 34 Deuterotineidae (= Eriocottinae), 26 Deuterotineinae (= Eriocottinae), 26 Dichocrociinae (= Pyraustinae), 61 Dichogaminae (= Dichogamini), 61 Dichogamini (Pyralidae), 61 Dichomeridae (= Dichomeridinae), 38 Dichomeridinae (Gelechiidae), 38 Dichomerinae [sic] (= Dichomeridinae), 38 Dichomerisinae [sic] (= Dichomeridinae), 38 Dichopinae (= Psaphidinae), 125 Dichordophorini (= Dichordophorina), 93 Dichordophorina (Geometridae), 93 Dicranurini (Notodontidae). 120 Dilobidae (= Dilobini), 125 Dilobinae (= Dilobini), 125 Dilobini (Noctuidae), 125 Dilophonotidae (= Dilophonotini), 119 Dilophonotinae (= Dilophonotini), 119 Dilophonotini (Sphingidae), 119 DIOPTIDAE, 121 Dioptinae (Dioptidae), 121 Diphtherinae (= Pantheinae), 125 Diplodomidae (= Psychidae). 27 Diplodominae (= Taleporimae). 27 Diplosaridae (= Cosmopterigidae), 43 Diptychinae (= Diptychini), 93 Diptychini (Geometridae), 93 Diptychini (Geometridae), 93 Diptychophorini (Pyralidae), 61 Dircennini (Nymphalidae), 101 Dirphiadae [sic] (= Hemileucinae), 118 Dirphiinae (= Hemileucinae), 118

Dismorphiinae (Pieridae), 97 Dismorphiini (Pieridae), 97 Dissoctenidae (= Psychidae), 27 Dissocteninae (= Penestoglossinae), 27 Dissoctenioidini (Psychidae), 27 DITRYSIA, 24 Diuerninae (= Chimabachinae), 34 Diurneidae (= Chimabachinae), 34 Divianini (= Phycitini), 61 Doinae (Dioptidae), 121 Doritinae (= Parnassiinae), 96 DOUGLASIIDAE, 31 Drasteriini (= Drasteriina), 125 Drasteriina (Noctuidae), 125 DREPANIDAE, 105 Drepaninae (Drepanidae), 105 DREPANOIDEA, 102 Dryadaulinae (Tineidae), 25 Dryocampinae (= Ceratocampinae), 118 Dryocampini (= Ceratocampinae), 118 **DUDGEONEIDAE**, 76 Dudusinae (Notodontidae), 120 Dudusini (Notodontidae), 120 Dysphaniinae (= Dysphaniini), 93 Dysphaniini (Geometridae), 93 Eariadini (Noctuidae), 125 Ectomyeloisini (= Phycitini), 61 Ectropinae (Chrysopolomidae), 82 ELACHISTIDAE, 36 Elachistinae (Elachistidae), 36 Elachistini (Elachistidae), 36 Elachistini (Elachistidae), 36 Elachistoidea (= Gelechioidea), 34 Elaphriinae (= Elaphriini), 125 Elaphriini (Noctuidae), 125 Eligmini (Noctuidae), 125 Elymniinae (= Elymniini), 101 Elymniini (Nymphalidae), 101 Elymniini (= Elymniina), 101 Elymniina (Nymphalidae), 101 Emesini (Riodinidae), 99 Emmeliinae (= Acontiinae), 125 Empedopsychinae (= Oiketicinae), 27 Emplociidae (= Ennominae), 93 Emplociini (Geometridae), 93 Emydiidae (= Arctiidae), 124 Endotheniini (= Bactrini), 84 Endotrichinae (= Endotrichini), 61 Endotrichini (Pyralidae), 61 ENDROMIDAE, 113 Endromididae (= Endromidae), 113 Endrominae (Endromidae), 113 Endrosinae (= Endrosini), 124 Endrosini (Arctiidae), 124 Enicostominae (= Depressariinae), 34 Ennomidae (= Ennominae), 93 Ennominae (Geometridae), 93 Ennomini (Geometridae), 93 Ennychiidae (= Pyraustinae), 61 Ennychiinae (= Pyraustinae), 61 Entephriinae (= Larentiini), 93 Eotaleporiini (Psychidae), 27 EPERMENIIDAE, 48 Epermeniinae (Epermeniidae), 48 Ephestiini (= Phycitini), 61 Ephyridae (= Cosymbiini), 93 Ephyrini unavail. (= Cosymbiini), 93 Epialidae [sic] (= Hepialidae), 12 Epibleminae (= Eucosmini), 84 Epicaliina (Nymphalidae), 101 Epicaliini (= Limenitidini), 101 Epichnopterigidae (= Psychidae), 27 Epichnopteriginae (= Psychinae), 27 Epichnopterygidae [sic] (= Psychidae), 27 Epichnopteryginae [sic] (= Psychinae), 27 Epichnopterygini (Psychidae), 27 Epicnopterygidae [sic] (= Psychidae), 27 Epicnopterygidae (= Heterogynidae), 69 Epicominae (Thaumetopoeinae), 120 EPICOPEIIDAE, 88 Epicopeiinae (Epicopeiidae), 88 Epigraphiidae (= Epigraphiina), 34 Epigraphiinae (= Depressariinae), 34 Epigraphiini (= Epigraphiina), 34 Epigraphiina (Oecophoridae), 34 Epiinae (Apatelodidae), 108 Epimarptini (Coleophoridae), 40 Epinephelini (= Maniolina), 101 Epionidae (= Epionini), 93

Epionini (Geometridae), 93 Epipaschiinae (Pyralidae), 61 Epiphilina (Nymphalidae), 101 Epiplemidae (= Eucosmini), 84 EPIPLEMIDAE, 92 Epipleminae (Epiplemidae), 92 EPIPYROPIDAE, 79 Epipyropinae (Epipyropidae), 79 Epirranthini (Geometridae), 93 Epirrhoinae (= Xanthorhoini), 93 Episemidae (= Oncocnemidini), 125 Epitimyrini (Tortricidae), 84 Eranninae (= Bistonini), 93 Erannini (= Erannina), 93 Erastrianae [sic] (= Acontiinae), 125 Erastriidae (= Acontiinae), 125 Erastriinae (= Acontiinae), 125 Erastriinae (= Caberini), 93 Erastriini (= Caberini), 93 Erebidae (= Catocalinae), 125 Erebiina (Nymphalidae), 101 Erebiinae [sic] (= Catocalinae), 125 Erebiinae (= Satyrini), 101 Erebiini (= Erebiina), 101 Erebinae (= Catocalinae), 125 Erechthiidae (= Erechthiinae), 25 Erechthiinae (Tineidae), 25 Eriocephalidae (= Micropterigidae), 1 ERIOCOTTIDAE, 26 Eriocottinae (Eriocottidae), 26 Eriocraniadae [sic] (= Eriocraniidae), 4 ERIOCRANIIDAE, 4 ERIOCRANIOIDEA, 4 ERIOCRANIOIDEA, 4
Eriopidae (= Eriopini), 125
Eriopini (Noctuidae), 125
Eriopygini (Noctuidae), 125
Erupini (Pyralidae), 61
Erycinidae (= Riodinidae), 99
Erycininae (= Riodininae), 99 Erynninae (= Pyrginae), 95 Ethmiidae (= Ethmiinae), 34 Ethmiinae (Oecophoridae), 34

Etimonotrysia (= Tischerioidea), 17 Eubagina (Nymphalidae), 101 Eublemminae (= Eublemmini), 125 Eublemmini (Noctuidae), 125 Euboliidae (= Xanthorhoini), 93 Euboliinae (= Xanthorhoini), 93 Eubolini (= Xanthorhoini), 93 Eubyjinae (= Bistonini), 93 Eubyjini (= Bistonina), 93 Eucestidae (= Gracillariidae), 33 Euchalciina (= Euchalciina), 125 Euchalciina (Noctuidae), 125 Eucheliidae (= Arctiidae), 124 Euchloeini (= Anthocharini), 97 Euchloini (= Anthocharini), 97 Euchlorini (= Comibaenina), 93 Euchromiinae (= Euchromiini), 124 Euchromiini (Arctiidae), 124 Euclastinae (= Pyraustinae), 61 Eucleidae (= Limacodidae), 81 Euclidiidae (= Euclidiina), 125 Euclidiini (= Euclidiina), 125 Euclidiina (Noctuidae), 125 Eucnemidophorinae (= Pterophorinae), 64 Eucocytiini (Noctuidae), 125 Eucosmidae (= Eucosmini), 84 Eucosminae (= Eucosmini), 84 Eucosmini (Tortricidae), 84 Eudamini (Hesperiidae), 95 Eudesmiini (Arctiidae), 124 Eudoniini (Pyralidae), 61 Eudulini (Geometridae), 93 Eulechriadae [sic] (= Oecophoridae), 34 Eulechriidae (= Oecophoridae), 34 Eulepidoptera (= Heteroneura), 15 Euliini (Tortricidae), 84 Eumaeini (Lycaenidae), 98 Eumaeina (Lycaenidae), 98 Eumeleinae (= Eumeleini), 93 Eumeleini (Geometridae), 93 Eumeninae (= Satyrini), 101 Euphydryina (Nymphalidae), 101 Euphydryini (= Melitaeini), 101 Euphyiini (= Xanthorhoini), 93 Eupistidae (= Coleophoridae), 40

Eupitheciinae (= Eupitheciini), 93

Eupitheciini (Geometridae), 93 Euplocamidae (= Euplocaminae), 25 Euplocaminae (Tineidae), 25 Euploeini (Nymphalidae), 101 Euploeina (Nymphalidae), 101 Euproctidini (Lymantriidae), 123 EUPTEROTIDAE, 107 Eupterotinae (Eupterotidae), 107 Euptychiina (Nymphalidae), 101 Eurhipidae (= Euteliinae), 125 Eurhipiinae (= Euteliinae), 125 Eurrhypinae (= Eurrhypini), 61 Eurrhypini (Pyralidae), 61 Eurybiini (Riodinidae), 99 Eurytelina (Nymphalidae), 101 Eurytelini (= Limenitidini), 101 Eurythmasiini (= Phycitini), 61 Euschemidae (= Dysphaniini), 93 Euschemini (= Dysphaniini), 93 Euselasiinae (Riodinidae), 99 Eustrotiinae (= Eustrotiini), 125 Eustrotiini (Noctuidae), 125 Eustrotiini (= Eublemmini), 125 Euteliinae (Noctuidae), 125 Euthaliini (Nymphalidae), 101 Euthalini [sic] (= Euthaliini), 101 Eutoeini (Geometridae), 93 Euxoinae (= Noctuinae), 125 Euzopherini (= Phycitini), 61 Evergestinae (Pyralidae), 61 Evergestini (Pyralidae), 61 Everinae (= Everina), 98 Everini (= Everina), 98 Everina (Lycaenidae), 98

Exapatidae (= Cnephasiini), 84 EXOPORIA, 8 Falkovitshiinae (= Coleophorinae), 40 Feraliini (Noctuidae), 125 Fernaldellinae (= Semiothisini), 93 Fernaldellini (= Semiothisini), 93 Fidoniidae (= Boarmiini), 93 Fidoniinae (= Boarmiini), 93 Fidoniini (= Boarmiina), 93 Fumariinae (= Psychinae), 27 Fumeidae (= Psychidae), 27 Fumeinae (= Psychinae), 27 Galacticinae (Urodidae), 67 Galleriidae (= Galleriinae), 61 Galleriinae (Pyralidae), 61 Galleriini (Pyralidae), 61 Gastropachini (Lasiocampidae), 112 Gatesclarkeanini (Tortricidae), 84 Gazerini (Castniidae), 83 GELECHIIDAE, 38 Gelechiinae (Gelechiidae), 38 Gelechiini (Gelechiidae), 38 GELECHIOIDEA, 34 GEOMETRIDAE, 93 Geometrinae (Geometridae), 93 Geometrini (Geometridae), 93 Geometrina (Geoemtridae), 93 GEOMETROIDEA, 93 Glaphyriinae (Pyralidae), 61 Glaphyriini (Pyralidae), 61 Glaucopsychinae (= Scolitantidina), 98 Glaucopsychini (= Scolitantidina), 98 GLOSSATA, 4 Glottulidae (Glottulini), 125 Glottulinae (Glottulini), 125 Glottulinie (Glottulini), 125 Glottulini (Noctuidae), 125 Glyphidocerini (Oecophoridae), 34 GLYPHIPTERIGIDAE, 50 Glyphipteriginae (Glyphipterigidae), 50
Glyphipterigoidea (= Yponomeutoidea), 49 Glyphipterygidae [sic] (= Glyphipterigidae), Gnophidae (= Boarmiini), 93 Gnophini (= Gnophina), 93 Gnophina (Geometridae), 93 Gnorimoschemini (Gelechiidae), 38 Godyridini (Nymphalidae), 101 Gonepterygini (Pieridae), 97 Goniacidaliini (= Sterrhini), 93 Gonodontini (Geometridae), 93 Gonometini (Lasiocampidae), 112 Gonopteridae (= Ophiderina), 125 Gonopterini (Noctuidae), 125 Gortynidae (= Eriopini), 125

Gortyninae (= Eriopini), 125 Gracilaridae [sic] (= Gracillariidae), 33 Gracilariidae [sic] (= Gracillariidae), 33 GRACILLARIIDAE, 33 Gracillariiformes, 29 Gracillariinae (Gracillariidae), 33 Gracillarioidea (= Gracillariiformes). 29 Graellsiini (= Saturniini), 118 Graphiina (Papilionidae), 96 Graphiini (= Graphiina), 96 Grapholithidae [sic] (= Grapholitini), 84 Grapholithinae [sic] (= Grapholitini), 84 Grapholitinae (= Grapholitini), 84 Grapholitini (Tortricidae), 84 Griveaudiinae (Callidulidae), 87 Grotellini (Noctuidae), 125 Habrosynini (Thyatiridae), 103 Hadapameini (= Apameini), 125 Hadenidae (= Hadeninae), 125 Hadeninae (Noctuidae), 125 Hadenini (Noctuidae), 125 Haemilidae (= Oecophoridae), 34 Haemylidae [sic] (= Oecophoridae), 34 Haeterini (Nymphalidae), 101 Haimbachiini (Pyralidae), 61 Hamearinae (Riodinidae), 99 Hamearinae (= Riodininae), 99 Hapaliinae (= Pyraustinae), 61 Hapigiini (Notodontidae), 120 Haploptilidae (= Coleophoridae), 40 Hapsiferinae (= Hapsiferini), 25 Hapsiferini (Tineidae), 25 Harmaclonidae (= Harmacloninae). 25 Harmacloninae (Tineidae), 25 Hazidae (= Dysphaniini), 93 Hazini (= Dysphanini), 93
Hazini (= Dysphanini), 93
Heccmeyerianae [sic] (= Psychinae), 27
Heckmeyeriinae (= Psychinae), 27
HEDYLIDAE, 94 Hedyloidea (= Hedylidae), 94 Heliconiidae (= Heliconiinae), 101 Heliconiinae (Nymphalidae). 101 Heliconiini (Nymphalidae). 101 Helicopini (Riodinidae), 99 HELIODINIDAE, 57 Heliotheiini [sic] (= Rhomboristina). 93 Heliotheini (= Rhomboristina), 93 Heliothelinae (= Heliothelini), 61 Heliothelini (Pyralidae), 61 Heliothidae (= Heliothinae), 125 Heliothidinae (= Heliothinae), 125 Heliothinae (Noctuidae), 125 HELIOZELIDAE, 23 Hemerophilidae (= Choreutidae), 68 Hemiceratinae (= Hemiceratini), 120 Hemiceratini (Notodontidae). 120 Hemileucinae (Saturniidae), 118 Hemileucini (Saturniidae), 118 Hemistolini (= Hemitheina), 93 Hemitheidae (= Hemitheina), 93 Hemitheinae (= Hemitheina), 93 Hemitheini (= Hemitheina), 93 Hemitheina (Geometridae), 93 Heodinae (= Lycaeninae), 98 Heodininae (= Lycaeninae), 98 HEPIALIDAE, 12 HEPIALOIDEA, 9 Hepiolidae [sic] (= Hepialidae). 12 Hercynidae (= Hercynini), 61 Hercyninae (= Hercynini), 61 Hercynini (Pyralidae), 61 Herminiidae (= Herminiinae), 125 Herminiinae (Noctuidae), 125 Herminiinina (Noctuidae), 125 Herrichiinae (= Herrichiina), 34 Herrichini (= Herrichina). 34 Herrichina (Oecophoridae). 34 HESPERIIDAE, 95 Hesperiiformes, 95 Hesperiinae (Hesperiidae), 95 Hesperioidea (= Hesperiiformes), 95 HETEROBATHMIIDAE, 3 HETEROBATHMIINA, 3 HETEROBATHMIOIDEA, 3 Heterocampinae (Notodontidae), 120 Heterocampini (Notodontidae), 120 Heterogeneidae (= Limacodidae), 81 HETEROGYNIDAE, 69

HETERONEURA, 15 Heteropsychinae (Epipyropidae), 79 Heteropterinae (Hesperiidae), 95 Hiberniidae [sic] (= Bistonini), 93 Hibrildinae (Pterothysanidae), 86 Hibrildini (Pterothysanidae), 86 Hierochthoniini (= Hierochthoniina), 93 Hierochthoniina (Geometridae), 93 Hieroxestidae (= Hieroxestinae), 25 Hieroxestinae (Tineidae), 25 Hilarographini (Tortricidae), 84 HIMANTOPTERIDAE, 71 Hipparchiidae (= Satyrinae), 101 Hipparchiinae (= Satyrini), 101 Hipparchiini (= Hipparchiina), 101 Hipparchiina (Nymphalidae), 101 Hoffmanniinae (= Yponomeutinae), 53 Holcocerinae (Blastobasidae), 39 Holcopogonidae (= Symmocini), 34 Hollandiidae (= Metarbelidae), 77 Holocerinae (= Ludiinae), 118 Hotocerinae (= Ludiinae), 118
Homalochroinae (= Pyralinae), 61
Homoeosomini (= Phycitini), 61
Homophysidae (= Glaphyrinae), 61
Horagini (Lycaenidae), 98
Horagina (Lycaenidae), 98
Hyberniini (= Bistonina), 93
HYBLAEIDAE, 59
Hydriamainae (= Hydriamanini) 01 Hydriomeninae (= Hydriomenini), 93 Hydriomenini (Geometridae), 93 Hydririni (Pyralidae), 61 Hydrocampidae (= Nymphulinae), 61 Hydrocampinae (= Nymphulinae), 61 Hyemidae [sic] (= Oecophoridae), 34 Hylophilinae (= Chloephorinae), 125 Hymeniinae (= Pyraustinae), 61 Hypatiminae (= Chelariini), 38 Hypenidae (= Hypeninae), 125 Hypeninae (Noctuidae), 125 Hypenodinae (Noctuidae), 125 Hypenodini (Noctuidae), 125 Hypercalliinae (= Hypercalliini), 34 Hypercalliini (Oecophoridae), 34 Hypertrophidae (= Hypertrophinae), 34 Hypertrophinae (Oecophoridae), 34 Hypochrosidae (= Anagogini), 93 Hypochrosini (= Anagogini), 93 Hypocystina (Nymphalidae), 101 Hypogymnidae (= Lymantriidae), 123 Hypolimnini (= Nymphalini), 101 Hypolycaenina (Lycaenidae), 98 Hyponomeutidae [sic] (= Yponomeutidae), Hyponomeutinae [sic] (= Yponomeutinae), Hypoptinae (Cossidae), 75 Hypotheclina (Lycaenidae), 98 Hypotiinae (= Pyralinae), 61 Hypsidae (= Aganainae), 125 Hypsidophidae [sic] (= Plutellidae), 51 Hypsilophinae [sic] (= Ypsolophinae), 51 Hypsotropinae (= Phycitinae), 61 Hypsotropinae (= Peoriini), 61 Hysterosiinae unavail. (= Cochylini), 84 Hysterosiini (= Cochylini), 84 Idaeidae (= Sterrhini), 93 Idaeini (= Sterrhini), 93 Ideina (Nymphalidae), 101 IMMIDAE, 58 Imminae (Immidae), 58 IMMOIDEA, 58 INCURVARIIDAE, 19 **INCURVARIINA**, 19 Incurvariinae (Incurvariidae), 19 **INCURVARIOIDEA**, 19 Infurcitineinae (= Infurcitineini), 25 Infurcitineini (Tineidae), 25 Iolaini (Lycaenidae), 98 Iolaina (Lycaenidae), 98 Ipimorphinae (= Hadeninae), 125 Ipimorphini (= Hadenini), 125 Ithomiidae (= Ithomiinae), 101 Ithomiinae (Nymphalidae), 101 Ithomiini (Nymphalidae), 101 Itunina (Nymphalidae), 101 Janinae (Eupterotidae), 107 Janiodinae (Cercophanidae), 117

Jaspidiinae (= Acontiinae), 125 Jodiini [sic] (= Hemitheina), 93 Jodini (= Hemitheina), 93 Kallimini (Nymphalidae), 101 Lacariini (= Lithinini), 93 Lachneidae (= Lasiocampidae), 112 Lachnocnemini (Lycaenidae), 98 Lacosomidae (= Mimallonidae), 110 LACTURIDAE, 72 Lampidina (Lycaenidae), 98 Lampidinae (= Lampidina), 98 Lampidini (= Lampidina), 98 Lamproniidae (= Prodoxidae), 21 Lamproniinae (Prodoxidae), 21 Lampropterinae (= Leptocircini), 96 Lampropterini (Lampropterina), 96 Lamprysticini (Oecophoridae), 34 Larentiidae (= Larentiinae), 93 Larentiinae (Geometridae), 93 Larentiini (Geometridae), 93 Lariidae (= Lymantriidae), 123 LASIOCAMPIDAE, 112 Lasiocampinae (Lasiocampidae), 112 Lasiocampini (Lasiocampidae), 112 Lasiocampoidea (= Lasiocampidae), 112 Laspeyresiinae (= Lastocampidae) Laspeyresiini (= Grapholitini), 84 Laspeyresiini (= Grapholitini), 84 Lathrotelidae (= Nymphulinae), 61 Lavernidae (= Momphidae), 41 LECITHOCERIDAE, 35 Lecithocerinae (Lecithoceridae), 35 Lemoniadae [sic] (= Lemoniidae), 114 Lemoniadina (Riodinidae), 99 Lemoniadini (= Lemoniadina), 99 LEMONIIDAE, 114 Lemoniinae (Lemoniidae), 114 Lemoniini (= Lemoniadini), 99 Lepidarbelidae (= Metarbelidae), 77 Leptideinae (= Leptideini), 97 Leptideini (Pieridae), 97 Leptocercinae (= Leptocircini), 96 Leptocircini (Papilionidae), 96 Leptopteriginae (= Oiketicinae), 27 Leptopteryginae [sic] (= Oiketicinae), 27 Lethini (Nymphalidae), 101 Leucanidae (= Hadenini), 125 Leucaniinae (= Hadeninae), 125 Leucomidae (= Lymantriidae), 123 Leuconyctini (Noctuidae), 125 Leucopteriginae (= Cemiostominae), 55 Leucopterinae (= Cemiostominae), 55 LIBYTHEIDAE, 100 Libytheinae (= Libytheidae), 100 Ligidae [sic] (= Prosopolophini), 93 Ligiidae (= Prosopolophini), 93 Ligiini unavail. (= Prosopolophini), 93 LIMACODIDAE, 81 Limacodiformes, 78 Limacodoidea (= Limacodiformes), 78 Limenitidinae (Limenitidini), 101 Limenitidini (Nymphalidae), 101 Limenitinae (Limenitidini), 101 Limenitini [sic] (= Limenitidini), 101 Linostinae (Pyralidae), 61 Liparidae (= Lymantriidae), 123 Liphyrinae (Lycaenidae), 98 Lipteninae (Lycaenidae), 98 Liptenini (Lycaenidae), 98 Lipusinae [sic] (= Lypusinae), 27 Lithinini (Geometridae), 93 Lithocolletidae (= Lithocolletinae), 33 Lithocolletinae (Gracillariidae), 33 Lithophanini (= Xylenini), 125 Lithosiidae (= Lithosiinae), 124 Lithosiinae (Arctiidae), 124 Lithosiini (Arctiidae), 124 Litidae (Gelechiidae), 38 Lobophorinae (= Trichopterygini), 93 Lobophorini (= Trichopterygini), 93 Lophochoristini (= Lophochoristina), 93 Lophochoristina (Geometridae), 93 LOPHOCORONIDAE, 6 LOPHOCORONINA, 6 LOPHOCORONOIDEA, 6 Loxurina (Lycaenidae), 98 Lozoperidae (= Cochylini), 84 Luciini (Lycaenidae), 98 Ludiinae (Saturniidae), 118

Luehdorfiinae (= Zerynthiini), 96 Luehdorfiini (= Zerynthiini), 96 Luffiidae (= Psychidae), 27 Luffiinae (= Psychinae), 27 Lycaenesthini (Lycaenidae), 98 LYCAENIDAE, 98 Lycaeninae (Lycaenidae), 98 Lycaenoidea (= Lycaenidae), 98 Lycaenopsinae (= Lycaenopsina), 98 Lycaenopsini (= Lycaenopsina), 98 Lycaenopsina (Lycaenidae), 98 Lycoreinae (= Lycoreini), 101 Lycoreini (Nymphalidae), 101 Lycoreina (Nymphalidae), 101 LYMANTRIIDAE, 123 Lymantriinae (Lymantriidae), 123 Lymantriini (Lymantriidae), 123 LYONETIIDAE, 55 Lyonetiinae (Lyonetiidae), 55 Lypusidae (= Lypusinae), 27 Lypusinae (Psychidae), 27 Lyrceini unavail. (= Oenochromini), 93 Lythriini (Geometridae), 93 Macariidae (= Semiothisini), 93 Macariinae (= Semiothisini), 93 Macariini (= Semiothisini), 93 Macroglossinae (Sphingidae), 119 Macroglossini (Sphingidae), 119 Macromphaliinae (Lasiocampidae), 112 Macropiratinae (Pterophoridae), 64 Macrothecinae (= Galleriinae), 61 Macrothecini (= Cacotherapiini), 61 Macrothyatirini (Thyatiridae), 103 Madopidae (= Ophiderini), 125 Malacosomatinae (= Malacosomatini), 112 Malacosomatini (Lasiocampidae), 112 Maniidae (= Eriopini), 125 Maniolinae (= Satyrini), 101 Maniolini (= Maniolina), 101 Maniolina (Nymphalidae), 101 Marasmarchinae (= Pterophorinae), 64 Margarodidae (= Pyraustinae), 61 Margarodinae (= Pyraustinae), 61 Margaroniidae (= Pyraustinae), 61 Margaroniinae (= Pyraustinae), 61 Marpesiinae (= Marpesiini), 101 Marpesiini (= Cyrestidini), 101 Mechanitini (Nymphalidae), 101 Mecoceratini (= Ametridini), 93 Meessiinae (Tineidae), 25 Meessiini (Tineidae), 25 Megalophaninae (= Oiketicinae), 27 MEGALOPYGIDAE, 74 Megalopygina (= Megalopygidae), 74 Megalopyginae (Megalopygidae), 74 Megarthridini (Pyralidae), 61 Megathyminae (Hesperiidae), 95 Megathymini (Hesperiidae), 95 Meiceratinae (Notodontidae), 120 Melanargiinae (= Satyrini), 101 Melanargiina (Nymphalidae), 101 Melanchroiinae (= Melanchroiini), 93 Melanchroiini (Geometridae), 93 Melanippidae (= Rheumapterini), 93 Melanitini (Nymphalidae), 101 Melanolophidae (= Olethreutini), 84 Melanolophiini (Geometridae), 93 Melanthiini (Geometridae), 93 Melasininae (= Penestoglossinae), 27 Melicleptriinae (= Heliothinae), 125 Melinaeini (Nymphalidae), 101 Melitaeinae (= Melitaeini), 101 Melitaeina (Nymphalidae), 101 Melitaeini (Nymphalidae), 101 Melittiini (Sesiidae), 66 Mendesiini (= Perittiinae), 36 Mesosemiina (Riodinidae), 99 Mesosemiini (Riodinidae), 99 Metachandidae (= Oecophoridae), 34 Metachrostinae (= Bryophilinae), 125 METARBELIDAE, 77 Metasini (Psychidae), 27 Methonini (= Mechanitini), 101 Metriotinae (= Coleophorinae), 40 Metrocampidae (= Campaeini), 93 Metrocampini (= Campaeini), 93 Metzneriidae (Gelechiidae), 38 Metzneriinae (= Anomologini), 38

Micrarctiinae (= Callimorphini), 124 Microcorsini (Tortricidae), 84 Microloxiini (= Miroloxiina), 93 Microloxiina (Geometridae), 93 Microniinae (Uraniidae), 91 Micropinidae (= Cyllopodini), 93 Micropinini unavail. (= Cyllopodini), 93 Micropsychiidae (= Psychidae), 27 Micropsychinidae (= Psychidae), 27 MICROPTERIGIDAE, 1 Micropteriginae (Micropterigidae), 1 MICROPTERIGOIDEA. I Micropterygidae [sic] (= Micropterigidae), 1 Midilinae (Pyralidae), 61 Miletinae (Lycaenidae), 98 Miletini (Lycaenidae), 98 Miletina (Lycaenidae), 98 Milioniini (Geometridae), 93 Millieriinae (Choreutidae), 68 MIMALLONIDAE, 110 Mirininae (Endromidae), 113 MNESARCHAEIDAE, 8 MNESARCHAEOIDEA, 8 Moffatiinae (= Psychinae), 27 Mominae (= Paythnae), 25 MOMPHIDAE, 41 Monocteniadae (= Oenochromini), 93 Monocteniini (= Oenochromini), 93 Monopidae (= Tineidae), 25 Monopinae (= Tineinae), 25 MONOTRYSIA, 15 Morphidae (= Morphinae), 101 Morphinae (Nymphalidae), 101 Morphini (Nymphalidae), 101 Munychryiinae (Anthelidae), 111 Musotimidae (= Musotiminae), 61 Musotiminae (Pyralidae), 61 Mycalesini (Nymphalidae), 101 Myelobiini (Pyralidae), 61 MYOGLOSSATA, 7 Myrmecozelinae (Tineidae), 25 Myrmecozelini (Tineidae), 25 Nacophorini (Geometridae). 93 Napaeina (Riodinidae), 99 Napeogenini (Nymphalidae), 101 Naropini (Nymphalidae), 101 Naryciidae (= Taleporiinae), 27 Naryciinae (= Taleporiinae), 27 Naryciini (Psychidae), 27 Nemapogoninae (Tineidae), 25 Nematophoridae [sic] (= Adelidae), 22 Nematopogoninae (Adelidae), 22 Nemeobiidae (= Riodinidae), 99 Nemeobiinae (= Riodininae), 99 Nemophoridae (= Adelidae), 22 Nemoriinae (= Nemoriina), 93 Nemoriini (= Nemoriina), 93 Nemoriina (Geometridae), 93 Neoblastobasini (Blastobasidae), 39 Neocastniinae (Castniidae), 83 Neohipparchini (= Neohipparchina), 93 Neohipparchina (Geometridae), 93 NEOLEPIDOPTERA, 8 NEOPSEUSTIDAE, 7 NEOPSEUSTINA, 7 NEOPSEUSTOIDEA, 7 NEOTHEORIDAE, 9 Nephodiinae (= Ourapterygini), 93 NEPTICULIDAE, 15 NEPTICULINA, 15 Nepticulinae (Nepticulidae), 15 Nepticulini (Nepticulidae), 15 NEPTICULOIDEA, 15 Neptini (Nymphalidae), 101 Niasomini (Tortricidae), 84 Nidarinae (Drepanidae), 105 Niphonymphinae (= Niphonymphini), 53 Niphonymphini (Yponomeutidae), 53 Nocloini (Noctuidae), 125 NOCTUIDAE, 125 Noctuinae (Noctuidae), 125 Noctuini (Noctuidae), 125 Noctuinina (Noctuidae), 125 NOCTUOIDEA, 120 Nolidae (= Nolinae), 125 Nolinae (Noctuidae), 125 Nomophilinae (= Pyraustinae), 61 Nonagriinae (= Eriopini), 125

Noordinae (Pyralidae), 61 NOTODONTIDAE, 120 Notodontinae (Notodontidae), 120 Notodontini (Notodontidae), 120 Notodontinina (Notodontidae), 120 Nudariinae (= Nudariini), 124 Nudariini (Arctiidae), 124 Nyctemerinae (= Callimorphini), 124 Nyctemerini (= Callimorphini), 124 Nyctemerina (Arctiidae), 124 Nycteolidae (= Sarrothripiniae), 125 Nycteolinae (= Sarrothripiniae), 125 NYMPHALIDAE, 101 Nymphalinae (Nymphalidae), 101 Nymphalini (Nymphalidae), 101 Nymphalini (Nymphalidae), 101 Nymphaloidea (= Nymphalidae), 101 Nymphaloidea (= Nymphalidae), 101 Nymphidiina (Riodinidae), 99 Nymphidiini (Riodinidae), 99 Nymphuliniae (Pyralidae), 61 Nymphulini (Pyralidae), 61 Nystaleinae (Notodontidae), 120 Obeidiini (Geometridae), 93 Ochrognesiini (= Nemoriina), 93 Ochromolopinae (Epermeniidae), 48 OCHSENHEIMERIIDAE, 49 Ochyroticinae (Pterophoridae), 64 Oditinae (Lecithoceridae), 35 Oditini (Lecithoceridae), 35 Odontiinae (Pyralidae), 61 Odontiini (Pyralidae), 61 Odontodinae (= Stictopterinae), 125 Odontoperinae (= Odontoperini), 93 Odontoperini (= Ennomini), 93 Odontoperini (Geometridae), 93 Odopteridi (= Ennomini), 93 Odopterini (= Ennomini), 93 Oeciini (Oecophoridae), 34 OECOPHORIDAE, 34 Oecophorinae (Oecophoridae), 34 Oecophorini (Oecophoridae), 34 Oecophorina (Oecophoridae), 34 Oecophoroidea (= Gelechioidea), 34 Oecophyllembiinae (= Gracillariinae), 33 Oegoconiinae (= Oegoconiini), 34 Oegoconiini (Oecophoridae), 34 Oeneini (= Erebiina), 101
Oenochrominae (= Oenochromini), 93
Oenochrominae (Geometridae), 93 Oenochromini (Geometridae), 93 Oenophilidae [sic] (= Hieroxestinae), 25 Oenosandrinae (Notodontidae), 120 Oenosandrinina (Notodontidae), 120 Ogyrina (Lycaenidae), 98 Oiketicidae (= Oiketicinae), 27 Oiketicinae (Psychidae), 27 Oiketicini (Psychidae), 27 Oiketicoidinae (= Oiketicinae), 27 Oinophilidae (= Hieroxestinae), 25 Oinophilinae (= Hieroxestinae), 25 Oleriini (Nymphalidae), 101 Olethreutidae (= Olethreutinae), 84 Olethreutinae (Tortricidae), 84 Olethreutini (Tortricidae), 84 Oncocnemidini (Noctuidae), 125 Onychorini (Geometridae), 93 Operophterinae (= Operophterini), 93 Operophterini (Geometridae), 93 Operophtherini [sic] (= Operophterini), 93 Ophideridae (= Ophiderini), 125 Ophiderinae (= Catocalinae), 125 Ophiderini (Noctuidae), 125 Ophiderini (= Ophiderina), 125 Ophiderina (Noctuidae), 125 Ophiusidae (= Ophiusina), 125 Ophiusini (= Ophiusina), 125 Ophiusina (Noctuidae), 125 Opogoninae (= Hieroxestinae), 25 Oporiniinae (= Operophterini), 93 Oporiniini (= Operophterini), 93 OPOSTEGIDAE, 16 Oposteginae (Opostegidae), 16 Opostegoidinae (Opostegidae), 16 Orenaiini (Pyralidae), 61 Oreopsychinae (= Oiketicinae), 27 Oreopsychini (Psychidae), 27 Oretinae (Drepanidae), 105 Orgyiidae (= Orgyiinae), 123

Orgyiinae (Lymantriidae), 123 Orgyiini (= Orgyiinae), 123 Orgyinae [sic] (= Orgyinae), 123 Orneodidae (= Alucitidae), 46 Ornichichinae [sic] (= Gracillariinae), 33 Ornichinae (= Gracillariinae), 33 Ornigidae (= Gracillariidae), 33 Orophiinae (= Cryptolechiini), 34 Orophiini (= Chryptolechiina), 34 Ortholithinae (= Xanthorhoini), 93 Orthosiidae (= Eriopini), 125 Orthostixidae (= Orthostixinae), 93 Orthostixinae (Geometridae), 93 Orthostixini (= Orthostixinae), 93 Orthotaeliinae [sic] (= Orthoteliinae), 50 Orthoteliidae (= Glyphipterigidae), 50 Orthoteliinae (Glyphipterigidae), 50 Osminiini (Sesiidae), 66 Othreinae (= Ophiusina), 125 Ourapteryginae (= Ourapterygini), 93 Ourapterygini (Geometridae), 93 **OXYCHIROTIDAE**, 63 Oxydiidae (= Oxydiini), 93 Oxydiini (Geometridae), 93 Oxylidina (Lycaenidae), 98 Oxyptilinae (= Pterophorinae), 64 OXYTENIDAE, 116 Oxytripiinae (= Eriopini), 125 Pachycnemiidae (= Prosopolophini), 93 Pachycnemiidae (= Prosopolophini), 93 Pachycnemiini (= Prosopolophini), 93 PALAEOSETIDAE, 13 PALAEPHATIDAE, 18 PALAEPHATOIDEA, 18 Palimpsestidae (= Thyatiridae), 103 Palindidae (= Ophiderina), 125 Palpariidae (= Crambinae), 61 Palumbininae (= Anacampsini), 38 Palyadae (= Palyadini), 93 Palyadini (Geometridae), 93 Pamphilinae (= Hesperiinae), 95 Panacelinae (Eupterotidae), 107 Pangaptini [sic] (= Pangraptina), 125 Pangraptina (Noctuidae), 125 Pantheinae (Noctuidae), 125 Pantheini (Noctuidae), 125 Pantheridae (= Abraxini), 93 Pantherini (= Abraxini), 93 PAPILIONIDAE, 96 Papilioniformes, 96 Papilioninae (Papilioniidae), 96 Papilionini (Papilioniidae), 96 PAPILIONOIDEA, 95 Paramesiini (= Cnephasiini), 84 Parametriotinae (= Parametriotini), 42 Parametriotini (Agonoxenidae), 42 Paranthreninae (Sesiidae), 66 Paranthrenini (Sesiidae), 66 Parargina (Nymphalidae), 101 Pararginae (= Elymniini), 101 Parargini (= Paragina), 101 Parathyridae (= Arrhenophanidae), 28 Pardopsini (Nymphalidae), 101 Parnassiidae (= Parnassiinae), 96 Parnassiinae (Papilioniidae), 96 Parnassiini (Papilioniidae), 96 Pectinivalvinae (Nepticulidae), 15 Peleopodidae (= Peleopodini), 34 Peleopodinae (Oecophoridae), 34 Peleopodini (Oecophoridae), 34 Peloponnesini (Psychidae), 27 Penestoglossinae (= Penestoglossini), 27 Pennisetiini (Sesiidae), 66 Penthininae (= Eucosmini), 84 Pentilini (Lycaenidae), 98 Peoriinae (= Peoriini), 61 Peoriini (Pyralidae), 61 Pericopinae (Arctiidae), 124 Pericopinina (Arctiidae), 124 Pericymatini (= Polydesmina), 125 Perissomacticini (Tineidae), 25 Perissomasticinae (= Perissomasticini), 25 Perittiinae (Elachistidae), 36 Perittiini (= Perittiinae), 36 Perizomini (Geometridae), 93 Peroneidae (= Tortricini), 84 Perophoridae (= Mimallonidae), 110

Pexicopiinae (Gelechiidae), 38

Pexicopiini (Gelechiidae), 38 Phaegopterini (Arctiidae), 124 Phalacropterygini (Psychidae), 27 Phalerinae (Notodontidae), 120 Phaloniidae (= Cochylini), 84 Phaseliinae (= Phaseliina), 93 Phaseliini (= Phaseliina), 93 Phaseliina (Geometridae), 93 Phaselini [sic] (= Phaseliina), 93 Phaudinae (Zygaenidae), 70 Phiditiinae (Apatelodidae), 108 Philampeliadae [sic] (= Philampelini), 119 Philampelinae (= Philampelini), 119 Philampelini (Sphingidae), 119 Philereminae (= Rheumapterini), 93 Phileremini (Geometridae), 93 Philobotidae (= Oecophoridae), 34 Philobotinae (= Philobotini), 34 Philobotini (Oecophoridae), 34 Phlogophorinae (= Euteliinae), 125 Phosphilini (Noctuidae), 125 Phricanthini (Tortricidae), 84 Phryganopterygini (Arctiidae), 124 Phthoropoeinae (Tineidae), 25 Phycidae [sic] (= Phycitinae), 61 Phycitale [stc] (= Phycitinae), 6 Phyciodina (Nymphalidae), 101 Phycitidae (= Phycitinae), 61 Phycitinae (Pyralidae), 61 Phycitina (Pyralidae), 61 Phycitina (Pyralidae), 61 Phycodinae (Brachodidae), 65 Phyllobrostidae (= Lyonetiidae), 55 Phyllocnistidae (= Phyllocnistinae), 33 Phyllocnistinae (Gracillariidae), 33 Phyllonorycteridae (= Gracillariidae), 33 Phyllonorycterinae (= Lithocolletinae), 33 Phylloporiidae (= Incurvariidae), 19 Phyllorycteridae [sic] (= Gracillariidae), 33 Physoptilidae (= Physoptilinae), 38 Physoptilinae (Gelechiidae), 38 Phytometrinae (= Plusiinae), 125 PIERIDAE, 97 Pierididae (= Pieridae), 97 Pierinae (Pieridae), 97 Pierini (Pieridae), 97 Pigritiini (Blastobasidae), 39 Pingasini unavail. (= Pseudoterpnina), 93 Placodominae (= Taleporiinae), 27 Placodomini (Psychidae), 27 Plastistomidae [sic] (= Oecophoridae), 34 Platychasmatinae (Notodontidae), 120 Platyomidae nom. nud. (= Tortricidae), 84 Platyptericidae (= Drepanidae), 105 Platyptiliinae (= Platyptiliini), 64 Platyptiliini (Pterophoridae), 64 Plebeiinae (= Polyommatini), 98 Plebeiini (= Polyommatina), 98 Plebeiini (= Polyommatina), 98 Plebejinie [sic] (= Polyommatini), 98 Plebejini [sic] (= Polyommatina), 98 Pleurotinae (= Pleurotini), 34 Pleurotini (Oecophoridae), 34 Plusianae [sic] (= Plusiinae), 125 Plusiidae (= Plusiinae), 125 Plusiina (= Plusiina), 125 Plusiinae (Noctuidae), 125 Plusiini (Noctuidae), 125 Plusiina (Noctuidae), 125 PLUTELLIDAE, 51 Plutellinae (Plutellidae), 51 Plutelloidea (= Yponomeutoidea), 49 Plutodini (Geometridae), 93 Pococerinae (= Epipaschiinae), 61 Poecilocampinae (Lasiocampidae), 112 Poeciloptiliinae (= Gracillariinae), 33 Polydesmidae (= Polydesmina), 125 Polydesmini (= Polydesmina), 125 Polydesmina (Noctuidae), 125 Polyommatinae (Lycaenidae), 98 Polyommatini (Lycaenidae), 98 Polyommatina (Lycaenidae), 98 Polyorthini (Tortricidae), 84 Polyplocidae (= Polyplocinae), 103 Polyplocinae (Thyatiridae), 103 Polyplocini (Thyatiridae), 103 Polypogoninae (= Herminiinae), 125 Polythysanini (Saturniidae), 118 Poritiinae (Lycaenidae), 98

Praydinae (Plutellidae), 51

Preponini (Nymphalidae), 101 Prismostictinae (Bombycidae), 109 Problepsini (= Scopulini), 93 Procridae (= Zygaenidae), 70 Procridinae (Zygaenidae), 70 Procridini (Zygaenidae), 70 Procriinae (= Procridinae), 70 Prodidactidae (= Prodidactinae), 58 Prodidactinae (Immidae), 58 PRODOXIDAE, 21 Prodoxinae (Prodoxidae), 21 Pronophilina (Nymphalidae), 101 Prosopolophinae (= Prosopolophini), 93 Prosopolophini (Geometridae), 93 Prothoini (Nymphalidae), 101 PROTOTHEORIDAE, 11 Proutiinae (= Psychinae), 27 Psaphidinae (Noctuidae), 125 Psaphidini (Noctuidae), 125 Pseudapheliniini (= Urotini), 118 Pseudergolini (Nymphalidae), 101 Pseudocossinae (Brachodidae), 65 Pseudopontiinae (Pieridae), 97 Pseudoterpninae (= Pseudoterpnina), 93 Pseutoterpnini (= Pseudoterpnina), 93 Pseutoterpnina (Geometridae). 93 Psycheoidinae (= Psychinae), 27 PSYCHIDAE, 27 Psychideinae (= Psychinae), 27 Psychinae (Psychidae), 27 Psychini (Psychidae), 27 PTEROLONCHIDAE, 37 Pterolonchoidea (= Gelechioidea), 34 PTEROPHORIDAE, 64 Pterophorinae (Pterophoridae), 64 Pterophorini (Pterophoridae), 64 PTEROPHOROIDEA, 62 PTEROTHYSANIDAE, 86 Pterothysaninae (Pterothysanidae), 86 Ptilophorinae (= Notodontini), 120 Ptochopsychidae (= Mimallonidae), 110 Ptychopodinae (= Sterrhini), 93 Ptychopodini (= Sterrhini), 93 Pygaerinae (Notodontidae), 120 Pygaerini (Notodontidae), 120 PYRALIDAE, 61 Pyralididae (= Pyralidae), 61 Pyralidinae (= Pyralinae), 61 Pyralinae (Pyralidae), 61 Pyralini (Pyralidae), 61 Pyralinia (Pyralidae), 61 PYRALOIDEA, 59 Pyraustidae (= Pyraustinae), 61 Pyraustinae (Pyralidae), 61 Pyraustini (Pyralidae), 61 Pyrginae (Hesperiidae), 95 Pyrgini (Hesperiidae), 95 Pyromorphinae (= Procridinae), 70 Pyrrhopyginae (Hesperiidae), 95 Ramapesiini [sic] (= Cnephasiini), 84 Ramapeziini unavail. (= Cnephasiini), 84 Raphiinae (= Raphiini), 125 Raphiini (Noctuidae), 125 RATARDIDAE, 85 Remelanina (Lycaenidae), 98 Rhescyntidae (= Arsenurinae), 118 Rhescyntinae (= Arsenurinae), 118 Rheumapterini (Geometridae), 93 Rhodobatinae (= Myrmecozelinae), 25 Rhodoceridae (= Coliadinae), 97 Rhodocerinae (= Coliadinae), 97 Rhodogastriinae (= Arctiinae), 124 Rhodometrinae (= Rhodometrini), 93 Rhodometrini (Geometridae), 93 Rhodoneurini (Thyrididae), 60 Rhodostrophiicae [sic] (= Rhodostrophiini), Rhodostrophiini (Geometridae), 93
Rhodostrophini [sic] (= Rhodostrophiini), 93 Rhomboristini (= Rhomboristina), 93 Rhomboristina (Geometridae), 93 Ridiaschinidae (= Cecidosidae), 20 RIODINIDAE, 99 Riodininae (Riodinidae), 99 Riodinini (Riodinidae), 99

Risobini (Noctuidae), 125

Rivulidae (= Rivulinae), 125

Rivulinae (Noctuidae), 125

Roeslerstammiidae (= Amphitheridae), 29 Rosemini (Notodontidae), 120 Rumiinae (= Epionini), 93 Rumiini (= Epionini), 93 Ruralinae (= Lycaenidae), 98 Sabaliidae (= Sabaliinae), 114 Sabaliinae (Lemoniidae), 114 Sabatincinae (Micropterigidae), 1 Salassinae (Saturniidae), 118 Saridoscelinae (Yponomeutidae), 53 Sarotini (= Helicopini), 99 Sarrothripidae (= Sarrothripinae), 125 Sarrothripinae (Noctuidae), 125 Sarrothripini (Noctuidae), 125 SATURNIIDAE, 118 Saturniiformes, 113 Saturniinae (Saturniidae), 118 Saturniini (Saturniidae), 118 Saturnioidea (= Saturniiformes), 113 Satyridae (= Satyrinae), 101 Satyrinae (Nymphalidae), 101 Satyrini (Nymphalidae), 101 Satyrinia (Nymphalidae), 101 Satyrinia (Nymphalidae), 101 Satyrina (Nymphalidae), 101 Satyrinae), 101 Scaeosophidae (= Cosmopterigidae), 43 Scardamiinae (= Scardamiini), 93 Scardamiini (Geometridae), 93 Scardiidae (= Scardiinae), 25 Scardiinae (Tineidae), 25 Schistomitrinae (Epicopeiidae), 88 Schistonoeini (Oecophoridae), 34 Schoenobiidae (= Schoenobiinae), 61 Schoenobiinae (Pyralidae), 61 Schoenotenidae (= Schoenotenini), 84 Schoenotenini (Tortricidae), 84 SCHRECKENSTEINIIDAE, 30 Sciaphilidae (= Cnephasiini), 84 Sciaphilinae (= Cnephasiini), 84 Scirpophagidae (= Schoenobiinae), 61 Scoliopteryginae (= Ophirerina), 125 Scoliopterygini (= Ophiderini), 125 Scolitantidini (= Scolitantidina), 98 Scolitantidina (Lycaenidae), 98 Scopariidae (= Scopariinae), 61 Scopariinae (Pyralidae), 61 Scopariini (Pyralidae), 61 Scopulini (Geometridae), 93 Scoriodytinae (= Psychinae), 27 Scoriodytini (Psychidae), 27 Scranciini (Notodontidae), 120 Scythridae [sic] (= Scythrididae), 44 SCYTHRIDIDAE, 44 Scythropiinae (Plutellidae), 51 Scythropiodini (Lecithoceridae), 35 Seleniini (= Ennomini), 93 Selidosemidae (= Boarmiini), 93 Selidoseminae (= Boarmiini), 93 Semanophorae [sic] (= Macroglossinae), 119 Semanophorinae (= Macroglossinae), 119 SEMATURIDAE, 90 Semioscopidae (= Oecophoridae), 34 Semioscopidinae (= Depressariinae), 34 Semioscopinae [sic] (= Depressariinae), 34 Semiothisinae (= Semiothisini), 93 Semiothisini (Geometridae), 93 Semniinae (= Chrysauginae), 61
Sericoridae [sic] (= Eucosmini), 84
Sericorinae [sic] (= Eucosmini), 84
SESIIDAE, 66 SESIINA, 65 Sesiinae (Sesiidae), 66 Sesiini (Sesiidae), 66 SESIOIDEA, 65 Setomorphinae (Tineidae), 25 Siculidae (= Siculodidae), 60 Siculodinae (Thyrididae), 60 Siculodini (Thyrididae), 60 Siginae (= Pyraustinae), 61 Siloscinae (Tineidae), 25 Simaethidae (= Choreutidae), 68 Simaethistinae (Thyrididae), 60 Similipepsini (Sesiidae), 66 Sionidae (= Boarmiini), 93 Sionini (= Gnophina), 93 Smerinthinae (= Smerinthini), 119 Smerinthini (Sphingidae), 119

Solenobiidae (= Taleporiinae), 27

Solenobiinae (= Taleporiinae), 27 Solitaneini nom. nud. (= Operophterini), 93 Solitaneini (Geometridae), 93 Somabrachidae [sic] (= Somabrachyidae), 73 SOMABRACHYIDAE, 73 Spalgini (Lycaenidae), 98 Sparganothidae (= Sparganothini), 84 Sparganothinae (= Spargnaothini), 84 Sparganothini (Tortricidae), 84 Sphacelodini (Geometridae), 93 Sphingicampinae (= Ceratocampinae), 118 SPHINGIDAE, 119 Sphinginae (Sphingidae), 119 Sphingini (Sphingidae), 119 SPHINGOIDEA, 119 Spilomelidae (= Spilomelini), 61 Spilomelinae (= Pyraustinae), 61 Spilomelini (Pyralidae), 61 Spilonotinae (= Eucosmini), 84 Spilosominae (= Arctiinae), 124 Stalachtini (Riodinidae), 99 Stamnodini (Geometridae), 93 Stathmopodidae (= Stathmopodini), 34 Stathmopodinae (= Stathmopodini), 34 Stathmopodini (Oecophoridae), 34 Stathmopolitinae (= Tineidae), 25 Stauropini (Notodontidae), 120 Steniidae (= Pyraustinae), 61 Steniinae (= Pyraustinae), 61 Stenomatidae (= Stenomatinae), 34 Stenomatinae (Oecophoridae), 34 Stenomidae [sic] (= Stenomatinae), 34 Stenopterydae (= Pyraustinae), 61 Stenoptiliinae (= Pterophorinae), 64 Stephensiini (Elachistidae), 36 Sterrhinae (Geometridae), 93 Sterrhini (Geometridae), 93 Stichobasinae (= Psychinae), 27 Stichobasini (Psychidae), 27 Stictopterinae (Noctuidae), 125 Stigmellidae (= Nepticulidae), 15 Stigmellinae (= Nepticulinae), 15 Stigmonotidae (= Eucosmini), 84 Stigmonotinae (= Eucosmini), 84 Stilbiidae (= Apameini), 125 Stiriinae (Noctuidae), 125 Stiriini (Noctuidae), 125 Stomopteryginae (= Anacampsini), 38 Strepsimanidae (= Hypenodinae), 125 Strepsimanini (Noctuidae), 125 Striglininae (Thyrididae), 60 Striphnopterygidae (= Eupterotidae), 107 Strymoninae (= Theclini), 98 Strymonini (= Theclini), 98 Styginae (Riodinidae), 99 Syleptinae (= Pyraustinae), 61 Symmachiini (Riodinidae), 99 Symmocidae (= Symmocini), 34 Symmocinae (Autostichinae), 34 Symmocini (Oecophoridae), 34 Synanthedoninae (= Synanthedonini), 66 Synanthedonini (Sesiidae), 66 Synchlorini (= Synchlorina), 93 Synchlorina (Geometridae), 93 Synedini (= Drasteriina), 125 Synemoninae (Castniidae), 83 Syntomidae (= Syntominae), 124 Syntominae (Arctiidae), 124 Syntomini (Arctiidae), 124 Syricorinae (= Eucosmini), 84 Syricorinae (= Eucosmini), 84 Syringopainae (= Syringopaini), 34 Syringopaini (Oecophoridae), 34 Syssphingina (= Ceratocampinae), 118 Syssphingina (= Ceraocampinae), 118
Talaeporiinae [sic] (= Taleporiinae), 27
Talaeporiinae [sic] (= Taleporiinae), 27
Taleporiidae (= Taleporiinae), 27 Taleporiinae (Psychidae), 27 Taleporiini (Psychidae), 27 Tarakina (Lycaenidae), 98 Teichobiidae (= Teichobiinae), 25 Teichobiinae (Tineidae), 25 Teinopalpidae (= Teinopalpini), 96 Teinopalpini (Papilioniidae), 96 Telechrysidini (= Telechrysidina), 34 Telechrysidina (Oecophoridae), 34

Teleiodini (= Gelechiini), 38

Tellervinae (Nymphalidae), 101 Teragridae (= Metarbelidae), 77 Terpnini (= Pseudoterpnina), 93 Terpnini (= Pseudoterpnina), 93 Tetheini (Thyatiridae), 103 Thaidinae (Papilioniinae), 96 Thaiinae (Papilioniinae), 96 Thalamarchellidae (= Oecophoridae), 34 Thalamarchidae (= Oecophoridae), 34 Thalassodini (= Hemitheina), 93 Thalerini (= Hemitheina), 93 Thamnospheciini (= Synanthedonini), 66 Thaumetopoeinae (Notodontidae), 120 Thaumetopoeinina (Notodontidae), 120 Theclinae (Lycaenidae), 98 Theclini (Lycaenidae), 98 Theclina (Lycaenidae), 98 Theopina (Riodinidae), 99 Theriini (Geometridae), 93 Therini (= Cidariini), 93 Thestorinae (= Tomarina), 98 Thinopterygini (Geometridae), 93 THYATIRIDAE, 103 Thyatirinae (Thyatiridae), 103 Thyatirini (Thyatiridae), 103 Thymelicinae (= Hesperiinae), 95 THYRETIDAE, 122 Thyretini (= Thyretidae), 122 Thyridae [sic] (= Thyrididae), 60 THYRIDIDAE, 60 Thyridinae (Thyrididae), 60 Thyroidea (= Thyrididae), 60 Thysaniini (= Thysaniina), 125 Thysaniina (Noctuidae), 125 Tichobiidae [sic] (= Teichobiini), 25 Timandridi (= Timandrini), 93 Timandrini (Geometridae), 93 Timandromorphini (= Timandromorphina), Timandromorphina (Geometridae), 93 Timyridae (= Lecithoceridae), 35 Tinaegeriidae (= Stathmopodini), 34 TINEIDAE, 25 Tineiformes, 24 TINEINA, 24 Tineinae (Tineidae), 25 TINEODIDAE, 62 TINEOIDEA, 24 Tinissinae (= Scardiinae), 25 Tinthiinae (Sesiidae), 66 Tinthiini (Sesiidae), 66 Tirathabini (Pyralidae), 61 TISCHERIIDAE, 17 TISCHERIOIDEA, 17 Tissangini (Pterothysanidae), 86 Titaniini (= Odontiini), 61 Tithoreini (Nymphalidae), 101 Tolleophorini (= Coleophorinae), 40 Tomarina (Lycaenidae), 98 Torodorinae (Lecithoceridae), 35 TORTRICIDAE, 84 Tortricinae (Tortricidae), 84 Tortricini (Tortricidae), 84 Tortricodidae (= Cnephasiini), 84 TORTRICOIDEA, 84 Toxocampidae (= Toxocampina), 125 Toxocampini (= Toxocampina), 125 Toxocampina (Noctuidae), 125 Trapezitinae (Hesperiidae), 95 Trichophoridae unavail. (= Tortricidae), 84 Trichopterygini (Geometridae), 93 Trifurculinae (= Trifurculini), 15 Trifurculini (Nepticulidae), 15 Triocnemidini (Noctuidae), 125 Trochilidae (= Sesiidae), 66 Troidini (Papilioniidae), 96 Troidina (Papilioniidae), 96 Trosiinae (Megalopygidae), 74 Trypaninae (= Cossinae), 75 Typhoniinae (Psychidae), 27 Typhoniini (Psychidae), 27 Tytini (= Tytina), 125 Tytina (Noctuidae), 125 Udeini (= Pyraustini), 61 Ufeinae (= Ufeini), 125 Ufeini (Noctuidae), 125 URANIIDAE, 91 Uraniinae (Uraniidae), 91

URANIOIDEA, 88 Urapteridae (= Ouraptrygini), 93 URODIDAE, 67 Urodinae (Urodidae), 67 Urotini (Saturniidae), 118 Utetheisini (Arctiidae), 124 Uzuchidae (= Xyloryctinae), 34 Vanessidae (= Nymphalinae), 101 Varneriini (= Phycitini), 61 Vestermannianae [sic] (= Westermanniini), 125 Walshiidae (= Cosmopterigidae), 43 Walshiinae (= Chrysopeleiinae), 43 Westermanniinae (= Westermanniini), 125 Westermanniini (Noctuidae), 125 Whalleyaninae (Thyrididae), 60 Whitebreadiini (= Perittiinae), 36 Wilemaniina (Geometridae), 93 Wilemanini [sic] (= Wilemaniina), 93 Wurthiinae (Pyralidae), 61 Xanthiinae (= Acronictinae), 125 Xanthopastinae (= Glottulini), 125 Xanthorhoinae (= Xanthorhoini), 93 Xanthorhoini (Geometridae), 93 Xylenini (Noctuidae), 125 Xylenini (Noctuidae), 125 Xylenina (Noctuidae), 125 Xylinidae (= Xylenini), 125 Xyloryctidae (= Xyloryctinae), 34 Xyloryctiade (= Xyloryctinae), 34 Xyloryctinae (Oecophoridae), 34 Xysmatodominae (= Taleporiinae), 27 YPONOMEUTIDAE, 53 Yponomeutinae (Yponomeutidae), 53 Yponomeutini (Yponomeutidae), 53 YPONOMEUTOIDEA, 49 Ypsolophidae (= Plutellidae), 51 Ypsolophinae (Plutellidae), 51 Ypthimini (Nymphalidae), 101 Zanolidae (= Apatelodidae), 108 Zaretini (Nymphalidae), 101 Zelleriinae (= Yponomeutinae), 53 Zenobiinae (= Amphipyrinae), 125 Zenodoxinae (= Tinthiinae), 66 Zenodoxini (= Tinthiini), 66 Zephyrinae (= Theclinae), 98 Zerenidae (= Abraxini), 93 Zerenini (= Abraxini), 93 Zerynthiinae (= Zerynthiini), 96 Zerynthiini (Papilioniidae), 96 Zesiini (Lycaenidae), 98 ZEUGLOPTERA, 1 Zeuzeridae (= Cossidae), 75 Zeuzerinae (Cossidae), 75 Zizeerini (= Zizeerina), 98 Zizeerina (Lycaenidae), 98 Zizerini [sic] (= Zizeerina), 98 Zonosomatidi (= Cosymbiini), 93 Zonosomini (= Cosymbiini), 93 ZYGAENIDAE, 70 Zygaeninae (Zygaenidae), 70 ZYGAENOIDEA, 69

LEPIDOPTERA NEWS, 2003 (1-2): 23-24 (2005)

DISTRIBUTION NOTES ON TAIWAN MOTHS, 2. MIMICIA PSEUDOLIBATRIX (LEPIDOPTERA: PYRALIDAE: PYRALINAE)

J. B. HEPPNER 1

Florida State Collection of Arthropods, DPI, FDACS, P.O. Box 147100, Gainesville, Florida 32614-7100, USA

ABSTRACT.- The Chinese pyraline species, Minicia pseudolibatrix (Caradja), is newly recorded from Taiwan and named as a new subspecies, as M. p. taiwana n. subsp., with distribution records from the Taiwan Lepidoptera Survey. Notes also give specimen records from China and Japan.

KEYWORDS: Asia, Catocalinae, China, East Asia, Fujian, Guangdong, Höne Collection, Japan, Mimicia pseudolibatrix taiwana n.subsp., Noctuidae, Oriental, Pyralinae, Scoliopteryx, taxonomy.

This second report on Taiwan moths from the Taiwan Lepidoptera Survey (Heppner, 2003), begun in 1981, treats the large species, Mimicia pseudolibatrix (Caradja), subfamily Pyralinae, newly recorded herein for Taiwan. The species was originally described in the genus Hybolomia by Caradja (1925), based on a single male from the Höne Collection, collected at Lianping (formerly Lienping), Guangdong Province, southern China, from Höne's collecting in the Guangdong region during 1920-23. Höne was in China from 1917-46 and made collections over most areas of the country east of the Tibetan plateau (Niethammer, 1963). His vast collections of Lepidoptera from China, mostly moths, are now in museums in Germany (mostly in Bonn and Karlsruhe), other than what Caradja received from him and which is now preserved in the natural history museum in Bucharest, Romania.

Meyrick (1934) later named a new genus, Mellia, for this unusual pyralid species. Strand (1935) then coined a new name for the genus (Meltalia) because of some nomenclatural point, but his name is now listed as a synonym of the Meyrick name, with both names being junior to Mimicia Caradja (1925) (first named as a subgenus of Hybolomia) (Inoue, 1988; Sugi, 2000).

In subsequent notes about material from China, Caradja (1934) noted that among H. Höne (1883-1963) specimens collected in China, as well as in other collections by R. E. Mell (1878-1970) made in the years 1908-21, there were further specimens from Lianping, China, specifically at a nearby mountain pass called "Tsatmicku-gao", 18 km south of Lianping, at 900m elevation. The added specimens totaled 35 more collected at light from the same site in China, dated 11 Jul, 26 Sep, and 7 Oct (no year indicated), presumably collected by Mell. Caradja (1934) also mentioned two specimens from a location called "Kwanshien" (in Sichuan Prov., near Chengdu) in his German text, but this may be in error since this was in reference to the type "pair", but Caradja (1925) described the species from only a single male stated to be from Lianping. The variety "obscura" (Caradja, 1935) was also described from Guangdong Province but is merely a melanic form.

Since Caradja's (1925) original description of this interesting moth, few if any additional specimens have been taken in China, as far as is known (Luh and Kuan, 1953), and only recently has a survey in Fujian Province, at Wuyishan Nature Reserve, obtained the species again and extended its known range to another province of China (Wang et al., 2003). The Fujian data involves only a single female, collected 1 Sep 1981 (Wang et al., 2003).





Fig. 1. Paratype forewing of Mimicia pseudolibatrix taiwana 9, from Taiwan.

In 1988, the species was reported for the first time from Japan by Inoue (1988), for two specimens collected at Kiune, Niimi City, Okayama Prefecture: a male on 15 Sep 1986 and a female on 9 Aug 1987 (S. Yano coll.). In 2003, an additional female was found in Niigata Prefecture, Japan, on 25 Aug 2003 (Sakurai, 2003). Evidently, this species is very rare in Japan, since Japan is among the best surveyed of any nation for Lepidoptera yet only 3 specimens have been found there in the last 160 years, unless the species has somehow recently gotten to Japan and Taiwan.

In 1996 and again in 2005, the species was found in midelevation mountains in northern Taiwan, at about 1500m elevation in mid-July, in the Tamanshan range south of Taipei, near forest at the mountain resort town called Upper Palin, Taoyuan County (J. B. Heppner, collector). Both females were taken at blacklight in July. Another female was found in 1996 only about 20km northeast from there at the mountain area of Mingchih (1250m), while in mid-July 2005 another female was taken at Litonshan, Hsinchu Co. (1450m).

In the original description, Caradja (1925) noted the resemblance of the species to the noctuid, Scoliopteryx libatrix (Linnaeus), a European species of the subfamily Catocalinae that also occurs in East Asia, including Taiwan. The resemblance, of course, prompted Caradja (1925) to name the new pyralid after the noctuid. Although even a cursory examination clearly shows that M. pseudolibatrix belongs in Pyralidae, collections of unsorted noctuid moths should still be checked for any possible specimens of this pyralid inadvertantly left there due to this superficial resemblance to Scoliopteryx

The collection data from Japan and Taiwan, as well as China, indicate the species is active during several months from July to October, perhaps with two generations. However, nothing is otherwise known of its life cycle or what the larvae may feed on.

Given that only a single specimen has turned up in Taiwan in the last 25 years of intensive collecting in Taiwan with the Taiwan Lepidoptera Survey, besides the extensive collecting activities by the Japanese in Taiwan already since 1895 by both professional and amateur lepidopterists, *M. pseudolibatrix* would seem to be as rare in Taiwan as it is in Japan. However, at the right time and in the right place the species can be found in numbers, as was noted by Caradja (1934) for the 35 additional specimens taken at one locality in Guangdong, China, up to that time.

In Taiwan, the habitat of *M. pseudolibatrix* is typical Chinese subtropical montane forest, mostly hardwoods mixed with some tropical species, as well as cedars and other trees from higher elevations, and with numerous understory plants. The forests are, thus, similar to original forest in mountains of Guangdong and Fujian, where the moths have been found in mainland China.

The following new name is provided for the Taiwan population of *M. pseudolibatrix*:

Mimicia pseudolibatrix taiwana Heppner, n. subsp.

Adult similar in size (female 38mm wingspan) to mainland Chinese specimens, with similar wing maculation: forewing coloration reddish-brown (with orange highlights), with short white costal margin mark angled as in the nominate subspecies but more uniform in thickness (not narrowed at middle as in mainland China population); more basal central angular white mark also thicker than in nominate subspecies; postmedian curved white fascia with costal widening angled mesally (not angled slightly distal as in mainland China populations); hindwing uniformly gray but for slight brown tinge near margins. Males not known for Taiwan but in China they are much smaller (28mm) than the females.

Holotype female: TAIWAN.— Taoyuan Co.: Upper Palin (1500m), 11-18 Jul 1996, J.B.Heppner. Deposited with the National Taiwan Museum, Taipei, Taiwan (on indefinite loan to the Florida State Collection of Arthropods, Gainesville, Florida).

Paratypes (3 %): *Hsinchu Co.*: Litonshan (1450m), 13 Jul 2005 (1 %), J. B. Heppner. *Taoyuan Co.*:Upper Palin (1350m), 4-5 Jul 2005 (1 %). Mingchih For. Res. Area (1250m), 9-10 Jul 1996 (1 %), J. B. Heppner & H.-Y. Wang.

Etymology.- Named after the island of Taiwan.

Distribution.- China (Fujian and Guangdong), Japan, and Taiwan.

Notes.— The mainland China population of *M. pseudolibatrix* is darker brown than those of *M. p. taiwana* in Taiwan, as also in Japan. Adults have been figured by Wang *et al.* (2003) from Fujian, and by Inoue (1988) and Sakurai (2003) from Japan. The biology of the species remains to be studied more, but Yoshimatsu (1991) reported on the immature stages reared on artificial diet in Japan.

The Taiwan catalog update (Heppner, 2005) will list over 140 other new species records for Taiwan from the Taiwan Lepidoptera Survey, all added since publication of the original Taiwan catalog (Heppner and Inoue, 1992), but *M. pseudolibatrix* can already be added below as follows (with appropriate catalog number for insertion into the 1992 catalog):

MIMICIA Caradja, 1925

Mellia Meyrick, 1934 Meltalia Strand, 1935

394.1 **pseudolibatrix** (Caradja, 1925) (*Hybolomia*) China (Guangdong) var. *obscura* (Caradja, 1935) (*Mellia*) China (Guangdong) a) **taiwana** Heppner, **n. subsp.** Taiwan

ACKNOWLEDGMENTS

The Taiwan Lepidoptera Survey project began in 1980 with organizational meetings at the National Taiwan Museum (then called the Taiwan Provincial Museum), in Taipei, by J. F. G. Clarke (Smithso-

nian Institution, Washington, DC), and the first survey trip was by Dr. Don R. Davis, Smithsonian Institution, Washington, DC, in 1981. All subsequent trips were by the author, since 1982, and were primarily funded by two National Science Foundation grants (INT-8119539 and INT-8721716), plus survey grants from the Smithsonian Institution and added support from the Florida State Collection of Arthropods, Division of Plant Industry, Florida Dept. of Agriculture and Consumer Services, Gainesville, Florida. In Taiwan, project support has been primarily from the National Taiwan Museum, Taipei, and the Taiwan Forestry Research Institute, Taipei, and their entomology staffs, as well as a publication grant from the Pacific Cultural Foundation, Taipei, Special thanks to H.-Y. Wang, National Taiwan Museum, for facilitating all field trips in Taiwan since 1982. Principal housing of survey specimens is with the Florida State Collection of Arthropods, Gainesville, Florida, for the study of the collections, plus samples at the Smithsonian Institution and the National Taiwan Museum, Taipei.

REFERENCES

Caradja, A.

1925. Ueber Chinas Pyraliden, Tortriciden, Tineiden nebst kurze Betrachtungen, zu denen das Studium dieser Fauna veranlassung gibt. Mem. Sect. Stiint. Acad. Rom. (Bucharest), (3) 3:257-386, 2 pl.

1934. In A. Caradja and E. Meyrick, Materialien zu einer Microlepidopteren-Fauna Kwangtungs. Dtsch. Ent. Zeit. Iris (Dresden), 47:123-167.

Caradja, A., and E. Meyrick

1935. Materialien zu einer Microlepidopteren Fauna der chinesischen Provinzen Kiangsu, Chekiang und Hunan. Berlin: R. Friedländer & Son. 96pp, 3 pl.

Heppner, J. B.

 Distribution notes on Taiwan moths, 1. Cotachena and Uthinia (Lepidoptea: Pyralidae). Trop. Lepid. (Gainesville), 11:56. (2000)

2005. Part 1: Introduction and Faunal Synopsis. In J. B. Heppner and H. Inoue (eds.), Lepidoptera of Taiwan. Volume 1. Gainesville: Assoc. Trop. Lepid. 215+80pp (76 pl.).

Heppner, J. B., and H. Inoue (eds.)

 Lepidoptera of Taiwan. Volume 1. Part 2: Checklist. Gainesville: Assoc. Trop. Lepid. 276pp.

Inoue, H.

 An unrecorded Pyralinae from Japan. *Japan Heteroc. J.*, (Tokyo), 145:305-306.

Luh, C.-J., and C.-H. Kuan

1953. Recorded species of Pyralididae [sic] from China, a supplement to Wu's Catalogus Insectorum Sinensium, Part I. Subfamilies: Crambinae, Schoenobiinae, Anerastiinae and Phycitinae. Peking Nat. Hist. Bull., 20: 163-181 (1951-52).

Meyrick, E.

1934. In A. Caradja and E. Meyrick, Materialien zu einer Microlepidopteren-Fauna Kwangtungs. Dtsch. Ent. Zeit. Iris (Dresden), 47:123-167.

Niethammer, G.

1963. Zur Geschichte der "Sammlung Höne". Bonner Zool. Beitr. (Bonn), 14:234-244, 1 pl.

Sakurai, S.

 Records of some noteworthy moths from Niigata Prefecture. Yugato (Niigata), 174:117-119. [in Japanese]

Strand, E.

1935. Miscellanea nomenclatorica zoologica et palaeontologica. VII. Fol. Zool. Hydrobiol. (Riga), 7:300-306.

Sugi, S.

2000. Edition 2 'Post-MJ': Additions of Species and Changes in Names of Japanese Moths. Tokyo: Japan Heterocerist's Soc. 171pp.

Wang, J.-S., S.-M. Song, Y.-Y. Wu., and T.-M. Chen

2003. Fauna of Pyralidae of Wuyishan Nature Reserve in China. Beijing: China Sci. Tech. Pr. 328pp, 4 pl. [in Chinese]

Yoshimatsu, S.

1991. Immature stages of Mimicia pseudolibatrix (Caradja) (Pyralidae) reared on artificial diets. Japan Heteroc. J. (Tokyo), 162:194-196. [in Japanese]

LEPIDOPTERA NEWS, 2003 (1-2): 25-26 (2005)

DISTRIBUTION NOTES ON TAIWAN MOTHS, 3. PYRAUSTA MYSTICA

(LEPIDOPTERA: PYRALIDAE: PYRAUSTINAE)

J. B. HEPPNER 1

Florida State Collection of Arthropods, DPI, FDACS, P.O. Box 147100, Gainesville, Florida 32614-7100, USA

ABSTRACT.- Notes are provided on specimens from Taiwan of the Chinese (Sichuan) pyralid species Pyrausta mystica Caradja, including the first Taiwan records. The taxon Pyrausta mystica taitungensis n. subsp. is described as a new subspecies from Taiwan.

KEYWORDS: Asia, China, East Asia, Oriental, Pyrausta mystica taitungensis n.subsp., Pyraustinae, Sichuan, taxonomy.

Among many new moths recorded for Taiwan from the Taiwan Lepidoptera Project, this third note reports on a rare species of the pyralid subfamily Pyraustinae described from Sichuan, China, and now recorded for the first time from Taiwan. This is another of many to be added to the 1992 catalog (Heppner and Inoue, 1992), as will be noted in the additions to the catalog (Hepper, 2005).

Much of the Taiwan lepidopteran fauna is a disjunct remnant from the Himalaya Mountains of Nepal and western China, possibly isolated now in the higher mountains of Taiwan since the last Ice Age regression of habitats that once were continual from the Himalayas to Taiwan. Taiwan has the highest mountain area east of the Himalayas, other than for the isolated peak of Kinabalu in northern Borneo, so cooler habitats in Taiwan mountains have allowed these remnant lepidopterans to survive there. Likewise, many species from western Sichuan also have disjunct remnant populations in Taiwan. On the highest Taiwan mountains one also finds plants and Lepidoptera typical of Japan and even of northern Eurasian species.

The Himalayan and Sichuan lepidopteran faunas in Taiwan are sometimes at various elevations, depending on the species, and not necessarily all above 2000m elevation, where most of the Himalayan fauna begins in Taiwan in earnest. Such is the case with the small blackish pyralid moth found in Taiwan at only 720m elevation near the southeast region of Taiwan, in Taitung County. A similar moth was described from a single male from Sichuan, China, by A. Caradja (1932). He appropriately named it Pyrausta mystica Caradja, since he was unsure of its generic placement and had not seen anything similar from China or from the European fauna of his native Romania. The unique holotype of the species is in the natural history museum in Bucharest, Romania, along with the rest of the large Caradja Collection of Lepidoptera, much of it being Chinese moths.

The current Lepidoptera curator of the Muzeul National de Istoria Naturala 'Grigore Antipa', Bucharest, Romania, Dr. Mihai Stanescu, kindly helped reconfirm the label data and physical aspects of Caradja's type. The holotype male of P. mystica, type no. 177636 (Museum Bucharest, MGAB), has the following label data: Omei-shan (2300m), Sichuan, China, 20 Jul 1931, G. Franck. The rather worn male specimen has a wingspan of 19mm, and appears as illustrated in Fig. 1.

On earlier visits of mine to Romania and the Bucharest museum to study types there, in 1980 and again in 1984, I had already photographed the type specimen of P. mystica, along with many








Fig. 1-2. Adults of Pyrausta mystica: 1) holotype male, P. m. mystica, Omeishan, Sichuan, China (MGAB). 2) holotype female, P. m. taitungensis, Bishan Hot Springs, Taitung Co., Taiwan (FSCA).

others, but did not have time or need then to dissect it for genitalic examination. After sorting of the many collections obtained in Taiwan since 1981 from the Taiwan Lepidoptera Project, specimens of a small pyralid were found that could not be readily placed to any known species until the photograph of the Caradja type of P. mystica was checked to verify that the specimens unquestionably are the same species. Unfortunately, all the Taiwan specimens thus far are females, so no genitalic confirmation is as yet possible.

The Caradja collection has only the single male holotype of P. mystica and no other specimens of the species are known of in any other collection thus far, other than the 3 females from Taiwan. None of the more recent Chinese works on pyralid moths include the species and it was not even included in the last Chinese catalog

of Lepidoptera, by Wu (1938), although finally listed in the catalog update for Pyralidae by Luh and Kuan (1953). Thus, it is likely that no other specimens of the species have been collected in China.

The following new name is provided for the Taiwan population of P. mystica:

Pyrausta mystica taitungensis Heppner, n. subsp.

Adult similar in size (female 19-20mm wingspan) to mainland Chinese specimen, with similar wing maculation: body and forewing coloration brown-black, with 2 white (slight cream color) oval spots past mid-wing (more elongate than in nominate subspecies), angled from the end of cell equidistantly from termen border, the radial spot being somewhat larger than the cubital spot; also a much smaller white spot near the base; hindwing dark like forewing and white spots near midwing as in forwing except somewhat more elongated or oval than in forewing (males have these reduced and the inner spot elongated diffusely toward anal angle of wing margin); overall the wings and body have a bluish sheen in fresh specimens. Males not known for Taiwan but in China they are similar to females except for the hindwing spot pattern, where in the male the white marks are elongated instead of rounded as in the female.

Holotype female: TAIWAN.- Taitung Co.: Bishan Hot Springs (720m), 26-29 Oct 1984, J.B.Heppner & H.-Y. Wang. Deposited with the National Taiwan Museum, Taipei, Taiwan (on indefinite loan to the Florida State Collection of Arthropods, Gainesville, Florida).

Paratypes (2 9): same data as holotype (FSCA).

Etymology.- Named after the county of Taitung of the type locality.

Distribution.- China (Sichuan) and Taiwan.

Notes. – The Sichuan, China, population of P. mystica has forewing white spots somewhat more oval than those from Taiwan, but how much this is sexual variation is not known yet. The females from Taiwan show the "white" spots somewhat light-cream in coloration (not evident in the photos where the photo-flash has made the spots brighter white than on the actual specimen. The placement in the genus Pyrausta is still uncertain for P. mystica until examination of the male genitalia may clarify this. The biology of P. mystica remains unknown.

ACKNOWLEDGMENTS

The Taiwan Lepidoptera Survey project began in 1980 with organizational meetings at the National Taiwan Museum (then called the Taiwan Provincial Museum), in Taipei, by J. F. G. Clarke (Smithsonian Institution, Washington, DC), and the first survey trip was by Dr. Don R. Davis, Smithsonian Institution, Washington, DC, in 1981. All subsequent trips were by the author, since 1982, and were primarily funded by two National Science Foundation grants (INT-8119539 and INT-8721716), plus survey grants from the Smithsonian Institution and added support from the Florida State Collection of Arthropods, Division of Plant Industry, Florida Dept. of Agriculture and Consumer Services, Gainesville, Florida. In Taiwan, project support has been primarily from the National Taiwan Museum, Taipei, and the Taiwan Forestry Research Institute, Taipei, and their entomology staffs, as well as a publication grant from the Pacific Cultural Foundation, Taipei. Special thanks to H.-Y. Wang, National Taiwan Museum, for facilitating all field trips in Taiwan since 1982. Principal housing of survey specimens is with the Florida State Collection of Arthropods, Gainesville, Florida, for the study of the collections, plus samples at the Smithsonian Institution and the National Taiwan Museum, Taipei.

My thanks also to Dr. Mihai Stanescu, Muzeul National de Istoria Naturala 'Grigore Antipa', Bucharest, Romania, for his kind help in checking the Caradja holotype for me. Likewise, the former curator, the late Dr. Aurelian Popescu-Gorj, was very helpful during

my visits to Romania in 1980 and 1984, both at the museum in Bucharest and on field trips around the country. In Romania, my travel and research was supported in 1980 by the National Geographic Society, Washington, DC (grant 2136), the American Philosophical Society, Philadelphia, PA, and the Smithsonian Institution, Washington, DC. In 1984, a grant for study in Romania was received from the National Academy of Sciences, Washington, DC (scientific exchange program, Romania).

REFERENCES

Caradja, A.

Ueber Chinas Pyraliden, Tortriciden, Tineiden nebst kurze Betrachtungen, zu denen das Studium dieser Fauna veranlassung gibt. Mem. Sect. Stiint. Acad. Rom. (Bucharest), (3) 3:257-386, 2 pl.

Heppner, J. B.

2005. Part 1: Introduction and Faunal Synopsis. In J. B. Heppner and H. Inoue (eds.), Lepidoptera of Taiwan. Volume 1. Gainesville: Assoc. Trop. Lepid. 215+80pp (76 pl.).

Heppner, J. B., and H. Inoue (eds.)

Lepidoptera of Taiwan. Volume 1. Part 2: Checklist. Gainesville: Assoc. Trop. Lepid. 276pp.

Luh, C.-J., and C.-H. Kuan

Recorded species of Pyralididae from China, a supplement to Wu's Catalogus Insectorum Sinensium. Acta Ent. Sinica (Beijing), 3:91-

Wu. C.-F.

Catalogus Insectorum Sinensium. Volume IV [Lepidoptera]. Beijing: Fan Mem. Inst. Biol. 1009pp.

LEPIDOPTERA NEWS, 2003 (1-2): 27-28 (2005)

DISTRIBUTION NOTES ON TAIWAN MOTHS, 4. GENERA LIXA AND TAMRACA (LEPIDOPTERA: PYRALIDAE: PYRALINAE)

J. B. HEPPNER ¹

Florida State Collection of Arthropods, DPI, FDACS, P.O. Box 147100, Gainesville, Florida 32614-7100, USA

ABSTRACT.- Notes are provided on specimens from Taiwan in the Asian pyralid genera Lixa Walker and Tamraca Moore, including new Taiwan records, and description of the subspecies, Lixa productalis taiwana n. subsp. and Tamraca torridalis taiwana n. subsp. from Taiwan.

KEYWORDS: Asia, Borneo, China, East Asia, Japan, Lixa productalis taiwana n. subsp., Malaysia, Oriental, Pyralinae, Sarawak, Sulawesi, Tamraca torridalis taiwana n.subsp., taxonomy.

Among many new moths recorded for Taiwan from the Taiwan Lepidoptera Project, this note reports on some species in the pyralid genera Lixa Walker and Tamraca Moore, both in the subfamily Pyralinae. While Lixa has 6 known species in Asia, Tamraca includes only two species. The new Taiwan records below are added to the Taiwan catalog of 1992 (Heppner and Inoue, 1992), and are further listed in the catalog updates (Heppner, 2005).

All specimens reported on below are from the Taiwan Lepidoptera Survey, currently housed at the Florida State Collection of Arthropods (FSCA), Gainesville, Florida.

LIXA Walker

The genus Lixa Walker ([1866]) presently contains 6 known species, all from Southeast Asia (Robinson et al., 1994). The type species of the genus, Lixa productalis Walker, originally described from Sarawak, Borneo (Walker, [1866]), has until now remained known only from Borneo (thus far Sarawak and Brunei). Specimens from the Taiwan Lepidoptera Survey include 34 specimens of this species, which can now be reported for Taiwan for the first time, named herein as a new subspecies for Taiwan.

Lixa productalis taiwana Heppner, n. subsp.

Adults similar in size (male 20-25mm wingspan; female 27-28mm) to Bornean specimens, with similar wing maculation: body and forewing coloration reddish orange-brown, with HW more orange; FW with 1 fuscous line at 1/3 from base and a subterminal line angled near tornus, and costa with tan edge; HW with 2 curved fuscous lines (Taiwan specimens with HW more orange than from Borneo). Female similar to male.

Holotype male: TAIWAN.- Taoyuan Co.: Upper Palin (1500m), 11-18 Jul 1996, J.B.Heppner. Deposited with the National Taiwan Museum, Taipei, Taiwan (on indefinite loan to the Florida State Collection of Arthropods, Gainesville, Florida).

Paratypes (32 &, 3 9). : TAIWAN.- Chiayi Co.: Fennchihwu (1450m), 2-4 Jul 1985, J. B. Heppner & H.-Y. Wang (1 &). Hsinchu Co.: Litonshan (1450m), 13 Jul 2005, J. B. Heppner (2 &). Ilan Co.: Fushan Bot. Sta. (650m), 20-24 Jul 1996, J. B. Heppner (1 &, 1 9). Tuchan [Chilan For. Sta.] (480m), 1-2 Jul 1982, J. B. Heppner (1 &). Nantou Co.: Lo Farm, 8 km E. Puli (750m), 10-11 Jul 2005, J. B. Heppner (2 &). Taichung Co.: Kukuan (700m), 6 Jul 1985, J. B. Heppner & H.-Y. Wang (1 &). Taipei Co.: Wulai (200m), 17-19 Jun 1985, J. B. Heppne & H.-Y. Wang (4 &). Taoyuan Co.: Upper Palin (1350m), 7-9 Jul 1986, J. B. Heppner & H.-Y. Wang (1 &). Upper Palin (1500m), 11-18 Jul 1996, J. B. Heppner (19 &, 2 9). All FSCA. Etymology.- Named after the island of Taiwan.




Fig. 1-2. Adults: 1) Lixa productalis taiwana, holotype male. 2) Tamraca torridalis taiwana, holotype male.

Distribution.-Borneo (Sarawak, East Malaysia) and Taiwan. Notes.- The Taiwan population has the hindwings much more orange-brown than those from Borneo, which are more red-brown. The biology of the species remains uknown.

TAMRACA Moore

The genus Tamraca Moore (1887) was described for an unusual pyralid already described earlier from Sri Lanka, Asopia torridalis Lederer (1863). Walker (1965) described the same species again from northern India as Varnia incerta Walker. Since then the species, as T. torridalis, has been known from the Indian region and other parts of Southeast Asia (east at least to Sulawesi), as well as southern China (Hampson, 1896; Wang et al., 2003). Later, it was

^{1.} Contribution No. 997, Entomology Section, Bur. Ent. Nema. Plant Path., Div. Plant Industry, Florida Dept. Agric. & Consumer Serv., Gainesville, Florida.

reported from Japan (Inoue et al., 1982) and Taiwan (Heppner and Inoue, 1992).

The following subspecies is described for the Taiwan populations:

Tamraca torridalis taiwana Heppner, n. subsp.

Adults similar (male 20-22mm wingspan; female 20-34), with similar wing maculation to those from Southeats Asia: body and forewing coloration mostly fuscous on FW costa and radius, plus similar small spot at mid-wing (near costa) and on subterminal line, with remainder of wing red-brown intermixed with yellow, plus some fuscous along anal margin. HW mostly fuscous, with central light tan area surrounded with yellow and red-brown between 2 curved fuscous lines angled to anal margin and base.

Holotype male: TAIWAN.- Nantou Co.: Lienhuachih For. Sta. (700m), nr. Puli, 7-12 Sep 1983, J.B.Heppner, Deposited with the National Taiwan Museum, Taipei, Taiwan (on indefinite loan to the Florida State Collection of Arthropods, Gainesville, Florida).

Paratypes (25 &, 16 9): TAIWAN.- Ilan Co.: 5 km. N. Nanao (100m), 11-13 Oct 1984, J. B. Heppner & H.-Y. Wang (3 8). Tuchan [Chilan For. Sta.] (480m), 1-2 Jul 1982, J. B. Heppner (1 &, 1 9). Hualien Co.: Kuyuan (750m), Taroko Gorge, 6-7 Jul 2005, J. B. Heppner (1 9). Nantou Co.: Lienhuachih For. Sta. (700m), 7-12 Sep 1983, J. B. Heppner (218, 149). All FSCA.

Etymology.- Named after the island of Taiwan.

Distribution. - Sri Lanka and India/Nepal to Malayasia and Indonesia (Sulawesi), and China to Japan for the nominate subspecies; Taiwan for the Taiwan subspecies.

Notes.- Although the species was collected in Taiwan in numbers in 1983 (two were taken earlier in 1982), the Taiwan Lepidoptera Survey since then did not find many more, indicating that overall it may be fairly rare. It is unclear at this time, but the population in Japan can likely be named as a Japanese subspecies: it is much more orange-brown than the Taiwan populations. The biology of the species remains unknown.

ACKNOWLEDGMENTS

The Taiwan Lepidoptera Survey project began in 1980 with organizational meetings at the National Taiwan Museum (then called the Taiwan Provincial Museum), in Taipei, by J. F. G. Clarke (Smithsonian Institution, Washington, DC), and the first survey trip was by Dr. Don R. Davis, Smithsonian Institution, Washington, DC, in 1981. All subsequent trips were by the author, since 1982, and were primarily funded by two National Science Foundation grants (INT-8119539 and INT-8721716), plus survey grants from the Smithsonian Institution and added support from the Florida State Collection of Arthropods, Division of Plant Industry, Florida Dept. of Agriculture and Consumer Services, Gainesville, Florida. In Taiwan, project support has been primarily from the National Taiwan Museum, Taipei, and the Taiwan Forestry Research Institute, Taipei, and their entomology staffs, as well as a publication grant from the Pacific Cultural Foundation, Taipei. Special thanks to H.-Y. Wang, National Taiwan Museum, for facilitating all field trips in Taiwan since 1982. Principal housing of survey specimens is with the Florida State Collection of Arthropods, Gainesville, Florida, for the study of the collections, plus samples at the Smithsonian Institution and the National Taiwan Museum, Taipei.

REFERENCES

Butler, A. G.

1877-89. Illustrations of Typical Specimens of Lepidoptera Heterocera in the Collection of the British Museum. London: Br. Mus. 7 v. (Vol. 3, 1878).

Hampson G F

[1893]-96. The Fauna of British India, including Ceylon and Burma, Moths. London: Taylor & Francis. 4 v.

Haruta, T. (ed.)

1992-2000. Moths of Nepal. Tokyo: Japan Heteroc. Soc. 6 v. In Tinea (Tokyo), 13(Suppl. 2):1-122, 32 pl. (1992); 13(Suppl. 3):1-160, pl. 33-64 (1993); 14(Suppl. 1):1-171, pl. 65-96 (1994); 14 (Suppl. 2):1-206, pl. 97-128 (1995); 15(Suppl. 1):1-314, pl. 129-160 (1998); 16(Suppl. 1):1-163, pl. 161-174 (2000).

Heppner, J. B.

2005. Part 1: Introduction and Faunal Synopsis. In J. B. Heppner and H. Inoue (eds.), Lepidoptera of Taiwan. Volume 1. Gainesville: Assoc. Trop. Lepid. 215+80pp. (76 pl.).

Heppner, J. B., and H. Inoue (eds.)

1992. Lepidoptera of Taiwan. Volume 1. Part 2: Checklist. Gainesville: Assoc. Trop. Lepid. 276pp.

Inoue, H., S. Sugi, H. Kuroko, S. Moriuti, and A. Kawabe

Moths of Japan. Tokyo: Kodansha. 2 v. (392 pl.). [in Japanese]

Lederer, J.

1863. Beitrag zur Kenntnis der Pyralidinen. Wiener Ent. Monatsch. (Vienna), 7:243-280, 331-502, pl. 2-18.

Moore, F.

1880-[87]. The Lepidoptera of Ceylon. London: Reeves. 3 v., 215 pl.

Robinson, G. S., K. R. Tuck, and M. Shaffer

A Field Guide to the Smaller Moths of South-East Asia. Kuala Lumpur: Malaysian Nature Soc. 309pp (32 pl.).

Walker, F.

1854-66. List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. London: Brit. Mus. 35 pts. (Pat. 17, 1859; Pt. 34,

Wang, J.-S., S.-M. Song, Y.-Y. Wu., and T.-M. Chen

Fauna of Pyralidae of Wuyishan Nature Reserve in China. Beijing: China Sci. Tech. Pr. 328pp, 4 pl. [in Chinese]