

VAGINAL TUMORS

PATHOPHYSIOLOGY

VAGINAL, VESTIBULAR, VULVAR NEOPLASIA

- The majority of vaginal, vestibular & vulvar neoplasms are benign, w/ leiomyoma being the most common. Around ¼ of tumors are malignant, w/ leiomyosarcoma the most frequently diagnosed. Other reported malignancies include transmissible general tumor (TVT), transitional cell tumors & neuroendocrine tumors.
- The average age at diagnosis is around 11 years & there are no reported breed predispositions.
- Malignant neoplasia can metastasize to the local lymph nodes, spleen, lungs or cervical spinal cord.
- Clinical signs include bulging of perineum, vaginal prolapse, tenesmus, dysuria, urinary incontinence, difficulty copulation & sanguinous or purulent discharge.
- Diagnosis is made by vaginoscopy, digital vaginal exam & rectal palpation in small dogs.
- Cytology should be performed & thoracic radiography & abdominal radiography & ultrasound are recommended for full staging.
- Treatment of aggressive (ie malignant) tumors consists of surgical excision (vaginectomy) via an episiotomy or a combined (abdominal & perineal) approach. With the appropriate surgical expertise, splitting of the pelvic is not always needed, though this is a technically challenging surgery & referral to a specialist center is recommended. TVT is best treated with vincristine, though radiotherapy is reported to be successful.
- Surgical excision carries a high rate of reoccurrence & therefore is considered a poor alternative. Treatment of benign tumors is performed by marginal excision via an episiotomy.
- Prognosis is good for both TVT & a completely excised leiomyoma & guarded for malignant tumors. Adjunctive treatment is probably necessary after resection of malignant tumors.

+ GENERAL CONSIDERATIONS

- Vaginal & vulval tumors are the 2nd most common canine reproductive tumor & account for 2.4%-3.0% of all canine tumors •

Majority of vaginal & vulval tumors are benign:

- Leiomyoma & fibroma in cat
- Leiomyoma & lipoma in dog

+ LEIOMYOMA

- 86% of vaginal & vulvar tumors are benign smooth muscle tumors (i.e., leiomyoma & fibroma)
- Majority of leiomyomas arise from vestibule of vulva rather than vagina
- Extraluminal & intraluminal forms have been described:
- Extraluminal present with a slow-growing perineal mass
- Intraluminal are attached to vestibular or vaginal wall by variably sized pedicle & can be multiple
- Mucosa is generally intact, but ulceration may occur with exposure & irritation
 - Leiomyoma may be hormone dependent:
- Leiomyoma has not been diagnosed in dogs ovariectomized < 2 years
- 15% local tumor recurrence rate in intact & 0% in dogs following ovariectomy
- Mean age 10.8 years
- Breed predisposition: Boxer
- Incidence of leiomyoma higher in nulliparous bitches
- 33% associated with cystic endometrial hyperplasia, ovarian cysts, & mammary gland tumors

+ LIPOMA

- Lipoma can arise from perivascular or perivaginal adipose tissue & attach to tuber ischii
- Lipoma can lie within pelvic canal & are usually well-circumscribed & relatively avascular
- Mean age 6.3 years

+ TRANSMISSIBLE VENEREAL TUMOR (TVT)

- TVT occurs in ~4-year-old dogs with increased risk in free roaming dogs
- Cell origin of transmissible venereal tumors is unknown, but an undifferentiated round cell tumor of reticuloendothelial origin is most likely
- Transmissible venereal tumors have a common origin as chromosomal aberrations are constant & highly specific
- Virus particles have been identified in transmissible venereal tumors
- However, viral etiology is unlikely as the disease cannot be transmitted by cellfree filtrates

- Transmissible venereal tumors are transmitted by contact with genital mucous membranes during coitus
- Transmissible venereal tumors appear as multiple tumors along the mucosal lining of the vagina & vestibule

+ OTHER

- Benign tumors: sebaceous adenoma, fibrous histiocytoma, benign melanoma, myxoma & myxofibroma
- Malignant tumors: leiomyosarcoma, ADC, SCC, TCC, HSA, OSA, MCT, & epidermoid carcinoma
- Carcinoma of bladder or urethra may present with palpably enlarged urethral papilla

CLINICAL FEATURES

+ CLINICAL SIGNS

- Duration of clinical signs longer for extraluminal compared to intraluminal leiomyoma
- Intraluminal leiomyoma often presents as mass extruding between vulval lips, particularly during estrous
- Other clinical signs include vulval bleeding or discharge, enlarging vulvar mass, dysuria, hematuria, tenesmus, excessive vulval licking, & dystocia
- Lipomas usually present with a slowly growing mass impinging on adjacent structures

+ DIAGNOSIS

- Vaginoscopic examination, retrograde vaginography, & urethrocystography may delineate mass
- Ultrasonography, FNA, & histopathology

TREATMENT

+ MEDICAL MANAGEMENT

- Local tumor recurrence rate is high for dogs with transmissible venereal tumor & surgery is not recommended
- Transmissible venereal tumors are very responsive to chemotherapy & radiation therapy
- Chemotherapy: vincristine 0.5-0.7 mg/m² IV 4-8 times ± doxorubicin

+ SURGICAL MANAGEMENT – BENIGN TUMORS

- Exploratory celiotomy for ovariohysterectomy (due to hormonal dependence & local tumor recurrence)
- Conservative surgical resection
- Wide resection probably not required if ovariohysterectomy performed concurrently
- Dorsal episiotomy may be required to provide adequate visualization & ensure complete resection
- Dorsal episiotomy indicated for extraluminal vaginal & vulvar tumors as tumors are usually well-circumscribed & poorly vascularized resulting in good probability of complete excision
- Perineal approach or pubic split is rarely required

+ SURGICAL MANAGEMENT – MALIGNANT TUMORS

- Malignant infiltrative vaginal tumors treated with complete vulvovaginectomy & perineal urethrostomy
- Sternal recumbency in perineal stand with perineum elevated
- Urethra catheterized
- Fusiform skin incision performed around vulva
- Deeper tissues sharply dissected from labia & vestibule
- Constrictor vestibuli & constrictor vulvae muscles are dissected from the vestibule
- Dorsal labial branches of the ventral perineal artery are ligated or bleeding controlled with electrocautery
- Catheterized urethra identified & dissected free from encircling constrictor vestibuli muscles

- Vagina dissected with transection of ischiocavernosus & ischiourethralis muscles
- Dissection continued cranially between paired levator ani muscles to level of cervix
- Vaginal branches of vaginal & uterine arteries & veins ligated
- Vagina transected immediately caudal to cervix in intact bitches or cervix & uterine stump removed in spayed dogs
- Deep tissues closed to reduce dead space
- Perineal urethrostomy performed with transected urethra tractioned caudally, distal end spatulated, & closed in 2 layers with final layer mucosa to skin

+ PROGNOSIS

- Complete surgical excision is usually curative
- Guarded to poor prognosis with ADC, TCC, & SCC due to high local tumor recurrence & metastatic rates




FIG. 26.16 (A) For episiotomy, place noncrushing forceps on each side of the proposed incision site. Make a full-thickness incision from the dorsal vulvar commissure to near the external anal sphincter muscle. Explore the vagina and vestibule. (B) Appose vaginal mucosa with a simple continuous suture pattern. (C) Appose muscle, subcutis, and connective tissue with a second layer of sutures and skin with a third layer of appositional sutures.

Surgery of the reproductive tract 1




Fig. 5.12a,b Episiotomy. 1–4. The incision is made at the level of the dorsal vulvar commissure (between non-crushing forceps) a suturing of the incision is performed in three layers (mucosa, muscle and connective tissues, skin). b. Episiotomy performed to access vaginal leiomyoma. A urinary catheter is placed. *Illustrator: Elaine Leggett*




Figure 3 - Vaginal procedure: Positioning, episiotomy and placement of a urinary catheter

A: Positioning of the dog for episiotomy.

B: Midline episiotomy incision with urinary papilla visualisation.

C: Urinary catheter insertion.

1) Vulva 2) Anus purse string 3) Urinary papilla 4) Urinary catheter




Figure 4 - Vaginectomy with tumor removal

A: Placement of stay sutures on the ventral vaginal wall.

B: Inverting the cranial aspect of the vagina.

C: Externalised vaginal tumor by inverting the cranial aspect of the vagina through the episiotomy.

D: Tumor after excision (low grade leiomyosarcoma).

E: Cross-sectional view of the tumor (low grade leiomyosarcoma).

F: Ventral vaginal wall after tumor excision.

1) Stay suture 2) Urinary catheter 3) Urinary papilla 4) Ventral vaginal wall 5) Dorsal vaginal wall

TREAT Veterinary Surgery Service

Dr. Le-Nguyen, DVM, SAS (GPCert Canine/Feline Surgery)

(Practice Limited to Surgery)

(916) 230-8103

TREATVeterinarySurgeryService@gmail.com

TREATVeterinarySurgeryService.com