

Preventative/Wellness Care for Dogs & Puppies



Spay Neuter RoCo
Veterinary Clinic, PLLC

Spay/Neuter • Dentistry • Wellness

www.SpayRoCo.com

Distemper/Adenovirus/Parainfluenza/Parvovirus Vaccine (CORE Vaccine)

- Puppies ≤16 Weeks of Age (Boosters)
 - At least 3 doses between 6 & 16 weeks, 3-4 weeks apart
- Dogs >16 Weeks of Age
 - 2 doses, 3-4 weeks apart

Leptospirosis Vaccine (CORE Vaccine)

- Puppies ≤16 Weeks of Age (Boosters)
 - 2 doses starting at 8-9 weeks, 3-4 weeks apart
- Dogs >16 Weeks of Age
 - 2 doses, 3-4 weeks apart

Rabies Vaccine (CORE Vaccine)

- Initial vaccine at 12-16 weeks of age, booster in 1 year & then every 3 years

Bordetella Vaccine (Non-core, but recommended if has any exposure to other dogs)

- A single (intranasal) dose as a puppy & booster annually

Lyme Vaccine (Non-core, but recommend if exposed to ticks)

- Puppies ≤16 Weeks of Age (Boosters)
 - 2 doses starting at 8-9 weeks, 3-4 weeks apart
- Dogs >16 Weeks of Age
 - 2 doses, 3-4 weeks apart

Heartworm Test for dogs 6 months & older who have not been on consistent monthly heartworm prevention since 6-8 weeks of age

Heartworm prevention which includes a monthly intestinal parasite dewormer is strongly recommended each month, year-round on all dogs & puppies 6 weeks & older.

Millbeguard is available for purchase

Flea & tick control is strongly recommended each month, year-round on all dogs & puppies

Credelio is available for purchase

DA2PP Vaccine-Distemper/Adenovirus/Parainfluenza/Parvovirus Vaccine

Canine Distemper: A severe and often fatal viral illness that attacks the nervous, respiratory, and gastrointestinal systems.

Adenovirus (Types 1 and 2): Causes infectious canine hepatitis (Type 1) and contributes to kennel cough (Type 2).

Parainfluenza: A virus that contributes to kennel cough.

Parvovirus: A highly contagious and potentially fatal virus, particularly dangerous for puppies.

Leptospirosis Vaccine

Leptospirosis is a bacterial infection that can affect dogs and humans (Zoonotic**). It is spread through contact with infected water or soil that has been contaminated by the urine of infected animals, including rodents, dogs, cattle, and wildlife.

Rabies Vaccine

Rabies is an incurable virus that attacks the brain and spinal cord. All mammals, including dogs and humans, can be infected with rabies.

Boretella Vaccine (Kennel Cough)

Kennel Cough is a highly contagious respiratory infection in dogs. Our vaccine is an intranasal vaccine for dogs that protects against canine adenovirus type 2, canine parainfluenza virus, and Bordetella bronchiseptica.

Lyme Vaccine

Lyme disease is a bacterial infection transmitted by blacklegged ticks that can affect dogs. Vaccination is an effective way to protect dogs from Lyme disease