

LONG-TERM TRAINING ADAPTATIONS IN ELITE MALE VOLLEYBALL PLAYERS

JEREMY M. SHEPPARD^{1,2} AND ROBERT U. NEWTON¹

¹School of Exercise, Biomedical, and Health Sciences, Edith Cowan University, Joondalup, Western Australia; ²Australian Volleyball Federation, Canberra, Australia

ABSTRACT

Sheppard, JM, and Newton, RU. Long-term training adaptations in elite male volleyball players. *J Strength Cond Res* 26(8): 2180–2184, 2012—Several investigations have demonstrated differences in anthropometry, jump performance, and strength variables between developmental and elite-level volleyball players. However, within the elite level of play, the magnitude of change that can occur with training is unclear. The purpose of this investigation was to examine the anthropometric, vertical jump, and strength quality changes over 2 years in a group of national team volleyball players. Fourteen national team volleyball players (age, 23.0 ± 4.1 years; height, 1.98 ± 0.07 m; weight, 91.7 ± 7.9 kg) began and completed this study. Participants had all played international matches (representing Australia) before the examination time period and continued to do so during the international season. Anthropometry (stature, mass, and sum of 7 skinfolds), vertical jump measures (counter-movement vertical jump; depth jump from 0.35 m, DJ; spike jump, SPJ, all including arm swing), and lower-body power (jump squat at body mass, and jump squat + 50% body weight, JS50) measures were tested before and at the conclusion of the investigation period. Significant ($p < 0.05$) improvements were observed in sum of 7 skinfolds, DJ, SPJ, and JS50 performance, with large magnitude changes ($d > 0.70$) in the sum of 7 skinfolds reduction, SPJ, and leg extensor power. This study has demonstrated that elite male volleyball players can improve leanness and power, which contribute to improvements in vertical jump.

KEY WORDS jump, spike jump, elite, training

INTRODUCTION

Volleyball is an intermittent and explosive sport, involving brief but frequent movements such as jumping and diving (3,8). Jumping activities can include both horizontal approach movements (spike jumps, SPJ) and movements without a horizontal approach (jump setting, jousts, and blocking). Considering the importance of these jumping activities to the performance outcome in volleyball, and the frequency that they occur in a typical match (7), both countermovement vertical jump (CMVJ) ability and SPJ ability are considered important qualities contributing to game performance by elite volleyball players, and they are the primary targets that training programs aim to enhance (4,6,12).

Recent studies have examined the difference between junior and senior elite volleyball players (7), highlighting the importance of vertical jump improvement for developing athletes to progress successfully from lower to higher levels. Cross-sectional examinations of junior and senior groups provides strong evidence for the need to improve both CMVJ and SPJ, and presenting useful performance data for setting standards and benchmarks for athletes aspiring to play at the international level (6–8,11). Despite the acknowledged significance, little is known as to the trainability of CMVJ and SPJ and other potentially related qualities within senior elite populations of volleyball players. Of great interest to the sport scientist and strength and conditioning coach working with elite volleyball athletes is the magnitude of trainability of these qualities and which specific variables require development to underpin these improvements in vertical jump ability. As studies with elite populations of athletes is rare, the magnitude of improvement possible with elite athletes is poorly understood, as is the interplay of these trainable aspects. The purpose of this investigation was to examine potential changes in anthropometry, vertical jump ability, and fundamental neuromuscular capacities over a 2-year time period in elite male volleyball players to gain insight into the trainability of these qualities within an elite population.

METHODS

Experimental Approach to the Problem

To assess the changes that occurred in jump performance, speed-strength, and anthropometric variables, a longitudinal

This study was completed without the assistance of any grant or external financial support.

Address correspondence to Jeremy M. Sheppard, physicalpreparation@yahoo.com.

26(8)/2180–2184

Journal of Strength and Conditioning Research
© 2012 National Strength and Conditioning Association

TABLE 1. Changes (mean \pm SD) in anthropometric variables over 2 years in elite male volleyball players ($n = 14$).

	Pre-Test	Post-Test	ES (d)	Magnitude	Alpha (p)
Height (cm)	198.1 \pm 7.0	187.8 \pm 7.1	0.04	No change	0.415
Mass (kg)	91.7 \pm 7.9	92.7 \pm 6.9	0.14	No change	0.225
Skinfolds (mm)	56.7 \pm 9.0	50.8 \pm 8.5	0.7	Large	<0.001
Lean mass ratio	1.7 \pm 0.3	1.9 \pm 0.3	0.65	Moderate	<0.001

Alpha <0.05 is considered significant.

analysis was performed over a 2-year time period with a cohort of elite national team volleyball players.

Subjects

Fourteen national team volleyball players (age, 23.0 \pm 4.1 years; height, 1.98 \pm 0.07 m; weight, 91.7 \pm 7.9 kg) began and completed this study. Participants had all played international matches (representing Australia) before the examination time period and continued to do so during the international season. During professional seasons, match play volume varied depending on club, with training time typically involving 4–7 court and 2 strength sessions per week. During international seasons, participants typically engaged in 7–9 court and 2–4 strength sessions per week, with match involvement varying by player from 15–40 per season.

Strength training sessions varied throughout the 2-year time period, based on individual and to a large extent their professional club commitments. However, the main components of the sessions typically involved 1–2 Olympic lifts and their variations (e.g., Snatch, power clean) (3–10 sets total), 1–3 maximal strength training exercises (squat, front squat, pull-up) (1–6 repetitions, 4–10 sets total), 2–4 supplementary exercises (unilateral leg strength, rows, etc.), followed by any

prescribed medical exercise. Total sets for workouts ranged greatly depending on the athlete and the time of year, from 8 to 22 main “working” sets.

All participants received a clear explanation of the study, including the risks and benefits of participation, and if following this explanation their decision was to not be included in the analysis, it did not adversely affect any current or future team selection. All included participants provided written informed consent for testing and data analysis. Approval for this investigation was granted from the Institutional Human Ethics Committee.

Procedures

Participants performed a maximum-effort CMVJ, depth jump from a 0.35-m box (DJ), and an SPJ (with approach) using a vane jump and reach apparatus, which allowed for recording of the maximum height reached to the nearest centimeter (Yardstick; Swift Systems, Lismore, Australia). In the CMVJ, no horizontal approach was allowed, whereas in the SPJ, an approach ranging from 3 to 4 steps was used, based on the athlete’s preference. For the DJ, participants stepped off a 0.35-m box, and immediately upon landing, attempted to jump as high as possible. The population-specific intraclass correlation coefficients (ICC) (%Typical Error in parenthesis) of the height of the CMVJ, DJ, and SPJ was 0.98 (2.5%), 0.97 (3.0%), and 0.97 (3.2%) respectively.

Kinetic and kinematic assessments of jump squat performance were conducted with the subjects standing on a commercially available force plate (400 Series Performance Force Plate, Fitness Technology, Adelaide, Australia).




Figure 1. Mean (\pm SD) of 2-year changes in countermovement vertical jump, depth jump, and spike jump of elite male volleyball players ($n = 14$). All changes were statistically significant ($p < 0.01$).




Figure 2. Mean (\pm SD) of 2-year changes in unloaded (jump squat at body mass) and loaded jump squat (body weight + 50% body weight, JS50) performance of elite male volleyball players ($n = 14$). *Indicates change is statistically significant ($p < 0.001$).

TABLE 2. Changes (mean ± SD) in speed-strength variables over 2 years in elite male volleyball players (n = 14).

	Pretest	Posttest	ES (<i>d</i>)	Magnitude	Alpha (<i>p</i>)
Unloaded jump squat					
Peak power (W)	6,978 ± 1256	7,440 ± 890	0.51	Moderate	0.043
Relative peak power (W/kg)	76.2 ± 12.3	80.4 ± 9.1	0.47	No change	0.069
Force (N)	2,140 ± 243	2,228 ± 209	0.42	No change	0.104
Jump height (m)	0.502 ± 0.065	0.512 ± 0.066	0.15	No change	0.365
Loaded jump squat (+50% BM)					
Peak power (W)	5,793 ± 771	5,957 ± 742	0.22	No change	0.287
Relative peak power (W/kg)	63.2 ± 6.3	64.4 ± 7.7	0.16	No change	0.393
Force (N)	2,579 ± 283	2,640 ± 232	0.26	No change	0.246
Jump height (m)	0.340 ± 0.038	0.376 ± 0.052	0.7	Large	<0.001

Alpha <0.05 is considered significant.

A position transducer (PT5A, Fitness Technology, Adelaide, Australia) was connected to a 400-g wooden pole (body weight jumps, BWJS) or weightlifting bar and weightlifting lifting plates (body weight + 50% of body weight as additional mass, JS50) held across the shoulders. Both the force plate and position transducer sampled at 200 Hz and were interfaced with computer software (Ballistic Measurement System; Fitness Technology, Adelaide, Australia) that allowed direct measurement of force-time characteristics (force plate) and displacement-time and velocity-time (position transducer) variables.

Before all data collection procedures, the force plate was calibrated using a spectrum of known loads, and then assessed against 3 criterion masses. The position transducer was calibrated using a known distance of 1 m. The ICC and %TE of the displacement, force, and power measures used in the assessment methodology, with this population group, were 0.95–0.97 (3.1–3.9%), 0.95–0.97 (3.1–4.0%), and 0.80–0.98 (3.0–9.5%) respectively.

All subjects were assessed for height, mass, standing reach, and the sum of 7 skinfolds. The sum of 7 skinfolds was determined after measurement of the triceps, subscapular, biceps, supraspinale, abdominal, quadriceps, and calf skinfold using a Harpenden skinfold caliper (British Indicator, West Sussex, United Kingdom). A composite ratio of body mass divided by the sum of 7 skinfolds was then determined to reflect the amount of mass that is made up of lean tissue, termed the lean mass ratio (LMR). All tests were conducted by a single researcher certified by the International Society for the Advancement of Kinanthropometry. The ICC and %TE for height, mass, and standing reach were 0.99 (1.5%), 0.99 (1.2%), and 0.98 (2.0%) respectively. The test-retest ICC and %TE for the skinfold assessment was 0.99 (2.2%).

All testing was conducted under the same conditions, at the same time of day, for each testing type. Although having elite athletes completely recovered for the purpose of testing is unrealistic, to minimize fatigue, testing took place after

a complete rest day and was spread over 2 days. Anthropometry was completed in the morning of day 1 to attempt to control for hydration status, with the vertical jump testing occurring in the afternoon. Speed-strength assessment was completed the following day, in the afternoon.

Statistical Analyses

Changes in the mean were used to assess the practical differences that occurred across the investigation time period. Paired *t*-tests were used to assess changes in the anthropometric and vertical jump variables over the 24-month period, with alpha set at *p* ≤ 0.05. Additionally, Cohen’s effect sizes (*d*) were calculated to reflect the magnitude of any changes that may be observed, with the criteria of <0.40 small, 0.40–0.70 moderate, and 0.70–1.00 large.

RESULTS

Anthropometric changes across the 2-year investigation period are presented in Table 1. A large magnitude reduction in skinfolds (*d* = 0.70, *p* < 0.001) contributed to a moderate increase in the LMR (*d* = 0.65, *p* < 0.001).

Jump height for CMVJ, DJ, and SPJ (*d* = 0.52, 0.51, and 0.70 respectively, *p* < 0.001) increased significantly over the 2-year period (Figure 1). Loaded jump squat height (JS50) also increased significantly (Figure 2) with large magnitude increases (*d* = 0.70, *p* < 0.001). With the exclusion of a moderate (*d* = 0.51, *p* = 0.043) increase in peak power in the BWJS, no other changes were observed in the measured kinetic and kinematic descriptors of the jump squat tests.

DISCUSSION

Of great interest to the strength and conditioning coach working with elite programs is the specific variables that can be developed to advance performance. This information is invaluable in decision-making and priority setting for optimizing the success of elite athletes. The results of this investigation are novel in that they examine the changes in

CMVJ and SPJ, as they relate to changes in anthropometry and speed-strength over 2 years in elite volleyball players (Table 2). We have demonstrated that large changes in vertical jump ability and speed-strength and improvements in lean mass can be realized, even in such a highly trained population. This investigation provides the strength coach and sport scientist with the basis of a rationale to prioritize: (a) Reducing fat mass; (b) increasing stretch load tolerance (through depth jumping and other jumping methods) and (c) increasing loaded jump performance, in elite national team players.

Total skinfold thickness was reduced by a large magnitude ($d = 0.70$, $p < 0.001$) over the course of this investigation, supporting a moderate and significant ($d = 0.65$, $p < 0.001$) LMR increase from 1.7 to 1.9 (Table 1). In a comparison of athletes transitioning from junior to senior elite competition over a similar time period to this study, LMR was observed to increase from 1.5 to 1.8 (10). Considered together, these results emphasize not only the importance of increasing lean mass in junior elite players to accommodate the large increases in strength observed (10) but also that further reductions in fat mass and increases in LMR should be expected for elite volleyball players. It stands to reason that in a sport where jumping ability is of primary importance (4–7,9), training should be aimed at increased strength and maximal power while maintaining very low fat mass.

During the course of this investigation, the participants improved their CMVJ and SPJ by 0.045 and 0.090 m, respectively, which are considered moderate to large, and practical, change magnitudes ($d = 0.52$ and $d = 0.70$, $p < 0.001$). These improvements suggest that not only are increases in jumping ability required for developing volleyball players (10,11), but also that further improvements can and should be made within elite populations.

Our present results support previous findings that demonstrate the importance of depth jump ability in volleyball players and the likely relationship between developing depth jump ability and improving both CMVJ and SPJ (4,6,9). In the current investigation, DJ scores improved by 0.041 m ($d = 0.51$, $p < 0.001$). Considering these results, and others that demonstrate strong association between improving depth jumping ability and increasing CMVJ and SPJ (1,4,6,9), strength and conditioning coaches should consider improving stretch load tolerance in depth jumping and related activities, as an important training priority in elite volleyball players.

The large increases in loaded jump squat height ($d = 0.70$, $p = 0.001$) demonstrate the importance of fast force production in elite volleyball players. This quality is highly trainable in developing volleyball players (10), and the results of the present study support the trainability of this quality even in elite players. Furthermore, the results of this study highlight the potential association between increased power output under load and improved vertical jump ability in volleyball players. As recently highlighted by Cormie et al (2),

increased force, impulse, and power output during jumping is in part a result of more effective use of the eccentric phase. Although not assessed in the current study, improvement in DJ performance indirectly lends support to this suggestion.

Examinations of training-induced changes are rare in elite populations of athletes. The results of this study suggest that even in elite populations, several impactful gains can be made in anthropometry, stretch-shorten cycle function, and speed-strength, which may underpin vertical jump improvement in elite volleyball players.

PRACTICAL APPLICATIONS

Large changes in vertical jump ability, lower-body power, and improvements in lean mass can be realized in already well-trained volleyball players. These results provide the strength coach and sport scientist with the basis of a rationale to prioritize: 1) Reducing fat mass; 2) increasing stretch load tolerance (through depth jumping and other jumping methods); and 3) increasing loaded jump performance, in elite national team players.

Volleyball players should be expected to increase their CMVJ and SPJ, even at the most elite level. This may be in part accomplished through reducing fat mass, improving force and power output during jumping, and development of high levels of stretch load tolerance in stretch shorten cycle activity. Strength and conditioning coaches should prioritize training for each individual volleyball player based on a testing profile, to highlight areas of relative weakness, and in turn, to best exploit trainable qualities in elite players.

REFERENCES

1. Bobbert, MF. Drop jumping as a training method for jumping ability. *Sports Med* 9: 7–22, 1990.
2. Cormie, P, McGuigan, MR, and Newton, RU. Changes in the eccentric phase contribute to improved stretch-shorten cycle performance after training. *Med Sci Sports Exerc* 42: 1731–1744, 2010.
3. Polgaze, T, and Dawson, B. The physiological requirements of the positions in state league volleyball. *Sports Coach* 15: 32–37, 1992.
4. Sheppard, JM, Chapman, D, Gough, C, McGuigan, MR, and Newton, RU. Twelve month training induced changes in elite international volleyball players. *J Strength Cond Res* 23: 2096–2101, 2009.
5. Sheppard, JM, Cormack, S, Taylor, KL, McGuigan, MR, and Newton, RU. Assessing the force-velocity characteristics of well trained athletes: the incremental load power profile. *J Strength Cond Res* 22: 1320–1326, 2008.
6. Sheppard, JM, Cronin, J, Gabbett, TJ, McGuigan, MR, Extebarria, N, and Newton, RU. Relative importance of strength and power qualities to jump performance in elite male volleyball players. *J Strength Cond Res* 22: 758–765, 2007.
7. Sheppard, JM, Gabbett, T, and Stanganelli, LC. An analysis of playing positions in elite mens' volleyball: considerations for competition demands and physiological characteristics. *J Strength Cond Res* 23: 1858–1866, 2009.
8. Sheppard, JM, Gabbett, TJ, Taylor, KL, Dorman, J, Lebedew, AJ, and Borgeaud, R. Development of a repeated-effort test for elite men's volleyball. *Int J Sports Physiol Perform* 2: 292–304, 2007.

9. Sheppard, JM, McGuigan, MR, and Newton, RU. The effects of depth-jumping on vertical jump performance of elite volleyball players: an examination of the transfer of increased stretch-load tolerance to spike jump performance. *J Aust Strength Condition* 16: 3–10, 2008.
10. Sheppard, JM, Nolan, E, and Newton, RU. Two year changes in strength and power qualities in volleyball players successfully progressing from junior to senior national team. In *Australian Strength and Conditioning Association—International Conference on Applied Strength and Conditioning*. Gold Coast, Australia, 2010.
11. Stanganelli, LC, Dourado, AC, Oncken, P, Mancan, S, and Cesar da Costa, S. Adaptations on jump capacity of Brazilian volleyball players prior to the under 19 World championships. *J Strength Cond Res* 22: 741–749, 2008.
12. Viitasalo, JT. Evaluation of physical performance characteristics in volleyball. *Int Volleyball Technique* 3: 4–8, 1991.