

PEDIATRICS

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Recommendations for COVID-19 Vaccines in Infants, Children, and Adolescents, 2026-2027: Policy Statement

Committee on Infectious Diseases

DOI: 10.1542/peds.2026-079045

Journal: *Pediatrics*

Article Type: AAP Policy Statement

Citation: American Academy of Pediatrics Committee on Infectious Diseases. Recommendations for COVID-19 Vaccines in Infants, Children, and Adolescents, 2026-2027: Policy Statement. *Pediatrics*. 2026; doi: 10.1542/peds.2026-079045

This is a prepublication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

AMERICAN ACADEMY OF PEDIATRICS

POLICY STATEMENT

Organizational Principles to Guide and Define the Child Health Care System and/or Improve the Health of All Children

Recommendations for COVID-19 Vaccines in Infants, Children, and Adolescents, 2026-2027: Policy Statement

Committee on Infectious Diseases

Address correspondence to Roberta L. DeBiasi, MD, MS, FAAP. Email: rdebiasi@childrensnational.org

All authors contributed substantially to the conception and design; review and interpretation of relevant literature; drafting and revisions; and final approval of the published version.

All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. Author disclosures are provided at the end of this article. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors.

ABSTRACT. This policy statement updates the recommendations of the American Academy of Pediatrics (AAP) for the use of COVID-19 vaccines in the prevention of severe COVID-19 in children. A review of evidence supporting these recommendations is in the accompanying technical report (<https://doi.org/10.1542/peds.2026-079046>). These COVID-19 vaccine recommendations may change in future seasons or as additional variants emerge. The AAP recommends all infants and children 6 through 23 months of age who do not have contraindications receive 2026–2027 COVID-19 vaccine. The AAP also recommends a single dose of age-appropriate

2026–2027 COVID-19 vaccine for all children and adolescents 2 through 18 years of age at increased risk of severe COVID-19, regardless of prior COVID-19 vaccination status. Children 2 through 18 years of age not considered at increased risk of severe COVID-19 whose parent or guardian desires their protection from COVID-19 should be offered a single dose of age-appropriate 2026–2027 COVID-19 vaccine. Any available COVID-19 vaccine appropriate by age and health status can be administered and the most updated version of the COVID-19 vaccine that is available should be used.

ABBREVIATIONS: AAP, American Academy of Pediatrics; COID, Committee on Infectious Diseases.

INTRODUCTION

This policy statement updates the recommendations of the American Academy of Pediatrics (AAP)¹ for the use of COVID-19 vaccines in the prevention of severe COVID-19 in children. A review of evidence supporting these recommendations is in the accompanying technical report (<https://doi.org/10.1542/peds.2026-079046>).² These COVID-19 vaccine recommendations may change in future seasons or as additional variants emerge.

COVID-19 continues to be a cause of hospitalization^{3,4} and death⁵ in the pediatric population. COVID-19 vaccines are safe^{6,7,8,9,10,11,12,13} and effective¹⁴ in protecting individuals and populations against serious outcomes associated with SARS-CoV-2 infection, including postacute sequelae of SARS-CoV-2 infection (long COVID)¹⁵ and multisystem inflammatory syndrome in children (MIS-C).¹⁶

METHODOLOGY

The AAP Committee on Infectious Diseases (COID) developed these recommendations through an evidence-based process informed by scientific literature, public health data, and expert review. For COVID-19, the COID convened a subgroup of COID members and external experts with relevant

expertise. The subgroup worked in collaboration with strategic partners including the Vaccine Integrity Project and professional societies such as the American Medical Association and the Council on Medical Specialty Societies to access salient data sources.

The subgroup identified clinical questions and reviewed available data on disease epidemiology; morbidity; mortality; and vaccine safety, efficacy, and uptake, considering relevant domains of the Evidence to Recommendations framework.¹⁷ Conflict of interest disclosures were collected from all participants, and members with relevant conflicts recused themselves from evidence review and recommendations meetings as well as voting as required. Proposed recommendations were presented to the full COID for discussion and vote by eligible members. Approved recommendations were incorporated into this policy statement and the accompanying technical report. Additional information about the 2026-2027 AAP immunization recommendation development process can be found at <https://www.aap.org/en/patient-care/immunizations/aap-immunization-recommendations-process-2026/>.

RECOMMENDATIONS

- Infants and children 6 through 23 months of age are at high risk for severe COVID-19.^{3,4,5} The AAP recommends all infants and children in this age group who do not have contraindications^a receive 2026–2027 COVID-19 vaccine, as follows:
 - Those who are previously unvaccinated should receive an initial vaccine series.
 - Those who are previously vaccinated but did not complete their initial vaccine series should complete their initial vaccine series.

^a COVID-19 vaccine contraindication includes a history of severe allergic reaction (eg, anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine.

- Those who are previously vaccinated and completed their initial series should receive a single dose. This dose should be administered at least 8 weeks after the last dose was received.
- Those with a previous asymptomatic infection or symptomatic disease caused by SARS-CoV-2 should also receive COVID-19 vaccination.

Additional evidence regarding the burden of COVID-19 among infants and young children 6 through 23 months of age is provided in the accompanying technical report.²

- Children 6 months through 18 years of age who are moderately or severely immunocompromised require 2 or more doses of age-appropriate 2026–2027 COVID-19 vaccine depending on previous vaccination status.¹⁸ Refer to the [AAP Recommended Child and Adolescent Immunization Schedule](#) for dosing recommendations at [AAP.org/ImmunizationSchedule](#). Conditions associated with moderate or severe immunocompromise, as well as evidence regarding risk for severe COVID-19 and immune responses to COVID-19 vaccination in this population, are described in the technical report.²
- The AAP recommends a single dose of age-appropriate 2026–2027 COVID-19 vaccine for all children and adolescents 2 through 18 years of age in the following risk groups (as described in Table 1), regardless of prior COVID-19 vaccination status:
 - Persons at high risk of severe COVID-19³
 - Residents of long-term care facilities or other congregate settings
 - Persons who have never been vaccinated against COVID-19
 - Persons whose household contacts are at high risk for severe COVID-19^{3,19}

Further information on these risk groups and the rationale supporting vaccination recommendations is provided in the technical report.²

- Children 2 through 18 years of age not included in the risk groups above whose parent or guardian desires their protection from COVID-19 should be offered a single dose of age-

appropriate 2026–2027 COVID-19 vaccine.²⁰ This dose should be administered at least 8 weeks after the last dose was received.

- Any available COVID-19 vaccine appropriate by age and health status that is approved by the US Food and Drug Administration through a biologics license application can be used. The most updated version of the COVID-19 vaccine that is available should be used.

TIMING CONSIDERATIONS

In recent years, SARS-CoV-2 has circulated year-round with recurring summer and winter peaks.²¹ Despite this year-round activity, COVID-19–associated hospitalization rates have consistently been highest during the winter months, and infants and young children experience the greatest burden, including hospitalization rates that are higher and more sustained than those observed among older children and adolescents.⁴

Updated COVID-19 vaccines have generally become available in late summer or early fall. Children for whom vaccination is recommended and those desiring protection from COVID-19 should receive the updated vaccine as soon as it becomes available and they are eligible, rather than waiting for a particular point in the season. For most children, vaccination soon after the updated vaccine becomes available can provide an opportunity to develop protection before the winter months, when COVID-19 disease burden has generally been highest. When indicated, COVID-19 vaccine may be administered at the same visit as seasonal influenza vaccine; coadministration is supported by safety and immunogenicity data and may improve vaccine uptake and timeliness.

However, because SARS-CoV-2 circulates year-round, COVID-19 vaccination provides protection even when administered at other times of the year. Variations in specific timing of administration may occur because of access to vaccine, individual risk assessments, local epidemiology,

and timing of prior doses of vaccine. Therefore, eligible individuals who do not receive the vaccine in the fall can still receive COVID-19 vaccination at any time during the season.

ADDITIONAL RESOURCES

For additional clinical guidance on COVID-19, refer to the *Red Book: 2024 Report of the Committee on Infectious Diseases* (<https://publications.aap.org/redbook>). For more information on the dosing and administration of COVID-19 vaccines, storage and handling, reporting, and patient education, visit www.aap.org/covid-19.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements, clinical reports, and technical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

FINANCIAL/CONFLICT OF INTEREST DISCLOSURES: *Authors disclosed relevant relationships in accordance with American Academy of Pediatrics (AAP) policy to promote transparency and allow readers to assess relevance. The AAP reviews disclosures through its conflict-of-interest process and implements management strategies when appropriate. The presence of a disclosed relationship does not necessarily indicate a conflict of interest. Dr DeBiasi has disclosed serving as a research funding recipient with Pfizer for collaboration on research related to Lyme Disease; as a scientific advisory board consultant with the Foundation for the NIH advising on clinical trial design for the RECOVER study of Long COVID; and as a research funding recipient with the Stephen and Alexandra Cohen Foundation (related to Lyme and Tickborne Diseases). Disclosures of all committee members were reviewed, and it was determined if disclosures were related to the work of the statement. If a member had disclosures determined to be relevant to the work related to the statement they did not participate in making recommendations or developing the statement.*

FUNDING: *No external funding.*

This article is free to read on the publisher's website but is not Open Access. No part of this article may be reproduced, distributed, or transmitted in any form or by any means without prior written permission from the American Academy of Pediatrics.

COVID-19 SUBGROUP

Roberta L. DeBiasi, MD, MS, FAAP, Lead

Betsy C. Herold, MD, FAAP

Chandy C. John, MD, MS, FAAP

Sean T. O’Leary, MD, MPH, FAAP

Matthew Zahn, MD, FAAP

Dean Blumberg, MD, FAAP, Infectious Diseases Society of America

Richard Dang, PharmD, Aph, FCPA, American Pharmacists Association

Teresa Salaway, MHA, Medical Writer and Staff

Sunnah Kim, MS, RN, Staff

COMMITTEE ON INFECTIOUS DISEASES, 2026-2027

James D. Campbell, MD, MS, FAAP, Chairperson

Pia S. Pannaraj, MD, MPH, FAAP, Vice Chairperson

Monica I. Ardura, DO, MSCS, FAAP

Kristina A. Bryant, MD, FAAP, *Red Book Online* Team Lead

Eric J. Chow, MD, MS, MPH, FAAP

Roberta L. DeBiasi, MD, MS, FAAP

Claudia Espinosa, MD, MSc, FAAP

Robert W. Frenck, Jr., MD, FAAP

C. Mary Healy, MD

Betsy C. Herold, MD, FAAP

Chandy C. John, MD, MS, FAAP

Latania Logan, MD, MSPH, FAAP

Aaron M. Milstone, MD, MHS, FAAP

Angela L. Myers MD, MPH, FAAP

José R. Romero, MD, FAAP

Matthew Zahn, MD, FAAP

EX-OFFICIO

Sean T. O’Leary, MD, MPH, FAAP, Immunization Schedule Editor

David W. Kimberlin, MD, FAAP, *Red Book* Editor

Ritu Banerjee, MD, PhD, FAAP, *Red Book* Associate Editor

Elizabeth D. Barnett, MD, FAAP, *Red Book* Associate Editor

Ruth Lynfield, MD, FAAP, *Red Book* Associate Editor

Adam J. Ratner, MD, MPH, FAAP, *Red Book* Associate Editor

LIAISONS

Amina Ahmed, MD, FAAP, American Thoracic Society

Michelle Barton, MD, Canadian Paediatric Society

Carolyn Cleary, MD, FAAP, AAP Committee on Practice and Ambulatory Medicine

Andrew Handel, MD, FAAP, Pediatric Infectious Diseases Society

Denee J. Moore, MD, American Academy of Family Physicians

Jennifer L. Thompson, MD, American College of Obstetricians and Gynecologists

Juan P. Torres, MD, PhD, Sociedad Latinoamericana de Infectología Pediátrica

STAFF

Gillian Gibbs, MPH

ACKNOWLEDGMENT

The Committee on Infectious Diseases gratefully acknowledges Roberta DeBiasi, MD, FAAP, for her leadership in drafting the policy statement.

TABLE 1. Pediatric Populations Recommended to Receive the COVID-19 Vaccine During the 2026–2027 Season

Population	Characteristics
Infants and children 6 through 23 months of age	Recommended for all in this age group.
Children and adolescents who are moderately or severely immunocompromised	See Box 1 for examples of qualifying conditions and treatments. These children may require additional vaccine doses. ^a
Children and adolescents at high risk of severe COVID-19	See below for examples of medical conditions for which COVID-19 vaccination is recommended.
Residents of long-term care facilities or other congregate settings	Congregate care settings refer to places where individuals live together in structured environments outside of their home, including residential treatment facilities, group homes, emergency shelters, juvenile detention centers, etc.
Children and adolescents who have never been vaccinated against COVID-19	Those who have not previously received vaccine-derived protection against COVID-19.
Children and adolescents with household contacts at high risk for severe COVID-19	Household contacts may include infants and young children less than 24 months of age, adults aged ≥ 65 years, persons who are moderately or severely immunocompromised, and children, adolescents, or adults with medical conditions associated with severe COVID-19. ^{3,19}
Underlying conditions with common examples^{3,b}	
Chronic pulmonary disease	Asthma/reactive airway disease Chronic lung disease of prematurity Compromised respiratory function (eg, abnormality of airway, tracheostomy, or ventilator dependent)
Cardiovascular disease	Congenital heart disease
Gastrointestinal disorders	Feeding tube dependent Inflammatory bowel disease
Hepatic disease	Chronic liver disease
Hematologic disease	Sickle cell disease
Metabolic disorders	Diabetes mellitus
Obesity	Body mass index (BMI) $\geq$ the 95th percentile in children
Neurologic and neurodevelopmental conditions	Cerebral palsy Epilepsy Intellectual developmental disorder Compromised mobility (eg, wheelchair dependent)
Immunosuppressive conditions	See Box 1.
Rheumatologic, autoimmune disease	Systemic lupus erythematosus Juvenile idiopathic arthritis

^a Additional doses may be administered at ≥ 8 week intervals, informed by the clinical judgment of a health care provider and personal preference and circumstances. Refer to [AAP Recommended Child and Adolescent Immunization Schedule](#) for dosing guidance at [AAP.org/ImmunizationSchedule](https://www.aap.org/ImmunizationSchedule).

^b List of examples is not exhaustive.

BOX 1. Moderate and Severe Immunocompromising Conditions and Treatment²²

Individuals may be considered moderately or severely immunocompromised if they have one or more of the following conditions or treatments. Examples include, but are not limited to, the following:

- **Active treatment for solid tumor and hematologic malignancies**
- **Hematologic malignancies associated with poor responses to COVID-19 vaccines** regardless of current treatment status (e.g., chronic lymphocytic leukemia, non-Hodgkin lymphoma, multiple myeloma, acute leukemia)
- **Receipt of solid-organ transplant or an islet transplant and taking immunosuppressive therapy**
- **Receipt of chimeric antigen receptor (CAR)-T-cell therapy or hematopoietic cell transplant (HCT)^a**
- **Moderate or severe primary immunodeficiency** (e.g., common variable immunodeficiency disease, severe combined immunodeficiency, DiGeorge syndrome, Wiskott-Aldrich syndrome)
- **Advanced^b or untreated HIV infection**
- **Active treatment with immunosuppressive or immunomodulatory therapies^c**

^a Within 2 years of transplantation or taking immunosuppressive therapy

^b Includes individuals with HIV and CD4 cell counts less than 200/mm³, history of an AIDS-defining illness without immune reconstitution, or clinical manifestations of symptomatic HIV.

^c Examples include high-dose corticosteroids (i.e., 20 mg or more of prednisone or equivalent per day when administered for 2 or more weeks), alkylating agents, antimetabolites, transplant-related immunosuppressive drugs, cancer chemotherapeutic agents classified as severely immunosuppressive, tumor necrosis factor (TNF) blockers, and other biologic agents that are immunosuppressive or immunomodulatory (e.g., B-cell-depleting agents).

REFERENCES

1. American Academy of Pediatrics, Committee on Infectious Diseases. Recommendations for COVID-19 vaccines in infants, children, and adolescents: policy statement. *Pediatrics*. 2025;156(5):e2025073924
2. American Academy of Pediatrics, Committee on Infectious Diseases. Recommendations for COVID-19 vaccines in infants, children, and adolescents: Technical report. *Pediatrics*. 2026; in press
3. Free RJ, Patel K, Taylor CA, et al. Hospitalization for COVID-19 and risk factors for severe disease among children: 2022–2024. *Pediatrics*. 2025;156(3):e2025072788. Doi: 10.1542/peds.2025-072788
4. Centers for Disease Control and Prevention. COVID-19 Hospitalization Surveillance Network (COVID-NET). Accessed August 4, 2026. <https://www.cdc.gov/covid/php/covid-net/index.html>
5. Ahmad FB, Cisewski JA. Provisional COVID-19 Mortality Surveillance. National Center for Health Statistics. 2026. Accessed July 10, 2026. Doi: <https://dx.doi.org/10.15620/cdc/20250305007>
6. Hause AM, Shay DK, Klein NP, et al. Safety of COVID-19 vaccination in United States children ages 5 to 11 years. *Pediatrics*. 2022;150(2):e2022057313. Doi: 10.1542/peds.2022-057313
7. Hause AM, Marquez P, Zhang B, et al. Safety monitoring of mRNA COVID-19 vaccine third doses among children aged 6 months–5 years—United States, June 17, 2022–May 7, 2023. *MMWR Morb Mortal Wkly Rep*. 2023;72(23):621–626. Doi: 10.15585/mmwr.mm7223a2

8. Hesse EM, Hause A, Myers T, et al. COVID-19 Vaccine safety first year findings in adolescents. *Pediatrics*. 2023;151(5):e2022060295. Doi: 10.1542/peds.2022-060295
9. Hause AM, Marquez P, Zhang B, et al. Safety monitoring of bivalent COVID-19 mRNA vaccine booster doses among children aged 5–11 years—United States, October 12–January 1, 2023. *MMWR Morb Mortal Wkly Rep*. 2023;72(2):39–43. Doi: 10.15585/mmwr.mm7202a5
10. Hause AM, Marquez P, Zhang B, et al. COVID-19 mRNA vaccine safety among children aged 6 months–5 years—United States, June 18, 2022–August 21, 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71(35):1115–1120. Doi: 10.15585/mmwr.mm7135a3
11. Goddard K, Donahue JG, Lewis N, et al. Safety of COVID-19 mRNA vaccination among young children in the Vaccine Safety Datalink. *Pediatrics*. 2023;152(1):e2023061894. Doi: 10.1542/peds.2023-061894
12. Cortese MM, Taylor AW, Akinbami LJ, et al. Surveillance for multisystem inflammatory syndrome in US children aged 5–11 years who received Pfizer-BioNTech COVID-19 vaccine, November 2021 through March 2022. *J Infect Dis*. 2023;228(2):143–148. Doi: 10.1093/infdis/jiad051
13. Yousaf AR, Cortese MM, Taylor AW, et al; MIS-C Investigation Authorship Group. Reported cases of multisystem inflammatory syndrome in children aged 12–20 years in the USA who received a COVID-19 vaccine, December, 2020, through August, 2021: a surveillance investigation. *Lancet Child Adolesc Health*. 2022;6(5):303–312. Doi: 10.1016/S2352-4642(22)00028-1

14. Irving SA, Rowley EA, Chickery S, et al. Effectiveness of 2024–2025 COVID-19 vaccines in children in the United States—VISION, August 29, 2024–September 2, 2025. *MMWR Morb Mortal Wkly Rep*. 2025;74(40):607–614.
DOI: <http://dx.doi.org/10.15585/mmwr.mm7440a1>
15. Yousaf AR, Mak J, Gwynn L, et al. COVID-19 vaccination and odds of post-COVID-19 condition symptoms in children aged 5 to 17 years. *JAMA Netw Open*. 2025;8(2):e2459672. Doi: 10.1001/jamanetworkopen.2024.59672
16. Hamad Saied M, van der Griend L, van Straalen JW, Wulffraat NM, Vastert S, Jansen MHA. The protective effect of COVID-19 vaccines on developing multisystem inflammatory syndrome in children (MIS-C): a systematic literature review and meta-analysis. *Pediatr Rheumatol Online*. 2023;21(1):80. Doi: 10.1186/s12969-023-00848-1
17. Ahmed F, Temte JL, Campos-Outcalt D, Schünemann HJ. Methods for developing evidence-based recommendations by the Advisory Committee on Immunization Practices (ACIP) of the U.S. Centers for Disease Control and Prevention (CDC). *Vaccine*. 2011;29(49):9171–9176. Doi: 10.1016/j.vaccine.2011.08.005
18. Roper LE, Godfrey M, Link-Gelles R, et al. Use of additional doses of 2024–2025 COVID-19 vaccine for adults aged ≥ 65 years and persons aged ≥ 6 months with moderate or severe immunocompromise: recommendations of the Advisory Committee on Immunization Practices—United States, 2024. *MMWR Morb Mortal Wkly Rep*. 2024;73(49):1118–1123.
Doi: 10.15585/mmwr.mm7349a2
19. Yek C, Warner S, Wiltz JL, et al. Risk factors for severe COVID-19 outcomes among persons aged ≥ 18 years who completed a primary COVID-19 vaccination series—465 health care facilities,

United States, December 2020–October 2021. *MMWR Morb Mortal Wkly Rep.* 2022;71(1):19–25. Doi: 10.15585/mmwr.mm7101a4

20. National Center for Immunization and Respiratory Diseases, US Centers for Disease Control and Prevention. Use of 2025-2026 COVID-19 Vaccines: Work Group Considerations. Accessed August 4, 2026. <https://www.cdc.gov/acip/downloads/slides-2025-04-15-16/05-Panagiotakopoulos-COVID-508.pdf>
21. Centers for Disease Control and Prevention. The National Respiratory and Enteric Virus Surveillance System (NREVSS). Accessed June 30, 2026. <https://www.cdc.gov/nrevss/php/dashboard/index.html>
22. Centers for Disease Control and Prevention. COVID-19 Vaccination Guidance for People Who Are Immunocompromised. Updated November 4, 2025. Accessed July 9, 2026. <https://www.cdc.gov/covid/hcp/vaccine-considerations/immunocompromised.html>