



# 抗人 CD8 流式单抗(iFluor™488)[OKT8]

---

货号: ZX038R1-025; ZX038R1-100                      种属反应: 人 CD8 蛋白  
抗体亚型: Mouse/IgG2a kappa                      荧光标记染料: iFluor™488 (Ex 499nm/ Em 520nm)  
抗体来源: 杂交瘤细胞 (OKT8) 悬浮培养, 亲和层析纯化  
保存液成分: PBS pH7.4, 0.01% NaN<sub>3</sub>  
应用: 免疫荧光, 流式细胞检测  
包装规格: 25μg/支 (100μg/支), 0.5mg/mL      贮存运输: 2~8°C, 避光      保质期: 2 年  
使用方法: 每 100μL 反应体系 (10<sup>6</sup> 个细胞) 加入 1μL 抗体, 即 1:100 稀释。建议优化稀释比例。

## 简介:

本公司抗体产品经先进的杂交瘤细胞 (或 CHO 细胞) 无血清悬浮培养技术表达, 避免腹水表达中宿主抗体的污染。经亲和层析纯化, 抗体纯度可达 95% 以上, 且可保持较高的抗体活性。本产品单克隆抗体 (OKT8) 可特异性结合人细胞表面蛋白 CD8, 并经 iFluor™488 标记。

**注意事项:** 本试剂只适用于科研应用, 不可用于临床检测。

## 参考文献:

1. Nyanhete TE, Frisbee AL, Bradley T, et al. HLA class II-Restricted CD8+ T cells in HIV-1 Virus Controllers. *Sci Rep.* 2019;9(1):10165. Published 2019 Jul 15.
2. Davenport B, Eberlein J, Nguyen TT, et al. Aging boosts antiviral CD8+T cell memory through improved engagement of diversified recall response determinants. *PLoS Pathog.* 2019;15(11):e1008144. Published 2019 Nov 7.
3. Perdomo-Celis F, Taborda NA, Rugeles MT. CD8+ T-Cell Response to HIV Infection in the Era of Antiretroviral Therapy. *Front Immunol.* 2019;10:1896. Published 2019 Aug 9.
4. Khan AA, Srivastava R, Lopes PP, et al. Asymptomatic memory CD8+ T cells: from development and regulation to consideration for human vaccines and immunotherapeutics. *Hum Vaccin Immunother.* 2014;10(4):945–963.

**企业名称:** 浙江正熙生物医药有限公司

**企业地址:** 中国浙江省湖州市德清县阜溪街道长虹东街 926 号千人计划产业园 1 号楼

**企业网址:** <https://www.zxbiopharm.com/>

**联系方式:** 0572-8232782



# Anti-Human CD8 Monoclonal Antibody (iFluor™488) [OKT8]

---

**Catalog:** ZX038R1-025; ZX038R1-100      **Antigen:** Against Human CD8  
**Antibody Host:** Mouse/IgG2a kappa      **Fluorescent Labeling Dye:** iFluor™488 (Ex 499nm/ Em 520nm)  
**Antibody Origin:** Hybridomas (Clone: OKT8) Culture, Affinity Purification  
**Formulation:** PBS pH7.4, 0.01% NaN<sub>3</sub>  
**Application:** FC (Flow Cytometry), IF (Immunofluorescence)  
**Package Specification:** 25µg/Vial(100µg/Vial), 0.5mg/mL      **Guarantee Period:** 2 years  
**Storage & Handling:** stored between 2°C and 8°C, protected from light exposure. **Do not freeze.**  
**Recommended Usage:** ≤1 µL /10<sup>6</sup> cells in 100 µl volume per test (Maximum dilution 1:100)

## Product Information:

To improve the quality of monoclonal antibodies for research purpose, ZXBio develops novel cell culturing technologies to grow hybridomas/CHO cells in serum-free condition. This method is superior to traditional ascites production method, which always contaminates the purified antibodies with host proteins. Using affinity purification, ZXBio is able to purify monoclonal antibodies at a purity higher than 95%.

Anti-Human CD8 Monoclonal Antibody [OKT8] specifically binds human T cell membrane CD8 surface antigen and the product is labeled with iFluor™488.

**Application Notes:** For research use only, Not for use in diagnostic procedures.

## Reference:

- 1.Nyanhete TE, Frisbee AL, Bradley T, et al. HLA class II-Restricted CD8+ T cells in HIV-1 Virus Controllers. *Sci Rep.* 2019;9(1):10165. Published 2019 Jul 15.
- 2.Davenport B, Eberlein J, Nguyen TT, et al. Aging boosts antiviral CD8+T cell memory through improved engagement of diversified recall response determinants. *PLoS Pathog.* 2019;15(11):e1008144. Published 2019 Nov 7.
- 3.Perdomo-Celis F, Taborda NA, Rugeles MT. CD8+ T-Cell Response to HIV Infection in the Era of Antiretroviral Therapy. *Front Immunol.* 2019;10:1896. Published 2019 Aug 9.
- 4.Khan AA, Srivastava R, Lopes PP, et al. Asymptomatic memory CD8+ T cells: from development and regulation to consideration for human vaccines and immunotherapeutics. *Hum Vaccin Immunother.* 2014;10(4):945–963.

**Corporation Name:** ZXBio Co., Ltd.

**Corporation Address:** No. 926 Changhong East Rd, Deqing Xian, Huzhou Shi, Zhejiang, China

**Corporate Website:** <https://www.zxbiopharm.com/>

**Corporate Phone:** 0572-8232782