



Generalized Anxiety Disorder

Overview

Occasional anxiety is a normal part of life. You might feel anxious when faced with a problem at work, before taking a test, or making an important decision. But anxiety disorders involve more than temporary worry or fear. For a person with an anxiety disorder, the anxiety does not go away and can get worse over time. The feelings can interfere with daily activities such as job performance, school work, and relationships. There are several different types of anxiety disorders. Examples include generalized anxiety disorder, panic disorder, and social anxiety disorder.

Signs and Symptoms

Generalized Anxiety Disorder

People with generalized anxiety disorder display excessive anxiety or worry for months and face several anxiety-related symptoms.

Generalized anxiety disorder symptoms include:

- Restlessness or feeling wound-up or on edge
- Being easily fatigued
- Difficulty concentrating or having their minds go blank
- Irritability
- Muscle tension
- Difficulty controlling the worry
- Sleep problems (difficulty falling or staying asleep or restless, unsatisfying sleep)

Panic Disorder

People with panic disorder have recurrent unexpected panic attacks, which are sudden periods of intense fear that may include palpitations, pounding heart, or accelerated heart rate; sweating; trembling or shaking; sensations of shortness of breath, smothering, or choking; and feeling of impending doom.

Panic disorder symptoms include:

- Sudden and repeated attacks of intense fear
- Feelings of being out of control during a panic attack
- Intense worries about when the next attack will happen
- Fear or avoidance of places where panic attacks have occurred in the past

Social Anxiety Disorder

People with social anxiety disorder (sometimes called “social phobia”) have a marked fear of social or performance situations in which they expect to feel embarrassed, judged, rejected, or fearful of offending others.

Social anxiety disorder symptoms include:

- Feeling highly anxious about being with other people and having a hard time talking to them
- Feeling very self-conscious in front of other people and worried about feeling humiliated, embarrassed, or rejected, or fearful of offending others
- Being very afraid that other people will judge them
- Worrying for days or weeks before an event where other people will be
- Staying away from places where there are other people
- Having a hard time making friends and keeping friends
- Blushing, sweating, or trembling around other people
- Feeling nauseous or sick to your stomach when other people are around

Evaluation for an anxiety disorder often begins with a visit to a primary care provider. Some physical health conditions, such as an overactive thyroid or low blood sugar, as well as taking certain medications, can imitate or worsen an anxiety disorder. A thorough mental health evaluation is also helpful, because anxiety disorders often co-exist with other related conditions, such as depression or obsessive-compulsive disorder.

Risk Factors

Researchers are finding that genetic and environmental factors, frequently in interaction with one another, are risk factors for anxiety disorders. Specific factors include:

- Shyness, or behavioral inhibition, in childhood
- Being female
- Having few economic resources
- Being divorced or widowed
- Exposure to stressful life events in childhood and adulthood
- Anxiety disorders in close biological relatives
- Parental history of mental disorders
- Elevated afternoon cortisol levels in the saliva (specifically for social anxiety disorder)

Treatments and Therapies

Anxiety disorders are generally treated with psychotherapy, medication, or both.

Psychotherapy

Psychotherapy or “talk therapy” can help people with anxiety disorders. To be effective, psychotherapy must be directed at the person’s specific anxieties and tailored to his or her needs. A typical “side effect” of psychotherapy is temporary discomfort involved with thinking about confronting feared situations.

Cognitive Behavioral Therapy (CBT)

CBT is a type of psychotherapy that can help people with anxiety disorders. It teaches a person different ways of thinking, behaving, and reacting to anxiety-producing and fearful situations. CBT can also help people learn and practice social skills, which is vital for treating social anxiety disorder.

Two specific stand-alone components of CBT used to treat social anxiety disorder are **cognitive therapy** and **exposure therapy**. Cognitive therapy focuses on identifying, challenging, and then neutralizing unhelpful thoughts underlying anxiety disorders.

Exposure therapy focuses on confronting the fears underlying an anxiety disorder in order to help people engage in activities they have been avoiding. Exposure therapy is used along with relaxation exercises and/or imagery. One study, called a meta-analysis because it pulls together all of the previous studies and calculates the statistical magnitude of the combined effects, found that cognitive therapy was superior to exposure therapy for treating social anxiety disorder.

CBT may be conducted individually or with a group of people who have similar problems. Group therapy is particularly effective for social anxiety disorder. Often “homework” is assigned for participants to complete between sessions.

Self-Help or Support Groups

Some people with anxiety disorders might benefit from joining a self-help or support group and sharing their problems and achievements with others. Internet chat rooms might also be useful, but any advice received over the Internet should be used with caution, as Internet acquaintances have usually never seen each other and false identities are common. Talking with a trusted friend or member of the clergy can also provide support, but it is not necessarily a sufficient alternative to care from an expert clinician.

Stress-Management Techniques

Stress management techniques and meditation can help people with anxiety disorders calm themselves and may enhance the effects of therapy. While there is evidence that aerobic exercise has a calming effect, the quality of the studies is not strong enough to support its use as treatment. Since caffeine, certain illicit drugs, and even some over-the-counter cold medications can aggravate the symptoms of anxiety disorders, avoiding them should be considered. Check with your physician or pharmacist before taking any additional medications.

The family can be important in the recovery of a person with an anxiety disorder. Ideally, the family should be supportive but not help perpetuate their loved one’s symptoms.

Medication

Medication does not cure anxiety disorders but often relieves symptoms. Medication can only be prescribed by a medical doctor (such as a psychiatrist or a primary care provider), but a few states allow psychologists to prescribe psychiatric medications.

Medications are sometimes used as the initial treatment of an anxiety disorder, or are used only if there is insufficient response to a course of psychotherapy. In research studies, it is common for patients treated with a combination of psychotherapy and medication to have better outcomes than those treated with only one or the other.

The most common classes of medications used to combat anxiety disorders are antidepressants, anti-anxiety drugs, and beta-blockers (visit [Mental Health Medications](#)). Be aware that some medications are effective only if they are taken regularly and that symptoms may recur if the medication is stopped.

Antidepressants

Antidepressants are used to treat depression, but they also are helpful for treating anxiety disorders. They take several weeks to start working and may cause side effects such as headache, nausea, or difficulty sleeping. The side effects are usually not a problem for most people, especially if the dose starts off low and is increased slowly over time.

Please Note: Although antidepressants are safe and effective for many people, they may be risky for children, teens, and young adults. A “black box” warning—the most serious type of warning that a prescription can carry—has been added to the labels of antidepressants. The labels now warn that antidepressants may cause some people to have suicidal thoughts or make suicide attempts. For this reason, anyone taking an antidepressant should be monitored closely, especially when they first start taking the medication.

Anti-Anxiety Medications

Anti-anxiety medications help reduce the symptoms of anxiety, panic attacks, or extreme fear and worry. The most common anti-anxiety medications are called benzodiazepines. Benzodiazepines are first-line treatments for generalized anxiety disorder. With panic disorder or social phobia (social anxiety disorder), benzodiazepines are usually second-line treatments, behind antidepressants.

Beta-Blockers

Beta-blockers, such as propranolol and atenolol, are also helpful in the treatment of the physical symptoms of anxiety, especially social anxiety. Physicians prescribe them to control rapid heartbeat, shaking, trembling, and blushing in anxious situations.

Choosing the right medication, medication dose, and treatment plan should be based on a person’s needs and medical situation, and done under an expert’s care. Only an expert clinician can help you decide whether the medication’s ability to help is worth the risk of a side effect. Your doctor may try several medicines before finding the right one.

You and your doctor should discuss:

- How well medications are working or might work to improve your symptoms
- Benefits and side effects of each medication
- Risk for serious side effects based on your medical history
- The likelihood of the medications requiring lifestyle changes
- Costs of each medication
- Other alternative therapies, medications, vitamins, and supplements you are taking and how these may affect your treatment
- How the medication should be stopped. Some drugs can’t be stopped abruptly but must be tapered off slowly under a doctor’s supervision.

For more information, please visit [Medications Health Topic webpage](#) developed by the National Institute of Mental Health (NIMH). Please note that any information on this website regarding medications is provided for educational purposes only and may be outdated. Information about medications changes frequently. Please visit the U.S. Food and Drug Administration ([FDA website](#)) for the latest information on warnings, patient medication guides, or newly approved medications.

Learn More

Free Booklets and Brochures

- **Generalized Anxiety Disorder (GAD): When Worry Gets Out of Control:** A brochure on GAD that explains the signs, symptoms, and treatment
- **Obsessive-Compulsive Disorder: When Unwanted Thoughts Take Over:** A brochure on OCD that explains the signs, symptoms, and treatments
- **Panic Disorder: When Fear Overwhelms:** A brochure on panic disorder that explains the signs, symptoms, and treatments
- **Social Anxiety Disorder: More Than Just Shyness:** This brochure discusses symptoms, causes, and treatments for social anxiety disorder (also called social phobia).

Multimedia

- **Watch: Bullying Exerts Psychological Effects into Adulthood**— Once considered a childhood rite of passage, bullying is no longer seen as benign. Its effects linger well into adulthood. Bullies and victims alike are at risk for psychiatric problems such as anxiety, depression, substance abuse, and suicide when they become adults, according to a study partially funded by the NIMH that was published in the April 2013 issue of *JAMA Psychiatry*.
- **See: Multimedia about Anxiety Disorders**

Federal Resources

- **Anxiety Disorders** MedlinePlus – also en [Español](#)
- **Anxiety Disorders** [NIHSeniorHealth.gov](#)
- **Specific Phobias:** U.S. Department of Veterans Affairs

Research and Statistics

- **Join a Study: Adults - Anxiety Disorders**
- **Join a Study: Children - Anxiety Disorders**
- **Journal Articles:** References and abstracts from MEDLINE/PubMed (National Library of Medicine).
- **Statistics: Anxiety Disorder in Adults**
- **Statistics: Anxiety Disorder in Children**

Last Revised: March 2016

National Institute of Mental Health (2013). *Eating Disorders*. Retrieved March 20, 2015, from <https://www.nimh.nih.gov/health/topics/anxiety-disorders/index.shtml>

Retrieved From: <https://www.nimh.nih.gov/health/topics/anxiety-disorders/index.shtml>