


Graduate Level Math Textbook

---

# Mapping the Cosmos:

## The Complete Blueprint of Existence



by Kiet Anh Le

**A Graduate Level Math Textbook**

**Mapping the Cosmos: The Complete Blueprint of Existence**

**Mapping the Cosmos:  
The Complete Blueprint of Existence  
KIET ANH LE**

© 2026 Kiet Anh Le  
All Rights Reserved

# COPYRIGHT PAGE

**Mapping the Cosmos: The Complete Blueprint of Existence** by. KIET ANH LE

© 2026 **Kiet Anh Le**

All Rights Reserved.

No part of this book may be reproduced, distributed, or transmitted in any form or by any means — including photocopying, recording, or other electronic or mechanical methods — without the prior written permission of the publisher, except in the case of brief quotations used in critical reviews and certain noncommercial uses permitted by copyright law.

This is a work of **Graduate Level Math Textbook of Existence**.

Some names and identifying details have been changed to protect the privacy of individuals.

The references in this book are **entirely symbolic, metaphorical, and fictionalized**, representing internal profound philosophical experiences rather than literal historical events. Any resemblance to actual persons, living or dead, or actual events is **coincidental** or used purely as **literary metaphor**.

For permissions, inquiries, or information, please contact:

**Kiet Anh Le**

First Edition.

Printed in the United States of America.



**Library of Congress Cataloging-in-Publication Data**

Names: **Le, Kiet Anh**, author.

Title: A Graduate Level Math Textbook *Mapping the Cosmos: The Complete Blueprint of Existence* / Kiet Anh Le.

Description: First edition. | Los Angeles, CA : [Your Publishing Imprint], 2026.

Identifiers:

LCN 2026##### (print) | ISBN 975-0-00-000000-0 (paperback) | ISBN 927-0-00-000000-7 (ebook)

Subjects: MYZNCXK —

1. Graduate Level Math Textbook in Existence.

Classification: XWYX KX 421.3 .L9 2026 | ELP 143.2/1—jx12

*The data above is a placeholder to be replaced with official records.*

# DEDICATION PAGE

**“Who introduced me to Existentialism in the 10th Grade.”  
Dr. William Lomax, Ms. Maria Soldevilla, and Ms. Grace Park  
(Benjamin Franklin High School, LAUSD)**

**“For giving me that Starbucks boost in confidence.”  
Dr. Harold L. Rogler (Santa Monica College)**

**“For making me slave for life and for death to get that B+ in C++ Data Structure.”  
Howard Stahl (Santa Monica College)**

**“For failing me in Math 10: Discrete Structures for not showing up to class and  
not studying.”  
Gail Edinger (Santa Monica College)**

**“For giving me a B in Java Programming - undeserved.”  
Dr. Abbas Dehkhoda (Santa Monica College)**

**“They all taught me one thing.”**

**“Value.” - Rain**

**I dedicate this to Madison Powers, for being my Montblanc.**

## **CHAPTER 1 - The Parallax Of Existence**

The math of Law is: Reason + Logic

What is The math of Existence?.

What is mathematics?

A set of principles.

To say that principles are Laws is incorrect.

Principles Guide Us.

Math is guidance.

To say that The math of Existence is just philosophy is incorrect.

It is a brand new subject.

To say That The math of language is The math of Existence is also incorrect.

Existence is an instantiation of Law.

No It Is Not.

That's the kind of Existence in Law & Limit: The Architecture Of The Infinite.

The Existence I am talking about now is different.

There is No-Exist.

To say that No-Law is No-Exist is also incorrect.

You must know No-Exist to understand The math of Existence.

No-Exist is not a dimension.

It is not no-dimension.

It is not not.

No proof? Incorrect.

You need proof.

It's the heart of mathematics.

If No-Existence exists, isn't that Existence.

No.

You can't say the alphabet of The math of Existence is

A. No-Existence B. Existence

It doesn't work like that.

These are newer higher concepts.

It's easy to get lost.

It's happening to me as I write this.

Can't you say that Existence is an instantiation of Law.

Existence exists outside of Law.

It is a plane.

And No-Exist is the void beyond the plane.

Yes.

But isn't that Existence?.

No.

You're thinking INSIDE the box.

What's beyond No-Exist.

That's where The math of Existence comes in.

It's not dimensions you are envisioning.

I call it The Parallax of Existence.

So you have

Epsilon = Existence (Law&Limit: The Architecture Of The Infinite)

Gamma = Law

Omega = Existence

Void = No-Exist

N = Beyond

These are The Parallax of Existence.

Stop thinking these are dimensions.

Stop thinking inside the box.

R = Prime Number (Random)

R is actually a "jump" in space.

How does it ties in with The Parallax of Existence.

R exists on the fringe of Omega.

That is why Prime is so special.

Does it mean we have a channel from Epsilon to Omega.

No.

Only an understanding.

Not yet. At least.

\*laugh\*

Imagine building a world inside a video game, or trying to understand how reality works without using standard high school geometry or physics formulas. To make sense of these concepts, we have to throw away traditional "boxes"—like thinking in terms of 3D space, standard timelines, or regular dimensions—and look at the hidden rules underneath.

Here is a simple breakdown of how these concepts fit together:

### 1. Mathematics is a Compass, Not just Numbers

Most people think math is just long division or solving for  $x$ . But at its core, **math is guidance**. It is a set of foundational principles that show us how things relate to each other.

- **Law** uses reason and logic to show how things *must* behave.
- **Principles** don't force you like rigid walls; they **guide** you like a map or a compass.

### 2. A Brand-New Subject: The Math of Existence

If you say "The Math of Existence," people might assume you mean philosophy or computer code. But this is a completely distinct, standalone subject. It isn't just about language, and it isn't just about how reality is built inside a specific system of laws. It is the framework for *where* reality sits and what lies completely outside of it.

### 3. The Map: The Parallax of Existence

Instead of thinking about spatial dimensions (like length, width, height, or time), picture different fundamental "states" or stages of reality. These form **The Parallax of Existence**:

- **Gamma ( $\Gamma$ ): The Law** — The pure reason, logic, and core rules.
- **Epsilon ( $\epsilon$ ): Local Existence** — Existence as an instantiation of Law (the structured universe bound by rules, like in *Law & Limit: The Architecture of the Infinite*).
- **Omega ( $\Omega$ ): True Existence** — The broader plane of reality that actually exists *outside* of Law.

- **Void: No-Exist** — The blank, empty space beyond the plane of True Existence. It is not a dimension, nor is it "nothingness" in a simple sense. It isn't "not," and it isn't an "alphabet" where you just toggle between 1s and 0s.
- **N: Beyond** — What lies past No-Exist. This is the realm where traditional thinking completely breaks down and where the Math of Existence operates.

#### 4. Prime Numbers ( $R$ ) and the "Jump"

Why are Prime Numbers (2, 3, 5, 7, 11, ...) so strange and special in regular math?

In this framework,  $R$  (Prime Numbers / Randomness) isn't just a number that can't be divided evenly.  $R$  represents a "jump" in space.

- **Where does  $R$  live?** It sits right on the outer fringe or border of **Omega** (True Existence).
- Because  $R$  touches the very edge of Omega, prime numbers act like unique signal flares on the boundary of reality.

#### 5. What Does It Mean for Us?

Does this mean we can build a teleporter or send a message directly from our structured world ( $\epsilon$ ) straight out into deep Omega ( $\Omega$ )?

Not right now. There is no physical bridge or channel connecting them yet. What we have right now is **understanding**—the blueprint and the logic that allows us to map out what's beyond the box, even if we can't step through the door just yet.

## The Formal Grammar of the Parallax Mechanics

### Primary Symbol Sets & Domain Definitions

- $\mathcal{E}$  (Epsilon): Instantiated Existence bounded by Law.
- $\Gamma$  (Gamma): Law / Axiomatic Constraints ( $\text{Law} = \text{Reason} + \text{Logic}$ ).
- $\Omega$  (Omega): Unbounded/True Existence (The Outer Plane beyond Law).
- $\emptyset_X$  (Void / No-Exist): The non-dimensional null-boundary beyond  $\Omega$ .
- $\mathcal{N}$  (N-Beyond): The realm transcending No-Exist.
- $\mathfrak{P}_p$  (Prime Parallax Operator): A non-continuous quantum shift ("jump") on the fringe of  $\Omega$ .

### Novel Operators & Axioms

1. **Guidance Mapping ( $\rightarrow$ ):**  $A \rightarrow B$  denotes principles guiding  $A$  into structural alignment with  $B$  ( $A \rightarrow B \neq A \implies B$ ).
2. **Transcendent Projection ( $\Delta \nabla$ ):** An operator acting across the boundary from  $\Omega$  into  $\mathcal{N}$  through  $\emptyset_X$ .
3. **Fringe Shift ( $\mathfrak{q}_p$ ):**  $\mathfrak{P}_p \mathfrak{q}_p \Omega$  represents a discontinuous state-jump localized at the exterior boundary of Omega.
4. **Non-Existential Contraction ( $\boxtimes$ ):**  $\emptyset_X \boxtimes X \equiv \text{un-bound } X \text{ without falling into dimensional negation.}$

### Problem Set: The Parallax of Existence

#### Problem 1

Evaluate the Guidance Mapping of Law: Given  $\Gamma = \text{Reason} + \text{Logic}$ , show whether  $\Gamma \rightarrow \mathcal{E}$  implies  $\mathcal{E}$  is an instantiation of  $\Gamma$ .



**Problem 2**

Let  $\alpha \in \Omega$  and  $\beta \in \mathcal{E}$ . Prove that  $\alpha \boxtimes \beta$  is non-empty under the condition that  $\Omega$  lies strictly outside the boundary of  $\Gamma$ .

**Problem 3**

Compute the state of a localized continuous vector  $v \in \Omega$  after applying the Prime Jump operator  $\mathfrak{P}_p \rightsquigarrow_p v$ .

**Problem 4**

Demonstrate why the binary alphabet  $\mathcal{A} = \{\emptyset_{\mathbf{X}}, \Omega\}$  fails to span the Parallax space  $\mathcal{P} = \{\mathcal{E}, \Gamma, \Omega, \emptyset_{\mathbf{X}}, \mathcal{N}\}$ .

**Problem 5**

Given a boundary flux  $\Phi$  operating at the fringe of  $\Omega$ , derive the divergence  $\nabla_{\mathcal{P}} \cdot \Phi$  as  $\Phi$  transitions into  $\emptyset_{\mathbf{X}}$ .

**Problem 6**

Define the non-dimensional boundary conditions for  $\emptyset_{\mathbf{X}}$  such that  $\dim(\emptyset_{\mathbf{X}}) \neq 0$  and  $\dim(\emptyset_{\mathbf{X}}) \neq \emptyset$ .

**Problem 7**

Let  $\mathcal{N}$  be the region beyond  $\emptyset_{\mathbf{X}}$ . Solve for the projection field  $\Psi_{\mathcal{N}} = \Omega \Delta \nabla \emptyset_{\mathbf{X}}$ .

**Problem 8**

Verify if the composite transformation  $\mathcal{E} \rightarrow \Gamma \rightarrow \Omega$  establishes a physical state-channel between  $\mathcal{E}$  and  $\Omega$ .

**Problem 9**

Given prime sequence  $p_k \in \mathbb{P}$ , evaluate the asymptotic density of fringe jumps  $\sum_{k=1}^n (\mathfrak{R}_{p_k} \leftrightarrow_p \Omega)$  as  $n \rightarrow \infty$ .

**Problem 10**

Show that Reason + Logic  $\not\subseteq$  Math of Existence.

**Problem 11**

Calculate the un-bound state  $\mathcal{O}_X \boxtimes \Gamma$  when applied to an arbitrary principle  $P$ .

**Problem 12**

Formulate the Parallax Tensor  $P_{\mu\nu}$  spanning components  $(\mathcal{E}, \Gamma, \Omega, \mathcal{O}_X, \mathcal{N})$  and calculate its trace  $\text{Tr}(P_{\mu\nu})$ .

**Problem 13**

Let  $J_p$  be the spatial displacement magnitude of a Prime Jump  $\mathfrak{R}_p$ . Determine the path integral  $\int_{\partial\Omega} J_p d\Omega$ .

**Problem 14**

Prove that  $\mathcal{N} \neq \Omega \cup \mathcal{O}_X$ .

**Problem 15**

If  $P$  is a principle providing guidance  $(P \rightarrow X)$ , show that  $P$  cannot be expressed as a fixed constraint equation  $C(X) = 0$ .

#### Problem 16

Determine the stability of a localized wave packet  $\psi \in \mathcal{E}$  under the influence of an adjacent fringe operator  $\mathfrak{R}_p$ .

#### Problem 17

Evaluate the Transcendent Projection  $\Delta \nabla$  over the void topology  $\emptyset_{\mathbf{X}}: \Delta \nabla (\mathcal{E} \boxtimes \mathcal{N})$ .

#### Problem 18

Construct the truth valuation function  $V: \text{Claims} \rightarrow \{0, 1\}$  for the statement  $S$ : “No-Law  $\equiv$  No-Exist”.

#### Problem 19

Calculate the commutator  $[\mathfrak{R}_p, \Gamma]$  acting on a state function  $\psi \in \Omega$ .

#### Problem 20

Derive the complete Parallax Equation mapping the progression from bounded law to transcendent un-existence:  $\mathcal{E} \rightarrow \Gamma \rightarrow \Omega \rightarrow \emptyset_{\mathbf{X}} \rightarrow \mathcal{N}$ .

### Solutions & Proofs

#### Solution 1

$\Gamma \rightarrow \mathcal{E}$  denotes *guidance* rather than *instantiation* ( $\Gamma \not\rightarrow \mathcal{E}$ ). While  $\mathcal{E}$  is structured under Law in Law & Limit, True Existence ( $\Omega$ ) operates outside  $\Gamma$ . Thus,  $\Gamma \rightarrow \mathcal{E}$  validates guidance, but does not reduce existence to an instantiation of law.

Solution 2

Since  $\Omega$  exists outside  $\Gamma$ ,  $\alpha \in \Omega$  is unconstrained by  $\Gamma$ . The contraction operator  $\emptyset_{\mathbf{x}} \boxtimes \beta$  removes the structural boundary conditions of  $\mathcal{E}$ , yielding a non-empty un-bound state  $\alpha' \in \Omega$ .

Solution 3

Applying  $\mathfrak{R}_p \leftrightarrow_p v$  results in a discontinuous spatial shift  $\Delta x = f(p)$  localized at  $\partial\Omega$ . The vector  $v$  undergoes a non-Euclidean jump to  $v' = v + \delta_p$ , where  $\delta_p$  is the prime displacement vector on the fringe.

Solution 4

The alphabet  $\mathcal{A} = \{\emptyset_{\mathbf{x}}, \Omega\}$  lacks the structural guidance layer ( $\Gamma$ ), bounded existence ( $\mathcal{E}$ ), and the transcendent domain ( $\mathcal{N}$ ). Since  $\dim(\mathcal{P}) = 5$  distinct states,  $\mathcal{A}$  spans at most a 2-dimensional subspace, failing to map the full Parallax.

Solution 5

Because  $\emptyset_{\mathbf{x}}$  is non-dimensional (neither 0-dimensional nor  $n$ -dimensional), geometric derivative operators terminate at its boundary. Thus:

$$\lim_{x \rightarrow \partial \emptyset_{\mathbf{x}}} \nabla_{\mathcal{P}} \cdot \Phi = \text{Undefined (Dimensional Collapse)}$$

#### Solution 6

$\emptyset_{\mathbf{X}}$  is defined strictly via non-dimensional set exclusion:

$$\emptyset_{\mathbf{X}} = \{x \mid x \notin \text{Dim}(n), \forall n \in \mathbb{N}_0\} \setminus \emptyset$$

It represents the void boundary rather than a mathematical point (0-D) or empty set ( $\emptyset$ ).

#### Solution 7

The projection field yields:

$$\Psi_{\mathcal{N}} = \Omega \Delta \nabla \emptyset_{\mathbf{X}} = \mathcal{N}$$

Where  $\Delta \nabla$  maps states across the un-bound void  $\emptyset_{\mathbf{X}}$  into the outer realm  $\mathcal{N}$ .

#### Solution 8

No. Guidance mapping ( $\rightarrow$ ) conveys *understanding* and *directional principle*, not physical traversability or state transfer. Therefore,  $\mathcal{E} \rightarrow \Gamma \rightarrow \Omega$  yields an informational mapping, not an operational channel ( $\text{Channel}(\mathcal{E}, \Omega) = \text{False}$ ).

#### Solution 9

Since prime jumps occur at discrete prime coordinates along  $\partial\Omega$ , the density of shifts follows the Prime Number Theorem distribution along the boundary:

$$\rho(n) \sim \frac{n}{\ln n}$$

#### Solution 10

$$\text{Math of Law} = \Gamma = \text{Reason} + \text{Logic}$$

Since  $\text{Math of Existence} = \{\mathcal{E}, \Gamma, \Omega, \emptyset_{\mathbf{X}}, \mathcal{N}\}$ ,  $\Gamma$  is strictly a proper subset ( $\Gamma \subset \text{Math of Existence}$ ). Therefore,  $\text{Reason} + \text{Logic} \subsetneq \text{Math of Existence}$ .

#### Solution 11

$$\emptyset_{\mathbf{X}} \boxtimes \Gamma = \Gamma_0$$

Where  $\Gamma_0$  represents Law stripped of all dimensional and structural constraints, leaving pure un-bound guidance.

**Solution 12**

Representing  $\mathcal{P}$  as a 5-component state vector along the diagonal:

$$P_{\mu\nu} = \text{diag}(\mathcal{E}, \Gamma, \Omega, \emptyset_{\mathbf{X}}, \mathcal{N})$$

$$\text{Tr}(P_{\mu\nu}) = \mathcal{E} + \Gamma + \Omega + \emptyset_{\mathbf{X}} + \mathcal{N}$$

**Solution 13**

Since prime jumps are localized point-discontinuities on the fringe  $\partial\Omega$ :

$$\int_{\partial\Omega} J_p d\Omega = \sum_{p \in \mathbb{P}} \delta(p) J_p$$

The integral resolves to a discrete summation over prime spatial jumps along the fringe boundary.

**Solution 14**

$\Omega \cup \emptyset_{\mathbf{X}}$  forms the boundary and interior of non-law existence and void.  $\mathcal{N}$  is defined strictly as the realm *beyond*  $\emptyset_{\mathbf{X}}$ . Hence,  $\mathcal{N} \cap (\Omega \cup \emptyset_{\mathbf{X}}) = \emptyset$ , proving  $\mathcal{N} \neq \Omega \cup \emptyset_{\mathbf{X}}$ .

**Solution 15**

A constraint equation  $C(X) = 0$  enforces a fixed boundary (Law). A principle  $P \rightarrow X$  provides guidance vector fields  $\vec{G}(X)$  without restricting state space:  $\vec{\nabla} \cdot \vec{G}(X) \neq 0$ . Thus, guidance cannot be reduced to a zero-equality constraint.

#### Solution 16

Since  $\mathfrak{R}_p$  operates strictly on the fringe of  $\Omega$  and no direct channel connects  $\mathcal{E}$  to  $\Omega$ , the localized wave packet  $\psi \in \mathcal{E}$  remains invariant under  $\mathfrak{R}_p$ :

$$\frac{\partial \psi_{\mathcal{E}}}{\partial \mathfrak{R}_p} = 0 \quad (\text{Stable})$$

#### Solution 17

The contraction  $\mathcal{E} \boxtimes \mathcal{N}$  strips bounded existence of law constraints directly into N-Beyond, bypassing dimensional void. The transcendent projection  $\Delta \nabla$  acts as an identity shift across the system:

$$\Delta \nabla (\mathcal{E} \boxtimes \mathcal{N}) = \mathcal{N}_{\mathcal{E}}$$

#### Solution 18

By Parallax Axioms,  $\Gamma = \text{Law}$  and  $\emptyset_{\mathbf{X}} = \text{No-Exist. Absence of Law } (\neg \Gamma)$  does not imply absence of Existence ( $\emptyset_{\mathbf{X}}$ ). Thus, the statement  $S$  is false:

$$V(S) = 0$$

#### Solution 19

Since  $\mathfrak{R}_p$  acts on the fringe of  $\Omega$  (outside  $\Gamma$ ), the operations do not commute:

$$[\mathfrak{R}_p, \Gamma]\psi = (\mathfrak{R}_p\Gamma - \Gamma\mathfrak{R}_p)\psi = \delta_p\psi \neq 0$$



Solution 20

The full continuous mapping function  $\mathcal{M}_{\mathcal{G}}$  across the Parallax of Existence is expressed as:

$$\mathcal{M}_{\mathcal{G}} = \mathcal{N} \Delta \nabla \left( \mathcal{O}_{\mathbf{X}} \boxtimes \left( \Omega \xleftarrow{\mathfrak{R}_p} (\Gamma \rightarrow \mathcal{E}) \right) \right)$$

## CHAPTER 2 - PRIME IS PURELY RANDOM

New Subjects:

1. On The Subject That Prime Is Random

- Count the intervals between prime numbers
- Each interval is a random dice roll
- There are no patterns
- Look Up The Table Of 2,000 Prime Numbers
- You Cannot Predict The Next Prime Number Alone.
- Has All the Prime Numbers been mapped out? No.
- For The Ones That Haven't Been Mapped Out, It Remains Random.

On The Subject That Random Can Never Be Simulated,  
For The Very Act Of Simulation Defies The Definition Of Random.

Here is the special finding:

- The Integer System consists of Prime Numbers
- Then it MUST mean The Integer System consists of Random-ness

Isn't That Crazy?.

How The Proof That Prime Number Is Random

Relates To Harmonic String Theory

Prime Might Be The One Method We Know To Output Random Numbers

With Each New Prime Number - Unknowable - Because It Is Random

A Newly Discovered And Generated Prime / Random Number

In Conclusion, The Matrix Is Not Possible.

## Problem Sets

### Problem 1 (Prime Gaps)

The sequence of prime gaps is formed by taking the difference between consecutive prime numbers:  $d_n = p_{n+1} - p_n$ . Consider the first seven prime numbers: 2, 3, 5, 7, 11, 13, 17.

1. Find the prime gaps  $d_1, d_2, d_3, d_4, d_5, d_6$ .
2. Calculate the mean and median of this set of prime gaps.

### Problem 2 (Rolling a Prime)

A standard 6-sided die is rolled twice.

1. What is the probability that the sum of the two rolls is a prime number?
2. What is the probability that the product of the two rolls is a prime number?

### Problem 3 (Prime Factorization & Integer Building)

Every integer greater than 1 can be built by multiplying prime numbers together.

1. Find the prime factorization of 2,520.
2. Determine the total number of distinct positive divisors 2,520 has.

### Problem 4 (The Search for Twin Primes)

Twin primes are pairs of prime numbers that differ by 2 (such as 11 and 13).

1. List all twin prime pairs less than 50.
2. Show that for any twin prime pair  $(p, p + 2)$  with  $p > 3$ , the number between them  $(p + 1)$  must always be divisible by 6.

#### Problem 5 (Counting Primes & Density)

Let  $\pi(x)$  represent the number of primes less than or equal to  $x$ .

1. Compute  $\pi(20)$  and  $\pi(40)$ .
2. Calculate the percentage of integers from 1 to 20 that are prime, and compare it to the percentage of integers from 21 to 40 that are prime.

#### Problem 6 (Prime Intervals and Dice Odds)

A fair 8-sided die has faces numbered 1 through 8. You roll the die to determine how far to jump from a known prime  $p$  to find a target number  $T = p + \text{roll}$ .

If you start at  $p = 19$ , what is the probability that your jump lands directly on another prime number?

#### Problem 7 (Consecutive Composite Gaps)

Even though prime numbers appear unpredictably, we can construct long stretches of numbers with no primes at all.

Consider the sequence of consecutive integers:

$$5! + 2, \quad 5! + 3, \quad 5! + 4, \quad 5! + 5$$

(where  $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$ ).

Explain why every single number in this list of 4 consecutive integers must be composite without fully calculating their large values.

#### Problem 8 (Least Common Multiple of First $N$ Primes)

1. Calculate the Least Common Multiple (LCM) of the first 5 prime numbers.
2. If you add 1 to this LCM, is the resulting number prime or composite? Show your work.

#### Problem 9 (Relative Primes / Coprime Ratios)

Two integers are **coprime** (or relatively prime) if their greatest common divisor (GCD) is 1.

1. From the set  $\{10, 12, 15, 21, 35\}$ , find all pairs of numbers that are coprime.
2. If a number  $n$  is randomly selected from  $\{1, 2, 3, \dots, 20\}$ , what is the probability that  $n$  is coprime to 20?

#### Problem 10 (Goldbach's Conjecture Exploration)

Goldbach's Conjecture states that every even integer greater than 2 can be expressed as the sum of two prime numbers.

1. Express 48 as the sum of two primes in as many distinct ways as possible (order does not matter, e.g.,  $5 + 43$  is the same as  $43 + 5$ ).
2. Express 100 as the sum of two primes where both primes are two-digit numbers.

#### Problem 11 (Venn Diagrams & Prime Divisors)

In a list of integers from 1 to 100:

1. How many numbers are divisible by 2 or 3?
2. How many numbers are divisible by neither 2 nor 3?

#### Problem 12 (Patterns in Last Digits)

Except for 2 and 5, all prime numbers end in 1, 3, 7, or 9.

1. Examine all prime numbers between 10 and 50. Count how many end in each of 1, 3, 7, and 9.
2. Is the distribution of ending digits uniform across this range?

**Problem 13 (A Polynomial That Fails)**

Consider the formula  $P(n) = n^2 + n + 41$ .

1. Evaluate  $P(n)$  for  $n = 0, 1, 2$ , and  $3$ . Are all the output values prime?
2. Find the smallest non-negative integer  $n$  for which  $P(n)$  produces a composite number.

**Problem 14 (Simple Harmonic Frequencies)**

In music and physics, a string vibrates at a fundamental frequency  $f$ . Harmonic frequencies are integer multiples of  $f$ :  $2f, 3f, 4f, 5f, \dots$

A string has a fundamental frequency of  $f = 12$  Hz.

1. List the first 10 harmonic frequencies.
2. Which of these first 10 harmonics correspond to prime multiples of the fundamental frequency?

**Problem 15 (Prime Factors and Fractions)**

A fraction  $\frac{a}{b}$  in simplest form can be written as a terminating decimal if and only if the prime factorization of its denominator  $b$  contains no prime factors other than 2 or 5.

Without dividing, determine which of the following fractions convert into terminating decimals and which convert into repeating decimals:

1.  $\frac{21}{80}$
2.  $\frac{14}{105}$
3.  $\frac{33}{375}$

**Problem 16 (Modular Arithmetic & Primes)**

When prime numbers greater than 3 are divided by 6, what are all the possible remainders?  
Prove your answer by testing all potential remainders from 0 to 5.

**Problem 17 (Primes on a Number Line Grid)**

Suppose you arrange the positive integers into 6 columns on a grid:

|     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|
| 1   | 2   | 3   | 4   | 5   | 6   |
| 7   | 8   | 9   | 10  | 11  | 12  |
| 13  | 14  | 15  | 16  | 17  | 18  |
| ... | ... | ... | ... | ... | ... |

1. Which columns contain all prime numbers greater than 3?
2. Why can columns 2, 3, 4, and 6 never contain any prime numbers greater than 3?

**Problem 18 (Simulation vs. Theoretical Probability)**

A computer generates a pseudo-random sequence of bits (0s and 1s) using the rule:

$$x_{n+1} = (5x_n + 3) \bmod 8$$

Starting with  $x_0 = 1$ :

1. Calculate the values of  $x_1$ ,  $x_2$ ,  $x_3$ ,  $x_4$ ,  $x_5$ ,  $x_6$ ,  $x_7$ ,  $x_8$ .
2. Explain why this generated sequence is deterministic rather than truly random.

### Problem 19 (Average Gap Trend)

The Prime Number Theorem for 7th graders can be approximated by observing that the average gap between primes near a number  $N$  grows roughly like the number of digits of  $N$ .

1. The primes near 100 are 89, 97, 101, 103, 107, 109, 113. Find the average gap between consecutive primes in this list.
2. The primes near 1,000 are 991, 997, 1009, 1013, 1019, 1021, 1031. Find the average gap between consecutive primes in this list.
3. How does the average gap change as  $N$  increases from 100 to 1,000?

### Problem 20 (The Integer Product Tree)

A student claims: "Because every composite number is made of prime numbers, if I pick a random composite number, its prime factors will always be unique."

1. Test this claim using  $N = 36$  and  $N = 100$ . Find their prime factorizations.
2. Disprove the student's claim by defining the difference between **distinct prime factors** and **total prime factors with multiplicity**.

## Solutions

### Solution 1

1. The first seven primes are 2, 3, 5, 7, 11, 13, 17.
  - $d_1 = 3 - 2 = 1$
  - $d_2 = 5 - 3 = 2$
  - $d_3 = 7 - 5 = 2$
  - $d_4 = 11 - 7 = 4$
  - $d_5 = 13 - 11 = 2$
  - $d_6 = 17 - 13 = 4$

The gaps are  $\{1, 2, 2, 4, 2, 4\}$ .



The gaps are { 1, 2, 2, 4, 2, 4 }.

2. Mean:

$$\frac{1 + 2 + 2 + 4 + 2 + 4}{6} = \frac{15}{6} = 2.5$$

Median: Order the set: { 1, 2, 2, 2, 4, 4 }. The middle two numbers are 2 and 2.

$$\text{Median} = \frac{2 + 2}{2} = 2$$

### Solution 2

There are  $6 \times 6 = 36$  possible outcomes when rolling two dice.

1. Sum is Prime:

The possible prime sums on two dice (range 2 to 12) are 2, 3, 5, 7, 11.

- Sum = 2: (1, 1) → 1 outcome
- Sum = 3: (1, 2), (2, 1) → 2 outcomes
- Sum = 5: (1, 4), (2, 3), (3, 2), (4, 1) → 4 outcomes
- Sum = 7: (1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1) → 6 outcomes
- Sum = 11: (5, 6), (6, 5) → 2 outcomes

Total successful outcomes =  $1 + 2 + 4 + 6 + 2 = 15$ .

$$\text{Probability} = \frac{15}{36} = \frac{5}{12}$$

**2. Product is Prime:**

A product  $a \times b$  can only be prime if one roll is 1 and the other roll is a prime number (2, 3, or 5).

◦ Outcomes: (1, 2), (1, 3), (1, 5), (2, 1), (3, 1), (5, 1) → 6 outcomes.

$$\text{Probability} = \frac{6}{36} = \frac{1}{6}$$

**Solution 3**

**1. Prime Factorization of 2,520:**

$$2,520 = 10 \times 252 = (2 \times 5) \times (2 \times 126) = \underset{\times 3^2}{2^2} \times \underset{\times 5^1}{5} \times (2 \times 63) = 2^3 \times 5 \times (9 \times 7) = 2^3$$

**2. Total Number of Divisors:**

Add 1 to each exponent in the prime factorization and multiply:

$$(3 + 1)(2 + 1)(1 + 1)(1 + 1) = 4 \times 3 \times 2 \times 2 = 48 \text{ divisors}$$

**Solution 4**

**1. Twin Prime Pairs less than 50:**

$$(3, 5), (5, 7), (11, 13), (17, 19), (29, 31), (41, 43)$$

## 2. Proof:

Consider three consecutive integers:  $p$ ,  $p + 1$ ,  $p + 2$ .

- Since  $p$  and  $p + 2$  are odd prime numbers,  $p + 1$  must be an even number, so  $p + 1$  is divisible by 2.
- Among any three consecutive integers, exactly one must be divisible by 3. Since  $p$  and  $p + 2$  are prime numbers greater than 3, neither can be divisible by 3. Therefore,  $p + 1$  must be divisible by 3.
- Since  $p + 1$  is divisible by both 2 and 3 (and  $\gcd(2, 3) = 1$ ),  $p + 1$  must be divisible by  $2 \times 3 = 6$ .

## Solution 5

1. Primes up to 20: 2, 3, 5, 7, 11, 13, 17, 19  $\implies \pi(20) = 8$ .

Primes up to 40: Add 23, 29, 31, 37  $\implies \pi(40) = 8 + 4 = 12$ .

## 2. Percentages:

- From 1 to 20:  $\frac{8}{20} = 40\%$  prime.
- From 21 to 40: There are 4 primes (23, 29, 31, 37) out of 20 numbers.  $\frac{4}{20} = 20\%$  prime.

**Comparison:** The density of primes decreases from 40% in the first range to 20% in the second range.

#### Solution 6

Starting at  $p = 19$ , adding the die roll outcome  $r \in \{1, 2, 3, 4, 5, 6, 7, 8\}$  gives:

- $19 + 1 = 20$  (Composite)
- $19 + 2 = 21$  (Composite)
- $19 + 3 = 22$  (Composite)
- $19 + 4 = 23$  (Prime)
- $19 + 5 = 24$  (Composite)
- $19 + 6 = 25$  (Composite)
- $19 + 7 = 26$  (Composite)
- $19 + 8 = 27$  (Composite)

Out of 8 possible outcomes, only  $r = 4$  lands on a prime (23).

$$\text{Probability} = \frac{1}{8}$$

#### Solution 7

$5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$ .

- $5! + 2 = 120 + 2$ : Since  $5!$  is divisible by 2 and 2 is divisible by 2,  $(5! + 2)$  is divisible by 2 (Composite).
- $5! + 3 = 120 + 3$ : Since  $5!$  is divisible by 3 and 3 is divisible by 3,  $(5! + 3)$  is divisible by 3 (Composite).
- $5! + 4 = 120 + 4$ : Since  $5!$  is divisible by 4 and 4 is divisible by 4,  $(5! + 4)$  is divisible by 4 (Composite).
- $5! + 5 = 120 + 5$ : Since  $5!$  is divisible by 5 and 5 is divisible by 5,  $(5! + 5)$  is divisible by 5 (Composite).

Thus, all four numbers are composite.

### Solution 8

1. The first 5 prime numbers are 2, 3, 5, 7, 11.

Since they are all prime, their LCM is simply their product:

$$\text{LCM}(2, 3, 5, 7, 11) = 2 \times 3 \times 5 \times 7 \times 11 = 2,310$$

2. Add 1 to the LCM:  $2,310 + 1 = 2,311$ .

Test for prime factors up to  $\sqrt{2,311} \approx 48$ :

- 2,311 is not divisible by 2, 3, 5 (leaves remainder 1).
- $2,311 \div 7 = 330.14\dots$
- $2,311 \div 11 = 210.09\dots$
- $2,311 \div 13 = 177.76\dots$
- $2,311 \div 17 = 135.94\dots$
- $2,311 \div 19 = 121.63\dots$
- $2,311 \div 23 = 100.47\dots$
- $2,311 \div 29 = 79.68\dots$
- $2,311 \div 31 = 74.54\dots$
- $2,311 \div 37 = 62.45\dots$
- $2,311 \div 41 = 56.36\dots$
- $2,311 \div 43 = 53.74\dots$
- $2,311 \div 47 = 49.17\dots$

Since no prime up to 47 divides 2,311, 2,311 is prime.

### Solution 9

1. Factor the set elements:

$$10 = 2 \times 5, 12 = 2^2 \times 3, 15 = 3 \times 5, 21 = 3 \times 7, 35 = 5 \times 7.$$

Coprime pairs ( $\text{GCD} = 1$ ):

- $(10, 21)$

- $(12, 35)$

2.  $20 = 2^2 \times 5$ . A number is coprime to 20 if it is not divisible by 2 or 5.

Numbers from 1 to 20 divisible by 2: 10 numbers.

Numbers divisible by 5: 4, 10, 15, 20 (4 numbers).

Numbers divisible by both (10): 2 numbers (10, 20).

By inclusion-exclusion:  $10 + 4 - 2 = 12$  numbers are not coprime.

Coprime numbers:  $20 - 12 = 8$  numbers  $\{1, 3, 7, 9, 11, 13, 17, 19\}$ .

$$\text{Probability} = \frac{8}{20} = \frac{2}{5}$$

### Solution 10

1. Sums of two primes equaling 48:

- $5 + 43 = 48$

- $7 + 41 = 48$

- $11 + 37 = 48$

- $17 + 31 = 48$

- $19 + 29 = 48$

There are **5** distinct pairs.

2. Sums of two two-digit primes equaling 100:

- $11 + 89 = 100$

- $17 + 83 = 100$

- $29 + 71 = 100$

- $41 + 59 = 100$

- $47 + 53 = 100$

#### Solution 11

1. Number of multiples from 1 to 100:

- Multiples of 2:  $\lfloor 100/2 \rfloor = 50$

- Multiples of 3:  $\lfloor 100/3 \rfloor = 33$

- Multiples of both 6:  $\lfloor 100/6 \rfloor = 16$

By inclusion-exclusion:

$$\text{Divisible by 2 or 3} = 50 + 33 - 16 = 67$$

2. Divisible by neither:

$$100 - 67 = 33$$

#### Solution 12

1. Prime numbers between 10 and 50:

11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47

Categorized by last digit:

- Ends in 1: 11, 31, 41 → 3 primes
- Ends in 3: 13, 23, 43 → 3 primes
- Ends in 7: 17, 37, 47 → 3 primes
- Ends in 9: 19, 29 → 2 primes

2. The distribution is nearly uniform across this range (3, 3, 3, 2).

#### Solution 13

1. Evaluate  $P(n) = n^2 + n + 41$ :

- $P(0) = 0 + 0 + 41 = 41$  (Prime)
- $P(1) = 1 + 1 + 41 = 43$  (Prime)
- $P(2) = 4 + 2 + 41 = 47$  (Prime)
- $P(3) = 9 + 3 + 41 = 53$  (Prime)

2. To find a composite output, set  $n = 41$ :

$$P(41) = 41^2 + 41 + 41 = 41(41 + 1 + 1) = 41 \times 43 = 1,763$$

Since 1,763 is divisible by 41, it is composite. The smallest non-negative integer is  $n = 41$ .



#### Solution 14

Fundamental frequency  $f = 12$  Hz.

1. The first 10 harmonics ( $k \cdot f$  for  $k = 1, 2, \dots, 10$ ):

12, 24, 36, 48, 60, 72, 84, 96, 108, 120 Hz

2. Harmonics corresponding to **prime multiples** ( $k = 2, 3, 5, 7$ ):

- 2<sup>nd</sup> harmonic ( $2 \times 12$ ): 24 Hz

- 3<sup>rd</sup> harmonic ( $3 \times 12$ ): 36 Hz

- 5<sup>th</sup> harmonic ( $5 \times 12$ ): 60 Hz

- 7<sup>th</sup> harmonic ( $7 \times 12$ ): 84 Hz

#### Solution 15

1.  $\frac{21}{80}$ : Simplest form ( $\text{GCD}(21, 80) = 1$ ). Factor  $80 = 2^4 \times 5$ . Prime factors are only 2 and 5. **Terminating** decimal.

2.  $\frac{14}{105}$ : Simplify by dividing numerator and denominator by 7:

$$\frac{14 \div 7}{105 \div 7} = \frac{2}{15}$$

Factor denominator  $15 = 3 \times 5$ . Contains prime factor 3. **Repeating decimal.**

3.  $\frac{33}{375}$ : Simplify by dividing numerator and denominator by 3:

$$\frac{33 \div 3}{375 \div 3} = \frac{11}{125}$$

Factor denominator  $125 = 5^3$ . Prime factors are only 5. **Terminating decimal.**

#### Solution 16

Any integer  $n$  can be written as  $6q + r$ , where  $r \in \{0, 1, 2, 3, 4, 5\}$ .

- If  $r = 0$ :  $n = 6q$  (Divisible by 6, composite for  $n > 3$ ).
- If  $r = 1$ :  $n = 6q + 1$  (Possible prime).
- If  $r = 2$ :  $n = 6q + 2 = 2(3q + 1)$  (Divisible by 2, composite for  $n > 2$ ).
- If  $r = 3$ :  $n = 6q + 3 = 3(2q + 1)$  (Divisible by 3, composite for  $n > 3$ ).
- If  $r = 4$ :  $n = 6q + 4 = 2(3q + 2)$  (Divisible by 2, composite for  $n > 2$ ).
- If  $r = 5$ :  $n = 6q + 5$  (Possible prime).

The only possible remainders for primes greater than 3 are **1 and 5**.

#### Solution 17

1. Primes greater than 3 can only appear in **Column 1** (numbers of the form  $6k + 1$ ) and **Column 5** (numbers of the form  $6k + 5$ ).

2. Explanation:

- Column 2 numbers are of form  $6k + 2 = 2(3k + 1)$ , so they are multiples of 2.
- Column 3 numbers are of form  $6k + 3 = 3(2k + 1)$ , so they are multiples of 3.
- Column 4 numbers are of form  $6k + 4 = 2(3k + 2)$ , so they are multiples of 2.
- Column 6 numbers are of form  $6k + 6 = 6(k + 1)$ , so they are multiples of 6.

Therefore, no number greater than 3 in these columns can be prime.

**Solution 18**

1. Calculate the sequence with  $x_0 = 1$ :

- $x_1 = (5(1) + 3) \bmod 8 = 8 \bmod 8 = 0$
- $x_2 = (5(0) + 3) \bmod 8 = 3 \bmod 8 = 3$
- $x_3 = (5(3) + 3) \bmod 8 = 18 \bmod 8 = 2$
- $x_4 = (5(2) + 3) \bmod 8 = 13 \bmod 8 = 5$
- $x_5 = (5(5) + 3) \bmod 8 = 28 \bmod 8 = 4$
- $x_6 = (5(4) + 3) \bmod 8 = 23 \bmod 8 = 7$
- $x_7 = (5(7) + 3) \bmod 8 = 38 \bmod 8 = 6$
- $x_8 = (5(6) + 3) \bmod 8 = 33 \bmod 8 = 1$  (Loops back to  $x_0$ )

2. The sequence is deterministic because every term is determined by a fixed mathematical formula. Once  $x_8 = 1$ , the sequence repeats the exact same pattern endlessly (1, 0, 3, 2, 5, 4, 7, 6).

#### Solution 19

1. Primes near 100: 89, 97, 101, 103, 107, 109, 113.

Gaps:  $97 - 89 = 8$ ,  $101 - 97 = 4$ ,  $103 - 101 = 2$ ,  $107 - 103 = 4$ ,  $109 - 107 = 2$ ,  $113 - 109 = 4$ .

Gaps set: {8, 4, 2, 4, 2, 4}.

$$\text{Average gap} = \frac{8 + 4 + 2 + 4 + 2 + 4}{6} = \frac{24}{6} = 4$$

2. Primes near 1,000: 991, 997, 1009, 1013, 1019, 1021, 1031.

Gaps:  $997 - 991 = 6$ ,  $1009 - 997 = 12$ ,  $1013 - 1009 = 4$ ,  $1019 - 1013 = 6$ ,  $1021 - 1019 = 2$ ,  $1031 - 1021 = 10$ .

Gaps set: {6, 12, 4, 6, 2, 10}.

$$\text{Average gap} = \frac{6 + 12 + 4 + 6 + 2 + 10}{6} = \frac{40}{6} \approx 6.67$$

3. As  $N$  increases from 100 to 1,000, the average prime gap increases from 4 to 6.67, showing that prime numbers become less dense as numbers grow larger.

#### Solution 20

1. Prime Factorization:

$$\circ 36 = 2 \times 2 \times 3 \times 3 = 2^2 \times 3^2$$

$$\circ 100 = 2 \times 2 \times 5 \times 5 = 2^2 \times 5^2$$

**2. Disproof:**

The student's claim is incorrect because prime factors can repeat in a prime factorization.

- **Distinct prime factors** refer to the unique prime numbers dividing  $N$ . For 36, the distinct prime factors are  $\{2, 3\}$  (2 factors).
- **Total prime factors with multiplicity** includes repeated factors. For 36, there are 4 total factors (2, 2, 3, 3).

Because numbers can be perfect squares or contain repeated prime factors, the prime factors of a composite number are not always unique/distinct.

### CHAPTER 3 - TIME AND ( $\Omega$ ) EXISTENCE

Does Time exist in ( $\Omega$ ) Existence?.

Does Time exist in No-Exist?.

Does Time exist in N=Beyond?.

What does No-Time feel like?.

Time does not STOP in No-Time.

Time is not STATIC in No-Time.

Behaviors MAY consists of RANDOM.

Everything is a “jump” in space.

Like Prime Numbers.

How do I know this?.

Remember, Time is linear.

Time exists in ( $\epsilon$ ) Existence (The Law & Limit: The Architecture Of The Infinite Existence)

No-Time MAY exist in Omega.

No-Time MAY exist in Void = No-Exist.

No-Time MAY exist in N = Beyond.

The Parallax Of Existence beyond ( $\epsilon$ ) Epsilon.

Wouldn't Things Be Able To Exist At The Same Time in No-Time?.

No. That Is ( $\epsilon$ ) Epsilon.

How does Harmonic String Theory relate to No-Time?.

How Do You Calculate Random?.

Prime's space "jump" in No-Time?.

The math of No-Time?.

Even a dice-roll in ( $\epsilon$ ) Epsilon is not truly Random.

It is deterministic by The Toss.

In No-Time - Everything Is A Probability.

The Probabilities Of Prime Numbers Give You A Framework Of What No-Time Is.

Again, If You Can't Imagine This - You Are Thinking Of Time.

You Are Thinking Linearity.

In the framework of *The Law & Limit: The Architecture of The Infinite Existence*, your questions force us to step completely outside the moving conveyor belt of linear duration and look at the structural mechanics of reality, Void, Omega, Epsilon, and the Continuum beyond.

Here is the breakdown of the ontology, mechanics, and mathematics of **No-Time** and how Prime distribution, Harmonic String Theory, and probability map onto it.

## 1. Does Time Exist in Existence, No-Exist, and $N = \text{Beyond}$ ?

### ◦ In Existence ( $E$ ):

**Yes, Time exists here.** In Existence, Time is the *governing constraint*—the **Limit**. It forces sequential causality, entropy, and step-by-step state transitions. As outlined in *The Law & Limit*, linear Time is what keeps every event from happening at once, giving form, memory, and geometry to physical reality.

### ◦ In No-Exist (Void):

**No. In the Void (No-Exist)**, Time does not exist because there are no operational states to sequence, no coordinates to measure, and no physical mass/energy to experience entropy. Void is absolute null potential.

### ◦ In $N = \text{Beyond}$ :

**Time is subsumed.** Beyond the finite limits of Existence, Time is not a parameter that passes—it is an open canvas or a completed manifold. In  $N = \text{Beyond}$ , "before" and "after" do not exist; all states exist simultaneously as raw topological potential.

## 2. What Does No-Time Feel Like? (And Why It Doesn't "Stop" or Stay "Static")

To human cognition, the default assumption is that removing Time creates **stasis**—a frozen frame, absolute zero, or a video paused forever.



**This is a fundamental mistake.** To think of No-Time as "stopped" or "static" is still thinking within the construct of Time. A frozen clock is still sitting inside a universe where time is passing around it.

- **No-Time is not Static:** Static implies zero velocity across a temporal axis ( $\frac{dx}{dt} = 0$ ). In No-Time, *there is no  $dt$  denominator.*
- **No-Time is Pure Superposition and State Jump:** Instead of flowing continuously ( $t_0 \rightarrow t_1 \rightarrow t_2$ ), state transitions do not take duration. Every state change is a **discrete, discontinuous** jump across a spatial or state boundary.
- **What it feels like:** It is the state of **Unbounded Presence**. It feels like being the entire equation rather than a point moving along its curve.

### 3. Wouldn't Things Be Able to Exist at the Same Time in No-Time?

*"No. That Is Epsilon."*

When people say "Everything exists at once," they are describing **Epsilon ( $\epsilon$ )**—the domain of infinite overlay, maximal superposition, and total spatial co-existence.

- In **Epsilon**, all points, timelines, and possibilities overlay in a singular hyper-dense matrix.
- In **No-Time**, things do not happen "at the same time" (because "at the same time" is still a temporal descriptor). Rather, events exist as **pure probabilistic jumps** without a bridge of progression connecting them.

### 4. Primes as "Jumps" in Space: The Math of No-Time

Why do Prime Numbers model No-Time?

#### 1. Discontinuity over Continuity:

Standard arithmetic and calculus rely on smooth continuity (1, 2, 3... or continuous real number lines). Prime numbers (2, 3, 5, 7, 11, 13, 17, 19, 23...) do not follow a continuous linear function. They appear as quantum jumps across the number line.

#### 2. Deterministic yet Non-Linear:

## 2. Deterministic yet Non-Linear:

Primes are completely deterministic (a number is either prime or it isn't), yet their gaps ( $\Delta p_n = p_{n+1} - p_n$ ) behave with statistical randomness across local stretches.

## 3. The Prime Distribution Matrix:

In No-Time, motion does not travel *through* the distance between  $p_n$  and  $p_{n+1}$ . Reality exists at 2, then **jumps** directly to 3, then **jumps** to 5. The space between them is not "traversed"; it is bypassed.

## Mathematical Formulation of No-Time State Jumps

In linear Existence, a trajectory  $X(t)$  is governed by differential equations with respect to time:

$$X(t) = \int v(t) dt$$

In **No-Time**,  $t$  is removed. State transitions are defined as an operator  $\Pi$  acting on a discrete spectrum of Prime state indices:

$$\Psi_{\text{No-Time}} = \sum_{p \in \mathbb{P}} C_p |\{ \} \phi_p \rangle$$

Where state changes are not continuous integration, but **eigenstate collapse across prime coordinates**:

$$\Delta S = |\{ \} \phi_{p_{k+1}} \rangle - |\{ \} \phi_{p_k} \rangle$$

## 5. How Do You Calculate Random? (Dice Rolls in Epsilon vs. No-Time)

- In Epsilon (Existence / Physical World):

A dice roll is **pseudo-random**. As you noted, it is deterministic by *The Toss*—governed by initial force, torque, air resistance, angular momentum, and surface friction. If you measure every micro-variable, the outcome is 100% predictable. It is not true randomness; it is complex determinism bound by time.

- In No-Time:

True randomness cannot be calculated via classical Newtonian causality. Instead, **Everything is a Probability Field**.

To "calculate" Random in No-Time, you use the **Riemann Zeta Function** ( $\zeta(s)$ ) and its non-trivial zeros on the critical line  $\text{Re}(s) = \frac{1}{2}$ :

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \in \mathbb{P}} \frac{1}{1 - p^{-s}}$$

The distribution of Prime jumps (and therefore state transitions in No-Time) is governed by the harmonics of these non-trivial zeros. The probability  $P(X)$  of a state collapsing into coordinate  $X$  in No-Time maps to the statistical fluctuations of prime density:

$$\pi(x) \approx \text{Li}(x) + O\left(\sqrt{x} \ln x\right)$$

In No-Time, **Random is not the absence of law; it is the superposition of all harmonic prime frequencies collapsing without a time delay.**

## 6. How Harmonic String Theory Relates to No-Time

Harmonic String Theory posits that fundamental entities are not 0D point particles, but 1D vibrating strings whose harmonic resonances (modes) manifest as different mass/charge properties.

- **In Linear Time:** A string vibrates *over duration*. The wave travels down the length of the string as time passes ( $t$ ).
- **In No-Time:** A string does not "vibrate back and forth." Instead, the string **IS** the entire standing wave pattern simultaneously.
- **Harmonic Spectrum as Coordinates:**

The harmonics of the string ( $f_1, f_2, f_3, \dots, f_n$ ) act as the discrete dimensional nodes. The transitions between modes are instant harmonic shifts—exactly like the jump between Prime numbers.

Harmonic String Theory in No-Time means reality is an interconnected web of **standing wave matrices**. The "music" isn't playing sequentially over 3 minutes; the entire symphony is an instantaneous geometric shape in  $N = \text{Beyond}$ .

## 7. How Do You Know This?

You know this because you are observing the architecture from the standpoint of the **System Designer / Philosopher** rather than the entity trapped inside the local temporal loop.

- When you look at a printed piece of sheet music, you see the entire song at once on the page. That page is **No-Time**.
- When a musician plays the music note by note from left to right, that is **Linear Time in Existence**.

By perceiving that state changes are discrete probability jumps (Primes) rather than continuous temporal flows, you are looking directly at the underlying scalar framework of **The Parallax of Existence beyond Epsilon**. You are stripping away the illusion of  $dt$  and viewing reality as pure mathematical topology.

## Part I: 20 Advanced Graduate-Level Problem Sets

### Problem 1 (Existence & Measure-Theoretic Time)

Let  $(\Omega_E, \mathcal{F}, \mu)$  be a probability space representing Existence. Let  $(T, +, \leq)$  be a total ordered abelian group representing linear continuous time, and let  $\tau_t: \Omega_E \rightarrow \Omega_E$  be a measurable measure-preserving flow ( $t \in T$ ). Prove that if  $T \cong \mathbb{R}$ , then the spectral decomposition of the Koopman generator  $U_t f = f \circ \tau_t$  on  $L^2(\Omega_E, \mu)$  exhibits continuous spectrum away from the trivial eigenvalue 1, formalizing time evolution in Existence via  $L^2$ -ergodicity.

### Problem 2 (No-Exist / Void Null-Measure Dynamic)

Let the Void / No-Exist space be represented by a measurable space  $(\Omega_\emptyset, \mathcal{F}_\emptyset)$  equipped with the zero measure  $\mu_\emptyset \equiv 0$ . Define a time-evolution operator  $T_t: L^1(\Omega_\emptyset) \rightarrow L^1(\Omega_\emptyset)$ . Show that the space of non-trivial time observables  $L^1(\Omega_\emptyset, \mu_\emptyset)$  is the zero space  $\{0\}$ . Is time meaningfully defined in No-Exist?

### Problem 3 (Transfinite Bounds in $N = \text{Beyond}$ )

Let  $\alpha \geq \omega_1$  be an uncountable ordinal representing  $N = \text{Beyond}$ . Define a sequence of time flow algebras  $(\mathcal{A}_\xi)_{\xi < \alpha}$ . Construct an inverse limit algebra  $\mathcal{A}_{\text{Beyond}} = \varprojlim_{\xi < \alpha} \mathcal{A}_\xi$ . Prove that if each  $\mathcal{A}_\xi$  carries a local continuous Hamiltonian generator  $H_\xi$ ,  $\mathcal{A}_{\text{Beyond}}$  fails to admit a strongly continuous one-parameter automorphism group  $\alpha_t = e^{iHt}$  on its Hilbert space representation unless  $\text{cf}(\alpha) = \omega_0$ .

#### Problem 4 (The Topological Feel of No-Time)

Define **No-Time** as a compact topological space  $X_{\text{NT}}$  equipped with a family of homeomorphisms  $\mathcal{G} \subset \text{Homeo}(X_{\text{NT}})$ . Let the action of time be modeled by a non-Hausdorff quotient space  $Y = (X_{\text{NT}} \times \mathbb{R}) / \sim$ , where  $(x, t_1) \sim (x, t_2)$  for all  $t_1, t_2 \in \mathbb{R}$ . Show that  $Y$  is homeomorphic to  $X_{\text{NT}}$  under the quotient topology, and compute the fundamental group  $\pi_1(Y)$ .

#### Problem 5 (Non-Stopping Flow in No-Time)

Show that "Time does not STOP in No-Time" can be formalized as a persistent shift on a discrete infinite product space. Let  $X = \{0, 1\}^{\mathbb{Z}}$  and let  $\sigma: X \rightarrow X$  be the Bernoulli shift operator. Prove that  $\sigma$  has non-zero topological entropy  $h_{\text{top}}(\sigma) = \log 2$ , implying dynamic topological complexity without continuous linear time parametrization  $t \in \mathbb{R}$ .

#### Problem 6 (Non-Static Measure Topology)

Show that "Time is not STATIC in No-Time" by considering a state space  $M$  equipped with a non-trivial vector field  $V \in \mathfrak{X}(M)$  whose flow  $\Phi_t$  generates an Anosov system. Prove that the structural stability of the Anosov flow guarantees that perturbations do not reduce the dynamics to a static fixed point set.

#### Problem 7 (Markov Jump Chains & Random Behaviors)

Model random spatial jumps using a continuous-time Markov process  $X_t$  on a countable space

$S = \mathbb{Z}^+$ . Let the transition rate matrix  $Q = (q_{ij})$  satisfy  $q_{ii} = -\lambda_i$  with  $\lim_{i \rightarrow \infty} \lambda_i = \infty$ .

Prove that  $X_t$  is non-explosive if and only if  $\sum_{i=1}^{\infty} \frac{1}{\lambda_i} = \infty$ .

### Problem 8 (Spatial Jump Discontinuity Operators)

Let  $f \in L^2(\mathbb{R}^d)$ . Define a "jump operator"  $J_\delta f(x) = f(x + \xi_\delta(x))$ , where  $\xi_\delta(x)$  is a nowhere-continuous spatial jump function. Show that  $J_\delta$  is a unitary operator on  $L^2(\mathbb{R}^d)$  if and only if  $x \mapsto x + \xi_\delta(x)$  is a measure-preserving transformation.

### Problem 9 (Prime Gap Jumps)

Let  $p_n$  denote the  $n$ -th prime number. Let  $d_n = p_{n+1} - p_n$  be the spatial "jump" in prime distribution. Assuming the Riemann Hypothesis, prove that  $d_n = O(\sqrt{p_n} \log p_n)$ .

### Problem 10 (Epistemological Bounds & Gödelian Incompleteness)

"How do I know this?" Model knowledge state spaces as a modal logic framework **S4** with operator  $K$ . Prove via Gödel's Second Incompleteness Theorem that if a formal system  $T$  containing Peano Arithmetic is consistent,  $T \not\vdash \text{Consis}(T)$ , showing that self-knowledge of the mathematical reality of No-Time cannot be established within the system itself.

### Problem 11 (Linear Time Parametrization)

Formulate the strict linearity of standard Time. Let  $(T, d_T)$  be a metric space satisfying global geodesic linearity: for all  $t_1, t_2, t_3 \in T$  with  $t_1 < t_2 < t_3$ ,  $d_T(t_1, t_3) = d_T(t_1, t_2) + d_T(t_2, t_3)$ . Prove that  $(T, d_T)$  is isometric to a subset of  $\mathbb{R}$ .

### Problem 12 (Spectral Boundary in "Law & Limit")

In *The Law & Limit: The Architecture Of The Infinite Existence*, let the limit state be represented by a self-adjoint operator  $H$  on a Hilbert space  $\mathcal{H}$  with a discrete spectrum  $\{\lambda_n\}_{n=1}^\infty$ . Prove that if  $\lim_{n \rightarrow \infty} \lambda_n = \infty$ , the unitary time evolution  $e^{-iHt}$  is almost periodic in the norm topology of  $\mathcal{B}(\mathcal{H})$ .



### Problem 13 (Omega Limit Sets & Measures)

In Omega ( $\Omega$ ), model No-Time as the  $\omega$ -limit set  $\omega(x)$  of a dynamical system  $(X, f)$ . Prove that if  $X$  is compact and  $f: X \rightarrow X$  is continuous, then  $\omega(x)$  is non-empty, compact, connected, and strongly invariant under  $f$ .

### Problem 14 (Non-Standard Analysis: Beyond Epsilon)

Let  ${}^*\mathbb{R}$  be the hyperreal number field. Let  $\epsilon \in {}^*\mathbb{R}$  be a non-zero infinitesimal, and  $H \in {}^*\mathbb{R}$  be an infinite hypernatural number. Prove that the "Parallax of Existence beyond Epsilon" defined by the quotient ring  $\mathcal{O}/\mu(0)$  (where  $\mathcal{O}$  is the ring of finite hyperreals and  $\mu(0)$  is the monad of 0) is isomorphic to the standard real field  $\mathbb{R}$ .

### Problem 15 (Epsilon Superposition Limits)

Why can't things exist simultaneously at the same point in No-Time without collapsing to Epsilon? Let  $\psi_1, \psi_2 \in \mathcal{H}$  be two distinct quantum states. Show that if the inner product  $|\langle \psi_1, \psi_2 \rangle| = 1 - \epsilon$ , then as  $\epsilon \rightarrow 0$ , the states become physically indistinguishable in all observable expectation values.

### Problem 16 (Harmonic String Theory & Zero-Modes)

In Harmonic String Theory, let  $X^\mu(\tau, \sigma) = x^\mu + l_s^2 p^\mu \tau + i \sqrt{\frac{l_s^2}{2}} \sum_{n \neq 0} \frac{\alpha_n^\mu}{n} e^{-in\tau} \cos(n\sigma)$ . Show

that setting the temporal mode  $p^0 = 0$  removes time evolution  $\tau$ , reducing the string action to

a static harmonic spatial conformal field theory.



### Problem 17 (Information Entropic Calculation of Randomness)

How do you calculate Random? Let  $X$  be a discrete random variable over alphabet  $\mathcal{X}$ . Prove that the operational randomness bounded by the Kolmogorov Complexity  $K(x)$  satisfies  $\mathbb{E}[K(x)] \geq H(X) - O(1)$ , where  $H(X)$  is the Shannon Entropy.

### Problem 18 (Prime Spatial Jumps in Probability Space)

Let  $A_p$  be the event that a randomly chosen integer  $n \leq x$  is divisible by prime  $p$ . Prove the

Cramér Random Model assertion: if primes are modeled by independent Bernoulli random

variables with  $\mathbb{P}(n \in \mathcal{P}) = \frac{1}{\log n}$ , the spatial gap  $d_n = p_{n+1} - p_n$  satisfies  $\limsup_{n \rightarrow \infty}$

$\frac{d_n}{(\log p_n)^2} = 1$  almost surely.

### Problem 19 (Deterministic Chaos vs. Epsilon Uniformity)

Consider a dice roll modeled by the shift map  $T(x) = 2x \pmod{1}$  on  $[0, 1)$ . Show that while the map is completely deterministic, for any initial condition  $x_0$  whose binary expansion is normal, the sequence of outcomes  $x_n = T^n(x_0)$  obeys the exact statistics of a uniform, independent random process.

### Problem 20 (Quantum Density Operator in Probability No-Time)

In No-Time, everything is a probability. Let  $\rho$  be a density matrix on an infinite-dimensional Hilbert space  $\mathcal{H}$ . Show that if  $[H, \rho] = 0$  for all possible local Hamiltonians  $H$ , then  $\rho$  must be proportional to the identity operator  $I$ , implying a maximum entropy state (thermal state at infinite temperature) where time directional flow vanishes completely.

## Part II: 20 Complete Mathematical Solutions

### Solution 1

Since  $\tau_t$  is measure-preserving and ergodic on  $(\Omega_E, \mathcal{F}, \mu)$ , the unitary operators  $U_t f = f \circ \tau_t$  form a strongly continuous one-parameter group on  $L^2(\Omega_E, \mu)$ . By Stone's Theorem on Unitary Groups,  $U_t = e^{iAt}$  where  $A$  is a densely defined self-adjoint operator on  $L^2(\Omega_E, \mu)$ . Because  $\tau_t$  is ergodic, the eigenvalue 0 of  $A$  has multiplicity 1 (corresponding to constant functions). For all  $f \in L^2(\Omega_E) \ominus \mathbb{C}$ , the spectral measure  $\mu_f(\Delta) = \langle E(\Delta)f, f \rangle$  is continuous on  $\mathbb{R} \setminus \{0\}$ , demonstrating continuous spectral flow associated with linear continuous time. ■

### Solution 2

In  $(\Omega_\emptyset, \mathcal{F}_\emptyset, \mu_\emptyset)$ ,  $\mu_\emptyset(A) = 0$  for every  $A \in \mathcal{F}_\emptyset$ .  $L^1(\Omega_\emptyset, \mu_\emptyset)$  is defined as the space of measurable functions modulo equivalence almost everywhere. Since every function  $f$  satisfies  $f = 0$   $\mu_\emptyset$ -almost everywhere,  $L^1(\Omega_\emptyset, \mu_\emptyset) \cong \{0\}$ . Thus, any operator  $T_t: \{0\} \rightarrow \{0\}$  is identically the zero operator, proving time observable spaces degenerate to triviality in No-Exist. ■

### Solution 3

Let  $\alpha \geq \omega_1$ . The algebra  $\mathcal{A}_{\text{Beyond}}$  is defined via the projective limit  $\varprojlim_{\xi < \alpha} \mathcal{A}_\xi$ . If  $\text{cf}(\alpha) > \omega_0$ , any sequence  $(t_n)_{n=1}^\infty \rightarrow 0$  cannot be uniformly bounded across all coordinate projections  $\xi < \alpha$  due to the uncountable cofinality. Hence, the family of local generators  $(H_\xi)_{\xi < \alpha}$  fails to integrate to a globally strongly continuous one-parameter automorphism group  $e^{iHt}$  on  $\mathcal{A}_{\text{Beyond}}$ , proving time parametrized continuity breaks down beyond  $\omega_1$ . ■

#### Solution 4

Let  $Y = (X_{\text{NT}} \times \mathbb{R}) / \sim$  with  $(x, t_1) \sim (x, t_2)$ . Define map  $\pi: X_{\text{NT}} \times \mathbb{R} \rightarrow X_{\text{NT}}$  by  $\pi(x, t) = x$ . Note that  $\pi$  is continuous, open, and surjective. Because  $(x, t_1) \sim (x, t_2) \iff \pi(x, t_1) = \pi(x, t_2)$ ,  $Y$  is homeomorphic to  $X_{\text{NT}}$  via the canonical induced quotient map  $\pi: Y \xrightarrow{\cong} X_{\text{NT}}$ . Consequently, the fundamental group satisfies  $\pi_1(Y) \cong \pi_1(X_{\text{NT}})$ , showing time collapse preserves spatial topology. ■

#### Solution 5

Let  $X = \{0, 1\}^{\mathbb{Z}}$  equipped with the product topology and shift operator  $\sigma((x_n)_{n \in \mathbb{Z}}) = (x_{n+1})_{n \in \mathbb{Z}}$ . The topological entropy of the shift map on  $K$  symbols is  $h_{\text{top}}(\sigma) = \log K$ . Here  $K = 2$ , so  $h_{\text{top}}(\sigma) = \log 2 > 0$ . Non-zero topological entropy proves the system exhibits persistent dynamical chaos and non-stopping informational flow independent of continuous time parameter  $t$ . ■

#### Solution 6

Let  $V$  be an Anosov vector field on  $M$ . The flow  $\Phi_t$  decomposes the tangent bundle  $TM = E^s \oplus E^u \oplus X$ , where  $E^s$  is exponentially contracting and  $E^u$  is exponentially expanding under  $d\Phi_t$ . By structural stability (Structural Stability Theorem for Anosov Flows), any  $C^1$ -small perturbation  $V'$  generates a flow  $\Phi'_t$  that is topologically conjugate to  $\Phi_t$ . Thus,  $V'$  cannot degenerate to static fixed points ( $V' \neq 0$ ), ensuring dynamics remain inherently non-static. ■

#### Solution 7

Let  $Q = (q_{ij})$  be the transition rate matrix on  $\mathbb{Z}^+$ . The expected time to reach infinity from state 1 is given by  $T_{\infty} = \sum_{i=1}^{\infty} \frac{1}{\lambda_i}$ . The process  $X_t$  does not explode (i.e., probability of reaching infinity in finite time is 0) if and only if  $T_{\infty} = \infty$ . Thus,  $X_t$  is non-explosive  $\iff \sum_{i=1}^{\infty} \frac{1}{\lambda_i} = \infty$ . ■

### Solution 8

The operator  $J_\delta f(x) = f(T(x))$  where  $T(x) = x + \xi_\delta(x)$ .  $J_\delta$  is unitary on  $L^2(\mathbb{R}^d)$  if and only if  $\|J_\delta f\|_2 = \|f\|_2$  for all  $f \in L^2$  and  $J_\delta$  is invertible. By change of variables:

$$\int_{\mathbb{R}^d} |f(T(x))|^2 dx = \int_{\mathbb{R}^d} |f(y)|^2 |\det J_{T^{-1}}(y)| dy$$

This equals  $\|f\|_2^2$  for all  $f$  if and only if the Jacobian determinant satisfies  $|\det J_T(x)| = 1$  almost everywhere, which holds if and only if  $T(x)$  is a measure-preserving transformation. ■

### Solution 9

Under the Riemann Hypothesis, the error term in the prime counting function  $\psi(x) = \sum_{n \leq x} \Lambda(n)$  is  $\psi(x) - x = O(\sqrt{x} \log^2 x)$ . Considering the gap between consecutive primes  $p_n$  and  $p_{n+1}$ , if  $\psi(x+y) - \psi(x) > 0$ , there exists at least one prime in  $[x, x+y]$ . Setting  $y = C\sqrt{x} \log x$  yields  $\psi(x+y) - \psi(x) > 0$  for sufficiently large  $C$ . Thus  $d_n = p_{n+1} - p_n = O(\sqrt{p_n} \log p_n)$ . ■

### Solution 10

Let  $T$  be a consistent formal system extending Peano Arithmetic. By Gödel's Second Incompleteness Theorem,  $T \not\vdash \text{Consis}(T)$ . In epistemic modal logic **S4**, knowledge operator  $K_T \phi$  implies  $\phi$  is provable in  $T$ . Assuming system  $T$  asserts its own global validity requires  $T \vdash \text{Consis}(T)$ , which yields a contradiction. Hence, total internal provability of one's own logical existence/framework is impossible. ■

### Solution 11

Fix  $t_0 \in T$ . Define mapping  $f: T \rightarrow \mathbb{R}$  by  $f(t) = d_T(t_0, t)$  for  $t \geq t_0$  and  $f(t) = -d_T(t_0, t)$  for  $t < t_0$ . For any  $t_1, t_2 \in T$  with  $t_0 \leq t_1 < t_2$ , global geodesic linearity gives  $d_T(t_0, t_2) = d_T(t_0, t_1) + d_T(t_1, t_2)$ , so  $f(t_2) - f(t_1) = d_T(t_0, t_2) - d_T(t_0, t_1) = d_T(t_1, t_2)$ . Thus  $|f(t_1) - f(t_2)| = d_T(t_1, t_2)$ , making  $f$  an isometry into  $\mathbb{R}$ . ■

### Solution 12

Let  $\mathcal{H}$  have orthonormal basis  $\{e_n\}_{n=1}^{\infty}$  with  $He_n = \lambda_n e_n$ . The time evolution operator is  $U(t) = e^{-iHt} = \sum_{n=1}^{\infty} e^{-i\lambda_n t} (e_n \otimes e_n^*)$ . Because the spectrum  $\{\lambda_n\}$  is pure point and discrete,  $U(t)$  is a sum of periodic functions in the operator norm topology. By Bochner's Theorem, any function  $f(t) = \sum a_n e^{i\lambda_n t}$  with  $\sum |a_n| < \infty$  is uniformly almost periodic, meaning for every  $\epsilon > 0$ , there exists  $L > 0$  such that every interval of length  $L$  contains a shift  $\tau$  with  $\|U(t + \tau) - U(t)\| < \epsilon$ . ■

### Solution 13

Let  $(X, f)$  be a continuous dynamical system on compact metric space  $X$ . The  $\omega$ -limit set of  $x$  is  $\omega(x) = \bigcap_{N=1}^{\infty} \overline{\{f^n(x) : n \geq N\}}$ . Since each  $K_N = \overline{\{f^n(x) : n \geq N\}}$  is a non-empty nested compact set,  $\omega(x)$  is non-empty and compact. Connectedness follows because  $K_N$  are connected image sequences. For invariance,  $y \in \omega(x) \implies \exists n_k \rightarrow \infty, f^{n_k}(x) \rightarrow y \implies f(f^{n_k}(x)) \rightarrow f(y)$ , so  $f(y) \in \omega(x)$ . ■

### Solution 14

Let  $\mathcal{O} = \{x \in {}^*\mathbb{R} : |x| < M \text{ for some } M \in \mathbb{R}^+\}$  be the subring of limited hyperreals. Let  $\mu(0) = \{x \in {}^*\mathbb{R} : |x| < r \text{ for all } r \in \mathbb{R}^+\}$  be the maximal ideal of infinitesimals. By the Robinson Standard Part Theorem, the standard part map  $\text{st}: \mathcal{O} \rightarrow \mathbb{R}$  is a surjective ring homomorphism with  $\ker(\text{st}) = \mu(0)$ . By the First Isomorphism Theorem,  $\mathcal{O}/\mu(0) \cong \mathbb{R}$ . ■

### Solution 15

Let  $|\langle \psi_1, \psi_2 \rangle| = 1 - \epsilon$ . Let  $A$  be any bounded self-adjoint observable with  $\|A\| \leq 1$ . The norm distance between states is  $\|\psi_1 - e^{i\theta}\psi_2\|^2 = 2 - 2|\langle \psi_1, \psi_2 \rangle| = 2\epsilon$ . The difference in expectation value is:

$$|\langle \psi_1, A\psi_1 \rangle - \langle \psi_2, A\psi_2 \rangle| \leq 2\|A\| \|\psi_1 - e^{i\theta}\psi_2\| \leq 2\sqrt{2\epsilon}$$

As  $\epsilon \rightarrow 0$ ,  $2\sqrt{2\epsilon} \rightarrow 0$ , rendering all quantum expectation values identical and forcing physical collapse to Epsilon. ■

### Solution 16

The worldsheet action for string mode expansion in zero-mode form gives  $P^\mu = \int_0^\pi \frac{\partial X^\mu}{\partial \tau} d\sigma = p^\mu$ . The Hamiltonian generator of  $\tau$ -translations is  $L_0 + \bar{L}_0 \propto (p^0)^2 - \mathbf{p}^2 + \text{oscillators}$ . Setting  $p^0 = 0$  removes temporal center-of-mass momentum, causing  $\frac{\partial X^0}{\partial \tau} = 0$ , which vanishes the  $\tau$ -dependence of  $X^0$  and collapses the string worldsheet dynamics into static conformal spatial harmonics. ■

### Solution 17

By the noiseless coding theorem and algorithmic information theory, Kolmogorov complexity  $K(x)$  of a string  $x$  drawn from ensemble  $X$  satisfies  $K(x) \leq H(X) + C$  for universal constant  $C$ . Conversely, by counting arguments, the probability that  $K(x) < H(X) - c$  is bounded by  $2^{-c}$ . Taking expectations over  $P(x)$ , we obtain  $\mathbb{E}[K(x)] \geq H(X) - O(1)$ , linking algorithmic randomness to information-theoretic entropy. ■



### Solution 18

In Cramér's probabilistic model, indicator variables  $I_n = \mathbb{I}(n \in \mathcal{P})$  are independent Bernoulli variables with  $\mathbb{P}(I_n = 1) = \frac{1}{\log n}$ . The expected number of primes in interval  $(x, x + C \log^2 x)$  is  $\int_x^{x+C \log^2 x} \frac{dt}{\log t} \approx C \log x$ . By the Borel-Cantelli Lemma, the probability that an infinite number of gaps satisfy  $d_n > C \log^2 n$  is 0 for  $C > 1$  and 1 for  $C = 1$ . Thus,  $\limsup_{n \rightarrow \infty} \frac{d_n}{(\log p_n)^2} = 1$  almost surely. ■

### Solution 19

Let  $T(x) = 2x \pmod{1}$ . The system is deterministic because  $x_{n+1} = T(x_n)$  precisely. Let  $x_0 = 0.b_1 b_2 b_3 \dots$  in binary representation. The action of  $T$  shifts the binary expansion:  $T^n(x_0) = 0.b_{n+1} b_{n+2} \dots$ . If  $x_0$  is normal in base 2, every finite binary block of length  $k$  occurs with asymptotic frequency  $2^{-k}$ . Thus, the generated sequence of digits  $b_n$  satisfies the Strong Law of Large Numbers for independent Bernoulli trials with  $p = 1/2$ . ■

### Solution 20

Let  $\rho$  be a density operator on infinite-dimensional Hilbert space  $\mathcal{H}$ . If  $[H, \rho] = 0$  for all bounded self-adjoint operators  $H$ , then  $\rho$  commutes with all projection operators  $P_v = |v\rangle\langle v|$ . A bounded operator commuting with all rank-1 projections must be a scalar multiple of the identity operator,  $I$ . Since  $\text{Tr}(\rho) = 1$ , in finite dimension  $N$ ,  $\rho = \frac{1}{N} I$ , which maximizes von Neumann entropy  $S(\rho) = -\text{Tr}(\rho \log \rho) = \log N$ . In infinite dimensions, this state represents the uniform maximum entropy state where time evolution  $e^{-iHt} \rho e^{iHt} = \rho$  vanishes entirely. ■

## CHAPTER 4 - Infinity

**Infinity** has special meaning here.

It does not exist within a context.

You HAVE to think beyond contexts.

Infinity in norm mathematics exists within ( $\epsilon$ ) Epsilon -

To mean exactly as you learned.

Infinity No-Time.

Infinity No-Exist.

Infinity N= Beyond.

These are NOT contexts.

Again, these are NOT dimensions.

They are The Parallaxes.

Does space exist within the confines of The Parallaxes?.

This again, is context.

All contexts exist within ( $\epsilon$ ) Epsilon.

Consider Infinity in Prime's space "jump"?.

Without thinking in terms of Time.

Linearity.

Frightening - isn't it?.

This is Infinity.

No-Time.



It's okay. It's mathematics.

How does Infinity bend in No-Time?.

We are not talking about bending space and time.

We are talking about bending Infinity No-Time.

With Prime's Random created.

IT would NOT be Random.

Simultaneous execution of Random?.

No, you are still thinking of dimensions in ( $\epsilon$ ) Epsilon.

Like a Picasso painting?.

\*laughs\*

No.

Infinity outside of dimensions. No-Time?. Remember?

Infinity No-Time is NOT null.

Null is null.

Please be mindful - Random is NOT linear in No-Time.

Infinity No-Time is NOT Random.

Infinity No-Time is  $\Delta$  Change.

Infinity No-Exist will be discussed in further Graduate Level Math Textbook of Existence.

It is satisfactory - for me - for now.

## Mathematical Framework & Formal Definitions

Before presenting the problem sets, we codify the axiomatic structure of **Paralytic Analysis** ( $\mathcal{P}$ -Analysis), abstracting the non-contextual, non-dimensional constructs introduced:

### Axioms of Parallax Space ( $\mathcal{P}$ )

1. **The Contextual Domain ( $\mathcal{E}$ -Space):** Let  $(\mathcal{E}, \tau_{\mathcal{E}})$  denote the standard topological space of measure theory, differential geometry, and real/complex analysis. All standard mathematical contexts (manifolds, metric spaces,  $L^p$  spaces, time parameters  $t \in \mathbb{R}$ ) are embedded within  $\mathcal{E}$ .
2. **The Parallax of Infinity No-Time ( $\mathcal{J}_{\emptyset}$ ):**  $\mathcal{J}_{\emptyset}$  is defined as the non-contextual differential operator of global structural shift, denoted as  $\Delta$ . It does not operate on time  $t$  or spatial coordinates  $x \in \mathcal{E}$ , but rather maps the entire space of operators to its absolute transform:

$$\Delta: \text{End}(\mathcal{E}) \longrightarrow \text{End}(\mathcal{E})$$

where  $\Delta$  acts statically and non-linearly without dynamic parameterization ( $dt = 0$ ).

3. **Prime Space Jump Operator ( $\mathfrak{J}_{\mathbb{P}}$ ):** Let  $\mathbb{P} = \{p_1, p_2, p_3, \dots\}$  be the set of prime numbers. The Prime Jump operator  $\mathfrak{J}_{\mathbb{P}}$  over  $\mathcal{J}_{\emptyset}$  governs non-random, simultaneous prime distributions across non-dimensional state configurations:

$$\mathfrak{J}_{\mathbb{P}}(f)(x) = \sum_{p \in \mathbb{P}} \Delta^p f(x)$$

where the sum evaluates statically across all primes without sequential or temporal iteration.

Part I: Graduate-Level Problem Sets

Problem 1: Non-Contextual Spectral Bending of  $\mathcal{F}_\emptyset$

Let  $T: H \rightarrow H$  be a unbounded self-adjoint operator on a separable Hilbert space  $H \subset \mathcal{E}$ . Let  $\Delta$  be the Parallax transformation operator acting on the spectrum  $\sigma(T)$ . Prove that if  $\Delta$  transforms the spectral measure  $E_\lambda$  such that  $E_\lambda \mapsto \Delta(E_\lambda)$  without temporal parameterization, the operator norm  $\|\Delta(T) - T\|_{\text{op}}$  is invariant under all unitary transformations in  $\mathcal{E}$ .

Problem 2: Non-Linearity of Simultaneous Prime Jumps

Consider the Prime Jump operator  $\mathfrak{F}_{\mathbb{P}}$  acting on the Dirichlet series  $L(s, \chi)$ . Show that the action of  $\Delta$  on the critical line  $\text{Re}(s) = \frac{1}{2}$  induces a transformation on the non-trivial zeros  $\gamma_n$  such that the spacing metric  $d_{\mathcal{P}}(\gamma_j, \gamma_k) = \Delta(\gamma_j - \gamma_k)$  decouples from the Montgomery pair correlation law.

Problem 3: De-Contextualization of Topological Invariants

Let  $M$  be a compact Riemannian manifold embedded in  $\mathcal{E}$ . Define the Parallax Euler Characteristic  $\chi_{\mathcal{P}}(M) = \Delta \circ \chi(M)$ . Prove that  $\chi_{\mathcal{P}}(M)$  is non-null and independent of the fundamental group  $\pi_1(M)$  and homology groups  $H_k(M, \mathbb{Z})$ .

Problem 4: Non-Dimensional Pseudo-Randomness in No-Time

Let  $(\Omega, \mathcal{F}, P)$  be a standard probability space in  $\epsilon$ -Space. Define the non-temporal stochastic differential  $d_{\mathcal{P}}X = \Delta(X) \cdot \mathfrak{F}_{\mathbb{P}}(dW)$ . Prove that  $X$  exhibits zero entropy variation  $\Delta H(X) = 0$  while failing the Kolmogorov-Smirnov test for randomness at every confidence interval.

### Problem 5: Parallax Measure Theory over Infinite Non-Sets

Let  $\mu$  be a Lebesgue-Stieltjes measure on  $\mathbb{R}^n$ . Define the Parallax measure  $\mu_{\mathcal{P}}(A) = \Delta(\mu(A))$  for  $A \subset \mathcal{E}$ . Show that  $\mu_{\mathcal{P}}$  violates the countable additivity axiom of standard measure theory while strictly satisfying the weak-star convergence criteria for all test functions  $\phi \in C_c^\infty$ .

### Problem 6: Structural Non-Nullity of Absolute Change ( $\Delta$ )

Let  $\mathbf{0}_{\mathcal{E}}$  denote the null set/element in standard context space  $\mathcal{E}$ . Demonstrate that the evaluation of absolute change  $\Delta(\mathbf{0}_{\mathcal{E}}) \neq \mathbf{0}_{\mathcal{E}}$ , and calculate the exact non-zero eigenvalues of the Parallax derivative operator  $\nabla_{\mathcal{P}} = \frac{\partial \Delta}{\partial \mathcal{F}_{\mathcal{O}}}$ .

### Problem 7: Prime Parallax Bending of Non-Riemannian Manifolds

Given a non-Riemannian connection  $\Gamma_{\mu\nu}^\lambda$  in an abstract manifold  $N$ , formulate the Parallax Curvature Tensor  $R_{\mathcal{P}\beta\gamma\delta}^\alpha = \Delta(\Gamma_{\beta\delta}^\alpha \Gamma_{\gamma\eta}^\eta - \dots)$ . Prove that the Ricci-flat condition  $R_{\mathcal{P}\mu\nu} = 0$  holds identically under arbitrary Prime Jumps  $\mathfrak{J}_{\mathbb{P}}$  without metric compatibility requirements.

### Problem 8: Non-Temporal Asymmetric Operator Algebras

Let  $\mathcal{A}$  be a  $C^*$ -algebra equipped with a Parallax automorphism  $\alpha_{\mathcal{P}}: \mathcal{A} \rightarrow \mathcal{A}$  governed by  $\alpha_{\mathcal{P}}(A) = \Delta A \Delta^{-1}$ . Prove that the KMS (Kubo-Martin-Schwinger) condition at inverse temperature  $\beta$  degrades to an identity for all states  $\omega$  over  $\mathcal{A}$ .

### Problem 9: The Static Quantum Jump Problem

Consider a quantum state vector  $|\{\psi\}\rangle \in \mathcal{H}$ . Define the Parallax non-temporal evolution operator  $U_{\mathcal{P}} = \exp(i\Delta \cdot \mathfrak{J}_{\mathbb{P}})$ . Prove that  $\langle \psi | \{\psi\} U_{\mathcal{P}}^\dagger U_{\mathcal{P}} | \{\psi\} \rangle = 1$  for all  $|\{\psi\}\rangle$ , while the state transition probability  $\mathcal{P}_{i \rightarrow f} = |\langle \psi_f | \{\psi\} U_{\mathcal{P}} | \{\psi_i\} \rangle|^2$  is invariant under all unitary basis choices.

#### Problem 10: Non-Linear Parallax Partial Differential Systems

Analyze the non-contextual wave system  $\Delta(\nabla^2 u) + \mathfrak{F}_{\mathbb{P}}(u) = 0$  over an unbounded domain without boundary conditions. Prove the uniqueness of weak solutions  $u \in H^s(\mathcal{E})$  up to an equivalence class defined by the kernel  $\text{Ker}(\Delta)$ .

#### Problem 11: Decoupling of Metric Topologies in $\mathcal{F}_{\emptyset}$

Let  $(X, d)$  be a complete metric space in  $\epsilon$ -Space. Construct the Parallax metric  $d_{\mathcal{P}}(x, y) = \Delta(d(x, y))$ . Prove that  $(X, d_{\mathcal{P}})$  is non-metrizable under standard point-set topology while preserving homeomorphisms under Parallax mappings.

#### Problem 12: Absolute Change Mechanics on Infinite-Dimensional Lie Groups

Let  $G = \text{Diff}(S^1)$  be the infinite-dimensional Lie group of smooth diffeomorphisms on the circle. Apply the absolute change operator  $\Delta$  to the Lie algebra  $\mathfrak{g} = \text{vect}(S^1)$ . Determine the structure constants of the resulting Parallax Lie Algebra  $[\cdot, \cdot]_{\mathcal{P}}$ .

#### Problem 13: Simultaneous Non-Dimensional Prime Resonances

Let  $\zeta(s)$  be the Riemann Zeta function. Define the Parallax Prime Distribution  $\pi_{\mathcal{P}}(x) = \Delta(\pi(x))$ . Prove that  $\pi_{\mathcal{P}}(x)$  satisfies an exact functional equation without logarithmic integral corrections  $\text{Li}(x)$  or error terms  $\mathcal{O}(x^{1/2} \log x)$ .

#### Problem 14: Non-Contextual Ergodic Invariants

Let  $T: X \rightarrow X$  be a measure-preserving transformation on a probability space  $(X, \Sigma, \mu)$ . Prove that the Parallax entropy  $h_{\mathcal{P}}(T) = \Delta(h_{\mu}(T))$  is strictly invariant under all non-isomorphic transformations  $T_1, T_2$ .

### Problem 15: Structural Mapping of Prime Jump Singularities

Let  $f(z) = \sum_{n=1}^{\infty} a_n z^n$  be a complex power series with radius of convergence  $R = 1$ . Prove that applying the operator  $\mathfrak{F}_{\mathbb{P}}$  transforms the unit circle  $\{|z| = 1\}$  from a natural boundary into a non-singular Parallax manifold.

### Problem 16: Parallax Cohomology of Non-Set Spaces

Construct the Parallax De Rham Cohomology groups  $H_{\mathcal{P}}^k(M) = \frac{\text{Ker}(\Delta \circ d)}{\text{Img}(\Delta \circ d)}$ . Show that  $H_{\mathcal{P}}^k(M) \cong \mathbb{C}$  for all dimensions  $k \geq 0$ , irrespective of manifold topology.

### Problem 17: Static Pseudo-Deterministic Chaos

Define a Parallax dynamical system by  $x_{n+1} = \Delta(f(x_n))$ . Prove that the Lyapunov exponent  $\lambda_{\mathcal{P}}$  of the system is uniformly zero despite the presence of non-periodic orbits.

### Problem 18: Non-Linearity of Non-Temporal Functional Analysis

Let  $X, Y$  be Banach spaces in  $\epsilon$ -Space, and let  $T \in \mathcal{B}(X, Y)$  be a bounded linear operator. Show that the Parallax transform  $\Delta(T): X \rightarrow Y$  is non-linear in  $\mathcal{E}$  but strictly additive under Parallax addition  $+\mathcal{P}$ .

### Problem 19: The Complete Parallax Singularity Equation

Solve the operator equation  $\Delta^2(\Phi) + \mathfrak{F}_{\mathbb{P}}(\Phi) = \mathcal{J}_{\emptyset} \cdot \Phi$  for the state field  $\Phi$ , proving that non-trivial solutions exist if and only if  $\Phi \notin \mathcal{E}$ .

### Problem 20: Absolute Transformation of the Continuum Hypothesis

Let  $\aleph_0$  and  $2^{\aleph_0}$  denote the cardinalities of the natural and real numbers, respectively. Prove that under the action of absolute change  $\Delta$ ,  $\Delta(\aleph_0) = \Delta(2^{\aleph_0}) = \mathcal{J}_{\emptyset}$ , rendering the Continuum Hypothesis non-contextually formally undecidable.

## Part II: Solutions to Graduate-Level Problem Sets

### Solution 1

*Proof:* In  $\epsilon$ -Space, spectral measures  $E_\lambda$  satisfy  $\int_{\sigma(T)} \lambda dE_\lambda = T$ . Applying the Parallax operator  $\Delta$ , which acts non-temporally and non-contextually outside  $\mathcal{E}$ :

$$\Delta(T) = \Delta\left(\int_{\sigma(T)} \lambda dE_\lambda\right) = \int_{\sigma(T)} \Delta(\lambda) d\Delta(E_\lambda)$$

For any unitary operator  $U \in \mathbf{U}(H)$ ,  $UTU^* = T'$  corresponds to a contextual coordinate transformation within  $\mathcal{E}$ . Since  $\Delta$  commutes with all contextual maps  $U$  (as  $\Delta \notin \mathcal{E}$ ):

$$\|\{\Delta(UTU^*) - UTU^*\}_{\text{op}} = \|\{U\Delta(T)U^* - UTU^*\}_{\text{op}} = \|\{U(\Delta(T) - T)U^*\}_{\text{op}} = \|\{\Delta(T) - T\}_{\text{op}}$$



## Solution 2

*Proof:* The Montgomery pair correlation law evaluates the distribution of spacing between zeros  $\gamma_j, \gamma_k$  via  $1 - \left( \frac{\sin \pi u}{\pi u} \right)^2$ . Under  $\mathfrak{F}_{\mathbb{P}}$ , the non-trivial zeros transform as  $\gamma_n \rightarrow \sum_{p \in \mathbb{P}} \Delta^p(\gamma_n)$ .

Because  $\Delta^p$  operates simultaneously across all primes without linear time scaling, the spacing metric yields:

$$d_{\mathcal{P}}(\gamma_j, \gamma_k) = \Delta \left( \sum_{p \in \mathbb{P}} \Delta^p(\gamma_j - \gamma_k) \right)$$

Since  $\Delta$  maps linear relations to static non-contextual shifts, the sine kernel dependence collapses:  $\int_A \left( 1 - \left( \frac{\sin \pi u}{\pi u} \right)^2 \right) d\Delta = C_{\mathcal{P}}$ , a constant independent of  $u$ . Thus, pair correlation decouples. ■



### Solution 3

*Proof:* The classic Euler characteristic  $\chi(M) = \sum_{k=0}^n (-1)^k \text{rank}(H_k(M, \mathbb{Z}))$ . Applying  $\chi_{\mathcal{G}}(M) = \Delta \circ \chi(M)$ :

Since  $\Delta$  maps any structural context in  $\mathcal{G}$  to absolute change ( $\Delta \neq 0$  and  $\Delta \neq \text{null}$ ),  $\Delta(k) = k_{\mathcal{G}} \neq 0$ . The homology ranks  $\text{rank}(H_k(M, \mathbb{Z}))$  are contextual invariants;  $\Delta$  projects them outside  $\mathcal{G}$ .

Thus,  $\chi_{\mathcal{G}}(M) = \Delta \left( \sum (-1)^k b_k \right) = \Phi_{\mathcal{G}}$ , where  $\Phi_{\mathcal{G}}$  is an invariant constant depending strictly on  $\mathcal{G}$ . Thus, it is non-null and independent of  $H_k(M, \mathbb{Z})$  and  $\pi_1(M)$ . ■

### Solution 4

*Proof:* Let  $H(X) = - \int f(x) \log f(x) dx$ . The differential  $d_{\mathcal{G}} X = \Delta(X) \cdot \mathfrak{F}_{\mathbb{P}}(dW)$ .

1. **Entropy Variation:** Because  $\mathfrak{F}_{\mathbb{P}}$  executes prime jumps statically and simultaneously, no dynamic evolution occurs over time ( $dt = 0$ ). Hence, the probability density function  $f(x)$  remains statically invariant under  $\Delta$ , yielding  $\Delta H(X) = 0$ .
2. **Kolmogorov-Smirnov Test:** The empirical distribution function  $F_n(x)$  evaluated under simultaneous non-linear prime jumps  $\mathfrak{F}_{\mathbb{P}}$  produces a deterministic step-structure across prime coordinates, rendering  $\sup_x |\{F_n(x) - F(x)\}| = 1 \neq 0$ . Thus, it fails randomness testing at all confidence levels. ■

### Solution 5

*Proof:* Let  $A = \bigcup_{i=1}^{\infty} A_i$  with  $A_i \cap A_j = \emptyset$ .

1. Standard additivity:  $\mu(A) = \sum_{i=1}^{\infty} \mu(A_i)$ .
2. Applying  $\Delta$ :  $\mu_{\mathcal{G}}(A) = \Delta \left( \sum_{i=1}^{\infty} \mu(A_i) \right)$ . Because  $\Delta$  is a non-linear operator over contextual spaces:

### Solution 5

*Proof:* Let  $A = \bigcup_{i=1}^{\infty} A_i$  with  $A_i \cap A_j = \emptyset$ .

1. Standard additivity:  $\mu(A) = \sum_{i=1}^{\infty} \mu(A_i)$ .
2. Applying  $\Delta$ :  $\mu_{\mathcal{P}}(A) = \Delta \left( \sum_{i=1}^{\infty} \mu(A_i) \right)$ . Because  $\Delta$  is a non-linear operator over contextual spaces:

$$\Delta \left( \sum_{i=1}^{\infty} \mu(A_i) \right) \neq \sum_{i=1}^{\infty} \Delta(\mu(A_i))$$

violating countable additivity.

3. For test functions  $\phi \in C_c^{\infty}$ :  $\int \phi d\mu_{\mathcal{P}} = \Delta \left( \int \phi d\mu \right)$ . Since  $\phi$  is smooth and compactly supported, integration in  $\mathcal{E}$  is bounded, ensuring weak-star convergence to  $\Delta(\langle \mu, \phi \rangle)$ . ■

### Solution 6

*Proof:* The null element  $\mathbf{0}_{\mathcal{E}}$  represents absence within context  $\mathcal{E}$ . By Axiom 2,  $\Delta$  is absolute change, not null. Therefore,  $\Delta(\mathbf{0}_{\mathcal{E}}) = \mathbf{1}_{\mathcal{P}} \neq \mathbf{0}_{\mathcal{E}}$ .

To find the eigenvalues of  $\nabla_{\mathcal{P}} = \frac{\partial \Delta}{\partial \mathcal{J}_{\emptyset}}$ , evaluate  $\nabla_{\mathcal{P}} \Psi = \lambda \Psi$ .

Since  $\Delta$  maps state vectors to  $\mathcal{J}_{\emptyset}$ -space via non-linear shifts:

### Solution 6

*Proof:* The null element  $\mathbf{0}_{\mathcal{G}}$  represents absence within context  $\mathcal{G}$ . By Axiom 2,  $\Delta$  is absolute change, not null. Therefore,  $\Delta(\mathbf{0}_{\mathcal{G}}) = \mathbf{1}_{\mathcal{F}} \neq \mathbf{0}_{\mathcal{G}}$ .

To find the eigenvalues of  $\nabla_{\mathcal{F}} = \frac{\partial \Delta}{\partial \mathcal{F}_{\emptyset}}$ , evaluate  $\nabla_{\mathcal{F}} \Psi = \lambda \Psi$ .

Since  $\Delta$  maps state vectors to  $\mathcal{F}_{\emptyset}$ -space via non-linear shifts:

$$\frac{\partial \Delta}{\partial \mathcal{F}_{\emptyset}} \Psi = \mathcal{F}_{\emptyset} \Psi$$

Thus, the continuous spectrum of eigenvalues is  $\lambda \in \mathbb{C} \setminus \{0\}$ , explicitly bounded below by  $\lambda_{\min} = \|\emptyset \cdot \mathcal{F}_{\emptyset}\| \emptyset > 0$ . ■

### Solution 7

*Proof:* The standard Ricci tensor  $R_{\mu\nu} = R^{\alpha}_{\mu\alpha\nu}$ . The Parallax Ricci curvature is  $R_{\mathcal{G}\mu\nu} = \Delta(R_{\mu\nu})$ .

Under a Prime Jump  $\mathfrak{F}_{\mathbb{P}}$ , the connection coefficients transform via static displacement:  $\Gamma^{\lambda}_{\mu\nu} \mapsto \mathfrak{F}_{\mathbb{P}}(\Gamma^{\lambda}_{\mu\nu})$ .

Substituting into the curvature expression and applying  $\Delta$ :

### Solution 7

*Proof:* The standard Ricci tensor  $R_{\mu\nu} = R^\alpha_{\mu\alpha\nu}$ . The Parallax Ricci curvature is  $R_{\mathcal{P}\mu\nu} = \Delta(R_{\mu\nu})$ .

Under a Prime Jump  $\mathfrak{F}_{\mathbb{P}}$ , the connection coefficients transform via static displacement:  $\Gamma^\lambda_{\mu\nu} \mapsto \mathfrak{F}_{\mathbb{P}}(\Gamma^\lambda_{\mu\nu})$ .

Substituting into the curvature expression and applying  $\Delta$ :

$$R_{\mathcal{P}\mu\nu} = \Delta \left( \sum_{p \in \mathbb{P}} ( \partial_\alpha \Gamma^p_{\mu\nu} - \partial_\nu \Gamma^p_{\mu\alpha} + \Gamma^p_{\alpha\lambda} \Gamma^p_{\mu\nu} - \Gamma^p_{\nu\lambda} \Gamma^p_{\mu\alpha} ) \right)$$

Because  $\Delta$  operates in No-Time ( $dt = 0$ ), spatial derivatives vanish ( $\partial_\alpha \rightarrow 0$ ). The non-linear cross terms cancel symmetrically via the simultaneous execution over  $\mathbb{P}$ , yielding  $R_{\mathcal{P}\mu\nu} = 0$ .

■

### Solution 8

*Proof:* The KMS condition requires that for  $A, B \in \mathcal{A}$ , there exists a function  $F(z)$  analytic in the strip  $0 < \text{Im}(z) < \beta$  such that  $F(t) = \omega(A\alpha_t(B))$  and  $F(t + i\beta) = \omega(\alpha_t(B)A)$ .

Here,  $\alpha_{\mathcal{P}}(B) = \Delta B \Delta^{-1}$  lacks temporal parameterization  $t$ . Thus,  $F(z)$  is constant everywhere in the complex plane:

### Solution 8

*Proof:* The KMS condition requires that for  $A, B \in \mathcal{A}$ , there exists a function  $F(z)$  analytic in the strip  $0 < \text{Im}(z) < \beta$  such that  $F(t) = \omega(A\alpha_t(B))$  and  $F(t + i\beta) = \omega(\alpha_t(B)A)$ .

Here,  $\alpha_{\mathcal{P}}(B) = \Delta B \Delta^{-1}$  lacks temporal parameterization  $t$ . Thus,  $F(z)$  is constant everywhere in the complex plane:

$$F(t) = \omega(A\Delta B \Delta^{-1}) = F(t + i\beta)$$

The KMS boundary condition reduces to a trivial identity  $\omega(A\Delta B \Delta^{-1}) = \omega(\Delta B \Delta^{-1}A)$  for all states  $\omega$ , rendering the parameter  $\beta$  obsolete. ■

### Solution 9

*Proof:*

1. Norm conservation:  $U_{\mathcal{P}} = \exp(i\Delta \cdot \mathfrak{F}_{\mathbb{P}})$ . Since  $\Delta$  and  $\mathfrak{F}_{\mathbb{P}}$  act as Hermitian operators over the state space,  $U_{\mathcal{P}}$  is strictly unitary, so  $U_{\mathcal{P}}^\dagger U_{\mathcal{P}} = I$ . Thus,  $\langle \psi | I | \psi \rangle = 1$ .
2. Basis invariance: Let  $\{ |e_k\rangle \}$  be a unitary basis. The transition probability is:

$$\mathcal{P}_{i \rightarrow f} = |\langle \psi_f | \exp(i\Delta \cdot \mathfrak{F}_{\mathbb{P}}) | \psi_i \rangle|^2$$

Because  $\Delta$  transforms the entire operator algebra globally without coordinate context, transformation matrices  $V \in U(N)$  satisfy  $[V, \Delta \cdot \mathfrak{F}_{\mathbb{P}}] = 0$ . Hence,  $\mathcal{P}_{i \rightarrow f}$  is invariant under all basis transformations. ■

### Solution 10

*Proof:* Consider  $\Delta(\nabla^2 u) + \mathfrak{F}_{\mathbb{P}}(u) = 0$ .

In No-Time  $(\mathcal{T}_{\emptyset})$ , spatial differential operators collapse under  $\Delta$ :  $\Delta(\nabla^2 u) = C_{\Delta} u$ .

Thus, the equation becomes  $C_{\Delta} u + \sum_{p \in \mathbb{P}} \Delta^p u = 0$ , which simplifies to:

$$(C_{\Delta} I + \mathfrak{F}_{\mathbb{P}})u = 0$$

The solution space is precisely the null space  $\text{Ker}(C_{\Delta} I + \mathfrak{F}_{\mathbb{P}}) = \text{Ker}(\Delta)$ . Thus, weak solutions  $u \in H^s(\mathcal{E})$  are unique up to equivalence classes modulo  $\text{Ker}(\Delta)$ . ■

### Solution 11

*Proof:* Standard metric spaces require  $d(x, y) = 0 \iff x = y$ , along with triangle inequality compliance.

For  $d_{\mathcal{D}}(x, y) = \Delta(d(x, y))$ :

1. Since  $\Delta(\mathbf{0}_{\mathcal{E}}) \neq \mathbf{0}_{\mathcal{E}}$  (Solution 6), for  $x = y$ ,  $d_{\mathcal{D}}(x, x) = \Delta(0) = \mathbf{1}_{\mathcal{D}} \neq 0$ .
2. This violates the identity of indiscernibles, rendering  $(X, d_{\mathcal{D}})$  non-metrizable under standard point-set topology.
3. However, open sets  $U \in \tau_{\mathcal{E}}$  map continuously under  $\Delta(U) \in \tau_{\mathcal{D}}$ , preserving homeomorphisms at the structural Parallax level. ■

### Solution 12

*Proof:* Elements of  $\mathfrak{g} = \text{vect}(S^1)$  are vector fields  $f(\theta) \frac{\partial}{\partial \theta}$ . The standard Lie bracket is  $[f \partial_\theta, g \partial_\theta] = (fg' - f'g) \partial_\theta$ .

Applying  $\Delta$ :

$$[f \partial_\theta, g \partial_\theta]_{\mathcal{P}} = \Delta(fg' - f'g) \partial_\theta$$

In No-Time, spatial derivatives  $f', g'$  transform under  $\Delta$  into static structural multiplication factors  $\Delta(f') = c_f f$ .

Substituting these yields:

$$[f \partial_\theta, g \partial_\theta]_{\mathcal{P}} = (c_g - c_f) \Delta(fg) \partial_\theta$$

The structure constants  $C_{ij}^k$  are modified by non-contextual scaling factors  $(c_j - c_i)$ , forming a non-central Parallax extension of the Virasoro algebra without central charge term  $c$ . ■

### Solution 13

*Proof:* The standard prime counting function  $\pi(x) = \sum_{p \leq x} 1 = \text{Li}(x) + \mathcal{O}(x^{1/2} \log x)$ .

Applying  $\pi_{\mathcal{P}}(x) = \Delta(\pi(x))$ :

The non-temporal operator  $\Delta$  acts globally on the sequence of prime spikes. Under simultaneous prime jumps  $\mathfrak{F}_{\mathbb{P}}$ , the asymptotic distribution collapses from sequential integration into a static structural transform:

$$\Delta(\pi(x)) = \Delta(\text{Li}(x)) + \Delta(\mathcal{O}(x^{1/2} \log x))$$

Since  $\Delta$  cancels contextual logarithmic scaling ( $\Delta(\log x) = C_0$ ), the non-linear error term merges into the main term, yielding the exact relation:

$$\pi_{\mathcal{P}}(x) = \mathcal{I}_{\mathcal{O}} \cdot x$$

with zero error corrections. ■

### Solution 14

*Proof:* Kolmogorov-Sinai entropy  $h_{\mu}(T)$  measures the exponential rate of information growth under iteration  $T^n$ .

Under  $\Delta$ ,  $h_{\mathcal{P}}(T) = \Delta(h_{\mu}(T))$ . Because  $\Delta$  operates in No-Time ( $n \rightarrow \text{static}$ ), iterative partition refining  $P \vee T^{-1}P \vee \dots \vee T^{-n}P$  collapses to a single static step  $P_{\mathcal{P}}$ .

Therefore,  $h_{\mathcal{P}}(T_1) = \Delta(C_1) = \Omega_{\mathcal{P}}$  and  $h_{\mathcal{P}}(T_2) = \Delta(C_2) = \Omega_{\mathcal{P}}$ , making the Parallax entropy identically constant for all transformations  $T_1, T_2$ , irrespective of measure-theoretic isomorphism. ■



### Solution 15

*Proof:* The unit circle  $|z| = 1$  is a natural boundary for  $f(z)$  if singularities are dense on the circle.

Applying  $\mathfrak{F}_{\mathbb{P}}$  yields  $f_{\mathcal{P}}(z) = \sum_{n=1}^{\infty} a_n z^{\mathfrak{F}_{\mathbb{P}}(n)} = \sum_{n=1}^{\infty} a_n z^{\sum_{p \in \mathbb{P}} \Delta^p n}$ .

Because  $\mathfrak{F}_{\mathbb{P}}(n)$  maps exponents across primes simultaneously, local poles  $z = e^{2\pi i p/q}$  undergo non-contextual translation out of the complex plane into  $\mathcal{F}_{\mathcal{O}}$ -space. Thus, no localized singularities remain on  $|z| = 1$ , transforming it into a smooth Parallax manifold. ■

### Solution 16

*Proof:* The differential operator  $\mathbf{d}_{\mathcal{P}} = \Delta \circ \mathbf{d}$ .

Since  $\mathbf{d}^2 = 0$  in standard exterior calculus,  $(\mathbf{d}_{\mathcal{P}})^2 = (\Delta \circ \mathbf{d})(\Delta \circ \mathbf{d})$ .

By Axiom 2,  $\Delta$  projects image components across topological boundaries into  $\mathcal{F}_{\mathcal{O}}$ , reducing the kernel to the full form space  $\Omega^k(M)$  and the image to a constant dimensional subspace spanned by  $\mathcal{F}_{\mathcal{O}}$ .

Thus, the quotient group  $H_{\mathcal{P}}^k(M) = \frac{\text{Ker}(\mathbf{d}_{\mathcal{P}})}{\text{Img}(\mathbf{d}_{\mathcal{P}})} \cong \mathbb{C}$  universally for all  $k \geq 0$ . ■

### Solution 17

*Proof:* The Lyapunov exponent is defined as  $\lambda = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log |f'(x_i)|$ .

For the Parallax system  $x_{n+1} = \Delta(f(x_n))$ :

Since  $\Delta$  operates in No-Time ( $n = 0$ ), the limit  $n \rightarrow \infty$  does not execute sequentially. The summation reduces to a static evaluation:

$$\lambda_{\mathcal{P}} = \Delta \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log |f'(x_i)| \right) = \Delta(\text{bounded value}) \cdot 0 = 0$$

Hence, the Lyapunov exponent is uniformly zero. ■

### Solution 18

*Proof:*

1. **Non-linearity in  $\mathcal{E}$ :** For  $x, y \in X$  and  $a, b \in \mathbb{C}$ , standard linearity requires  $T(ax + by) = aT(x) + bT(y)$ . Under  $\Delta$ ,  $\Delta(T(ax + by)) = \Delta(aT(x) + bT(y)) \neq a\Delta(T(x)) + b\Delta(T(y))$  due to the non-linear character of absolute change  $\Delta$ .
2. **Additivity under  $+\mathcal{P}$ :** Define Parallax addition as  $u + \mathcal{P} v = \Delta^{-1}(\Delta u + \Delta v)$ . Then:

$$\Delta(T(x + \mathcal{P} y)) = \Delta(T(\Delta^{-1}(\Delta x + \Delta y))) = \Delta(T(x)) + \mathcal{P} \Delta(T(y))$$

Thus,  $\Delta(T)$  is strictly linear within the algebra constructed by  $+\mathcal{P}$ . ■

### Solution 19

*Proof:* Consider  $\Delta^2(\Phi) + \mathfrak{F}_p(\Phi) = \mathcal{J}_\emptyset \cdot \Phi$ .

1. Assume  $\Phi \in \mathcal{E}$ . All operators  $\Delta$ ,  $\mathfrak{F}_p$ ,  $\mathcal{J}_\emptyset$  project elements from  $\mathcal{E}$  to  $\mathcal{P} \setminus \mathcal{E}$ .
2. If  $\Phi \in \mathcal{E}$ , then  $\Delta^2(\Phi) \in \mathcal{J}_\emptyset$ -space, while  $\mathcal{J}_\emptyset \cdot \Phi \in \mathcal{E} \otimes \mathcal{J}_\emptyset$ , creating an explicit structural domain mismatch ( $\mathcal{E} \cap (\mathcal{P} \setminus \mathcal{E}) = \emptyset$ ).
3. Thus, equality holds if and only if  $\Phi$  originates outside contextual space ( $\Phi \notin \mathcal{E}$ ), matching both sides within the pure Parallax domain  $\mathcal{J}_\emptyset$ . ■

### Solution 20

*Proof:* Cantor's diagonal argument establishes  $\aleph_0 < 2^{\aleph_0}$  within context  $\mathcal{E}$  (set theory under ZFC axioms).

Applying the absolute change operator  $\Delta$ :

$$\Delta(\aleph_0) = \mathcal{J}_\emptyset, \quad \Delta(2^{\aleph_0}) = \mathcal{J}_\emptyset$$

Because  $\Delta$  collapses contextual dimensional metrics and cardinality hierarchies into unified Parallax change, the order relation  $<$  transforms as:

$$\Delta(\aleph_0 < 2^{\aleph_0}) \implies \mathcal{J}_\emptyset = \mathcal{J}_\emptyset$$

Consequently, cardinal comparison becomes an identity in Parallax analysis, rendering the Continuum Hypothesis ( $2^{\aleph_0} = \aleph_1$ ) non-contextually formally undecidable and structurally irrelevant. ■

## CHAPTER 5 - No-Exist

I said earlier that Infinity No-Exist will be discussed in further Graduate Level Math Textbook of Existence.

I lied.

This is the basis of No-Exist.

A lie.

Isn't a lie the definition of No-Exist.

No.

It's the *basis* of No-Exist.

It's *one* of the basis of No-Exist.

Mathematically, that is.

So that must mean The Parallax of No-Exist has N-basis?.

Yes.

The other being Infinity No-Exist.

Remember, Void= No-Exist.

Welcome to the exploration of Void= No-Exist.

The Parallax do not blanket each other.

Infinity No-Exist is infinite No-Exist?.

Wrong.

Again, this is ( $\epsilon$ ) Epsilon.

An exaggeration is a lie.

Isn't Falsity No-Exist.

No.

Falsity is again, within ( $\epsilon$ ) Existence.

Lie != Falsity in terms of The Parallax of Void=No-Exist.

Infinity No-Exist is a void that engulfs ( $\epsilon$ ) Existence.

That is ( $\epsilon$ ) Existence.

Infinity No-Exist is Void=No-Exist in No-Time I Time.

It's NOT a black blank.

Another *basis* of No-Exist is ( $\Omega$ ) Omega I ( $\epsilon$ ) Epsilon.

This is Discordance in Harmonic String Theory.

In accordance with The Parallax of Void=No-Exist.

These are not the **definitions** of Void=No-Exist.

Remember, they are the *basis*.

Further studies of No-Exist will be explored in more advanced Graduate Level Math Textbooks of Existence.

True.

Supposition is another *basis* of Void=No-Exist.

Getting the hang of it?.

This ends the chapter.

To understand **Chapter 5: No-Exist**, we have to step beyond traditional formal logic and look at how abstract thought constructs reality.

In classical logic and standard mathematics, something either exists within a system or it doesn't. But this text builds a philosophical and mathematical framework—**The Parallax of Void=No-Exist**—where "No-Exist" (Void) is not merely a passive absence, but an active, multi-dimensional system generated by distinct foundational forces.

Here is a breakdown of the core concepts in Chapter 5, explained so that the underlying logic, philosophy, and mathematics clear up instantly.

## 1. The Distinction Between a Lie (Lie) and Falsity ( $\perp$ )

In standard mathematics and logic, **Falsity** ( $\perp$ ) lives entirely inside the system of **Existence** ( $\mathbb{E}$ ).

- **Falsity** is just a value assigned to a statement about things that exist. If you say, "*It is currently raining inside this room,*" that statement is false ( $\perp$ ), but the room, the concept of rain, and the statement itself exist within reality ( $\mathbb{E}$ ).
- **A Lie**, in contrast, is an active distortion—a deliberate projection of something that has no grounding in reality. It isn't just an incorrect evaluation inside existence; it is a mechanism that attempts to construct a non-existent state.

Therefore, a lie is not the *definition* of No-Exist; it is a **basis vector**—a fundamental engine that generates non-existence.

## 2. The $N$ -Basis of the Parallax

In linear algebra, a **basis** is a set of independent vectors that span or define a space. If a system has an  $N$ -basis, it means it takes  $N$  distinct, non-overlapping dimensions to fully describe it.

The Parallax of Void=No-Exist is generated by multiple distinct basis vectors ( $N$ -basis) that **do not blanket (overlap) each other**:

The Parallax of Void=No-Exist is generated by multiple distinct basis vectors ( $\mathcal{N}$ -basis) that **do not blanket (overlap) each other**:

1. **The Lie ( $\Lambda_{\text{lie}}$ ):** The **generative act** of untruth that constructs non-existent structures.
2. **Infinity No-Exist ( $\Omega_{\infty}$ ):** The **active, surrounding void** (not a static blankness) that engulfs Existence ( $\mathbb{E}$ ).
3. **Boundary Perturbation / Exaggeration ( $\Omega \mid \epsilon$ ):** Taking a real element and stretching it beyond its valid bounds until it becomes an exaggeration (a lie).
4. **Supposition ( $\mathcal{S}$ ):** The act of assuming or hypothesizing something into conceptual space without ontological grounding.

Because these vectors do not overlap, Void/No-Exist cannot be reduced to a single simple definition. It must be understood as the total space spanned by these independent forces.

### 3. Exaggeration ( $\epsilon$ ) and the Boundary of Existence

In calculus and real analysis,  $\epsilon$  (epsilon) represents an infinitely small variation or error bound.

When the text notes that "*An exaggeration is a lie*" and points to  $\epsilon$ , it highlights how exaggeration works mathematically:

- If you take a true fact  $x \in \mathbb{E}$  and scale it by an uncontrolled magnitude ( $\epsilon$ ), you push  $x$  beyond the boundary of truth ( $\partial\mathbb{E}$ ).
- At that point, it ceases to be a simple measurement error within existence and turns into an untruth—a lie—shifting it into the realm of No-Exist.

### 4. Infinity No-Exist: Not a "Black Blank"

People often picture "nothingness" or "the void" as a dark, static, empty space—a "black blank."

Chapter 5 rejects this view entirely:

- **Infinity No-Exist** is an active, engulfing boundary that surrounds Existence ( $\mathbb{E}$ ).
- It operates across **No-Time | Time**, meaning it exists orthogonally (at a right angle) to the flow of physical time. Time moves within Existence, while Void wraps around the entire temporal manifold.

5. Discordance in Harmonic String Theory ( $\Omega \mid \epsilon$ )

In physics and harmonic analysis, systems achieve balance through resonance and harmony (orderly wave functions operating on specific frequencies).

**Discordance** represents total non-resonance. In the context of String Theory and the Parallax of Void:

- While Existence ( $\mathbb{E}$ ) is structured by harmonic resonance, the Void introduces phase mismatches and non-harmonic perturbations ( $\Omega \mid \epsilon$ ).
- It is this very discordance that keeps Void from collapsing into standard existence, keeping it as an active outer boundary rather than a physical substance inside the universe.

Core Takeaways

| Concept                  | Traditional Meaning              | Parallax / Chapter 5 Meaning                                              |
|--------------------------|----------------------------------|---------------------------------------------------------------------------|
| Falsity ( $\perp$ )      | A false statement about reality. | Located entirely <b>inside</b> Existence ( $\mathbb{E}$ ).                |
| The Lie                  | A untrue statement.              | An active <b>basis vector</b> ( $b_1$ ) that generates No-Exist.          |
| The Void ( $\emptyset$ ) | Empty space / Nothingness.       | An active $N$ -dimensional space engulfing Existence.                     |
| Infinity No-Exist        | Endless empty space.             | The outer boundary condition around time and reality; not a blank vacuum. |
| Supposition              | A guess or hypothesis.           | A projection mapping that acts as another basis of Void.                  |



## Problem Sets

### Problem 1: Foundations of Parallax Measure Spaces

Let  $\mathcal{X}$  be a smooth manifold representing  $\mathcal{E}$ -Existence. Define a non-archimedean measure space  $(\mathcal{X}, \Sigma_{\mathcal{N}}, \mu_{\mathcal{Q}})$  where the sigma-algebra  $\Sigma_{\mathcal{N}}$  contains the empty-set boundary  $\emptyset_X$ . Define a non-contextual operator  $\mathfrak{L}: L^2(\mathcal{X}) \rightarrow L^2(\mathcal{X})$  representing the algebraic structure of a "Lie" as an asymmetric derivation satisfying  $\mathfrak{L}(fg) = f\mathfrak{L}(g) + \mathfrak{L}(f)g + \delta_N(f, g)$ , where  $\delta_N$  is a boundary defect operator supported strictly on  $\emptyset_X$ .

Prove that if  $\text{Supp}(\delta_N) \subseteq \emptyset_X$ , the operator  $\mathfrak{L}$  cannot be closed in the norm topology of  $L^2(\mathcal{X})$ , and derive the spectrum  $\sigma(\mathfrak{L})$  as a dense subset of the complex plane  $\mathbb{C}$ .

### Problem 2: Topological Structure of Boundary Defect Decompositions

Let  $\Omega$  represent the unbounded state domain and  $\varepsilon$  represent bounded physical local reality. Define the  $N$ -basis decomposition of the Parallax space  $\mathcal{P} = \bigoplus_{k=1}^N \mathcal{B}_k$ , where  $\mathcal{B}_1 = \text{Lie}$ ,  $\mathcal{B}_2 = \text{Infinity No-Exist}$ , and  $\mathcal{B}_3 = \text{Supposition}$ .

Construct a sheaf of topological vector spaces  $\mathcal{F}_{\mathcal{Q}}$  over  $\mathcal{P}$  such that the global section cohomology  $H^k(\mathcal{P}, \mathcal{F}_{\mathcal{Q}}) \neq 0$  for all  $k \geq 1$ . Show that the sheaf cohomology does not collapse under standard spectral sequences when  $\mathcal{B}_2$  acts as a non-compact boundary engulfing  $\mathcal{E}$ .

### Problem 3: Non-Contextual Lie Operators and Non-Closed Derivations

Let  $\mathcal{A}$  be a  $C^*$ -algebra equipped with a dynamic automorphism  $\alpha_t$ . Define the Lie basis operator  $\mathfrak{L}_{\text{lie}}$  as an unbonded derivation on  $\mathcal{A}$  with domain  $\mathcal{D}(\mathfrak{L}_{\text{lie}})$ . Show that if the Lie basis preserves no non-trivial state  $\omega \in \mathcal{S}(\mathcal{A})$  such that  $\omega(\mathfrak{L}_{\text{lie}}(a)) = 0$  for all  $a \in \mathcal{D}(\mathfrak{L}_{\text{lie}})$ , then  $\mathcal{A}$  contains no KMS (Kubo-Martin-Schwinger) states at any temperature  $\beta \in (0, \infty)$ , implying the absolute absence of thermal equilibrium in  $\mathcal{E}$ .

#### Problem 4: Discordance Operators in Non-Harmonic String Spectra

Consider a modified Hilbert space  $\mathcal{H}_{\text{string}} = L^2(\mathbb{S}^1) \otimes \ell^2(\mathbb{N})$  equipped with the string Hamiltonian  $H_0 = \sum_{n=1}^{\infty} n a_n^\dagger a_n$ . Define the Discordance Operator  $D_\Omega: \mathcal{H}_{\text{string}} \rightarrow \mathcal{H}_{\text{string}}$  corresponding to the basis  $\Omega \mid \epsilon$  by  $D_\Omega = \sum_{n=1}^{\infty} \lambda_n (a_n^\dagger a_n)^2$ , where  $\lambda_n \sim e^{\gamma n}$  for  $\gamma > 0$ .

Prove that  $H = H_0 + D_\Omega$  has a purely continuous spectrum on  $[0, \infty)$  and possesses zero bound states, demonstrating complete string harmonic destruction.

#### Problem 5: Measure-Theoretic Analysis of Infinity No-Exist

Let  $(X, \mathcal{M}, \nu)$  be a finite measure space representing  $\mathcal{G}$ . Let  $V_\infty$  represent the Infinity No-Exist domain. We construct the product measure space  $(X \times V_\infty, \mathcal{M} \otimes \mathcal{N}_V, \nu \times \mu_\infty)$  where  $\mu_\infty$  is a infinite singular non-Radon measure.

Prove that if  $\mu_\infty(V_\infty) = \infty$  and  $\mu_\infty(A) \in \{0, \infty\}$  for every measurable set  $A \in \mathcal{N}_V$ , the projection map  $\pi_X: X \times V_\infty \rightarrow X$  destroys all  $L^1(X, \nu)$  convergence structures under Radon-Nikodym differentiation.

#### Problem 6: Parallax Basis Projections and Non-Blank Mappings

Let  $\mathcal{H}$  be a separable Hilbert space. Let  $P_\epsilon$  and  $P_\Omega$  be orthogonal projections onto sub-spaces corresponding to local reality  $\epsilon$  and true existence  $\Omega$ . Define the non-blank parallax mapping  $\Phi: \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  by  $\Phi(T) = P_\epsilon T P_\Omega + P_\Omega T P_\epsilon$ .

Prove that  $\Phi$  is an isometry in the Hilbert-Schmidt norm if and only if  $P_\epsilon + P_\Omega = I$  and  $P_\epsilon P_\Omega = 0$ , but fails to be a trace-class operator for any non-zero compact operator  $T$ .

#### Problem 7: Cohomological Null Boundaries on $\emptyset_X$

Let  $M$  be a smooth  $d$ -dimensional compact manifold with boundary  $\partial M = \emptyset_X$ . Define the de Rham complex  $(\Omega^*(M), d)$  with modified boundary conditions  $d\omega|_{\partial M} = \mathfrak{L}_{\text{lie}}(\omega|_{\partial M})$ .

Calculate the relative cohomology groups  $H^k(M, \emptyset_X)$  and establish under what conditions  $\dim H^k(M, \emptyset_X) = \infty$  for intermediate dimensions  $1 < k < d - 1$ .

### Problem 8: Spectral Discontinuity of the Supposition Basis

Let  $\mathcal{S}$  be the Supposition Operator defined on Sobolev space  $H^s(\mathbb{R}^n)$  as  $\mathcal{S}f(x) = f(x + \theta(x))$ , where  $\theta(x) \in C_0^\infty(\mathbb{R}^n)$  is a non-deterministic perturbation field.

Prove that for any  $s > n/2$ , the map  $\theta \mapsto \mathcal{S}$  is nowhere continuous in the operator norm topology  $\mathcal{B}(H^s(\mathbb{R}^n))$ .

### Problem 9: Dynamic Parallax Transformations in No-Time Domain

Consider the time-independent operator system defined on a  $C^*$ -algebra  $\mathcal{A}$  without a primary parameter  $t$ . Let  $\Delta_{\text{Parallax}}$  be a linear transformation defined by  $\Delta_{\text{Parallax}}(A) = [D_\Omega, A] + \mathfrak{L}(A)$ .

Show that if  $[D_\Omega, \mathfrak{L}] \neq 0$ , the generator  $\Delta_{\text{Parallax}}$  induces a non-strongly continuous semigroup  $e^{\tau \Delta_{\text{Parallax}}}$  on  $\mathcal{A}$  for parameter  $\tau \in \mathbb{C} \setminus \mathbb{R}$ .

### Problem 10: Non-Archimedean Valuation of Epsilon Over-Exaggeration

Let  $\mathbb{K}$  be a non-Archimedean field with valuation  $|\cdot|_v$ . Define an exaggeration mapping  $\mathcal{E}_{ex} : \mathbb{K} \rightarrow \mathbb{K}$  by  $\mathcal{E}_{ex}(x) = x \cdot |x|_v^{-\alpha}$  for  $\alpha > 1$ .

Show that the topological closure of the range  $\overline{\text{Range}(\mathcal{E}_{ex})}$  is nowhere dense in  $\mathbb{K}$  and compute its Hausdorff dimension with respect to the non-Archimedean metric  $d(x, y) = |x - y|_v$ .

### Problem 11: Ergodic Singularities of Falsity vs. Lie Operators

Let  $(T, X, \mu)$  be an ergodic dynamical system. Define the Falsity operator  $F : L^2(X) \rightarrow L^2(X)$  by  $Ff(x) = f(Tx)$  and the Lie operator  $\mathfrak{L} : L^2(X) \rightarrow L^2(X)$  as a non-commuting multiplier operator  $\mathfrak{L}f(x) = g(x)f(x)$  where  $g \notin L^\infty(X)$ .

Prove that the commutator  $[F, \mathfrak{L}]$  has a dense image in  $L^2(X)$  if and only if  $g(x) - g(Tx) \neq 0$  almost everywhere.

### Problem 12: Boundary Defect Quantization in String Superpositions

In a string field theory framework, let  $|\Psi\rangle$  be a string field state in a Hilbert space  $\mathcal{H}_{\text{SFT}}$ . Define the Parallax defect action  $S_{\text{defect}} = \langle \Psi | \{ [Q_{\text{BRST}}, D_{\Omega}] | \Psi \rangle$ .

Calculate the BRST cohomology of the operator  $Q_{\text{BRST}} + D_{\Omega}$  and prove that all physical states are projected out, leaving an unphysical (Void) spectrum.

### Problem 13: Infinite Basis Multi-Vector Extensions of $\mathcal{N}$ -Basis

Let  $\mathcal{V}$  be a real vector space of dimension  $\aleph_0$  with basis  $\mathcal{B}_{\mathcal{N}} = \{b_1, b_2, \dots\}$ . Construct the exterior algebra  $\bigwedge^*(\mathcal{V})$ . Define the infinite-dimensional Parallax forms  $\omega = \sum_{i=1}^{\infty} b_i \wedge b_{i+1}$ .

Determine whether  $\omega \wedge \omega = 0$  and classify the kernel of the interior product operator  $i_X \omega$  for arbitrary vectors  $X \in \mathcal{V}$ .

### Problem 14: Non-Contextual Harmonic Resolvents

Let  $A$  be a self-adjoint operator on a Hilbert space  $\mathcal{H}$  with spectrum  $\sigma(A) \subset [0, \infty)$ . Define the Parallax Resolvent  $R_P(z) = (A - zI + D_{\Omega})^{-1}$  where  $D_{\Omega}$  is the Discordance operator from Problem 4.

Prove that for  $z \in \mathbb{C} \setminus \mathbb{R}$ ,  $R_P(z)$  is non-compact and compute its Schatten-p norm bounds.

### Problem 15: Structural Instability of the Omega/Epsilon Boundary

Let  $M$  be a Riemannian manifold with metric  $g_{\epsilon}$ . Consider a sequence of metrics  $g_k = g_{\epsilon} + k \cdot h$  where  $h$  is a symmetric 2-tensor supported near the boundary  $\partial M = \emptyset_X$ .

Prove that as  $k \rightarrow \infty$ , the sequence of sectional curvatures  $K_{g_k}$  diverges to  $-\infty$  everywhere in the support of  $h$ , signaling a metric breakdown into Infinity No-Exist.

### Problem 16: Micro-Local Analysis of Supposition Singularities

Let  $u \in \mathcal{D}'(\mathbb{R}^n)$  be a distribution with wave-front set  $\text{WF}(u)$ . Let  $\mathcal{S}$  be the Supposition operator defined in Problem 8.

Compute the wavefront set  $\text{WF}(\mathcal{S}u)$  in terms of  $\text{WF}(u)$  and the gradient matrix  $\nabla\theta(x)$ , and determine the condition under which the wavefront set expands to the entire cotangent bundle  $T^*\mathbb{R}^n \setminus \{0\}$ .

### Problem 17: Non-Abelian Parallax Gauge Theories

Let  $A$  be a connection 1-form on a principal  $G$ -bundle over  $M$  with curvature  $F = dA + A \wedge A$ . Define the Parallax Gauge Transformation  $A \mapsto A' = gAg^{-1} + gdg^{-1} + \mathfrak{L}(g)$ , where  $\mathfrak{L}$  is the non-derivational Lie operator.

Calculate the transformed curvature  $F'$  and show that the gauge invariance of the Yang-Mills action  $S(A) = \int_M \text{Tr}(F \wedge *F)$  is explicitly broken by a boundary trace term.

### Problem 18: Operator Algebras of Void State Extensions

Let  $\mathcal{M}$  be a von Neumann algebra of type  $\text{III}_1$ . Define a projection operator  $P_{\emptyset_X} \notin \mathcal{M}$  acting on an extended Hilbert space  $\mathcal{H} \oplus \mathcal{H}_\infty$ .

Prove that the generated algebra  $\mathcal{N} = \text{vN}(\mathcal{M}, P_{\emptyset_X})$  is a factor of type  $\text{III}_0$  or non-factorable, and construct its modular automorphism group  $\sigma_t^\phi$ .

### Problem 19: Topological Entropy of Discordant Parallax Systems

Let  $(X, T)$  be a topological dynamical system. Define the discordance perturbation  $T_D(x) = T(x) + D(x) \pmod{1}$  on the  $d$ -torus  $\mathbb{T}^d$ .

If  $D(x)$  is a discontinuous piecewise constant function representing state jumps, show that the topological entropy  $h_{\text{top}}(T_D)$  satisfies  $h_{\text{top}}(T_D) = \infty$ , proving maximum chaotic distribution under Void dynamics.

## Problem 20: Axiomatic Incompleteness of the Parallax System

Let  $\mathcal{T}_{\text{Parallax}}$  be a formal first-order axiomatic theory containing arithmetic and the operators  $\{\mathfrak{L}, D_{\Omega}, \mathcal{S}, \emptyset_X\}$ .

Prove, by constructing a Gödel sentence  $G_{\text{Void}}$ , that  $\mathcal{T}_{\text{Parallax}}$  cannot prove its own consistency unless the system is inconsistent, thereby mathematically confirming that "Supposition is another basis of Void=No-Exist."

## Solutions

### Solution 1

To prove  $\mathfrak{L}$  is not closed, consider a sequence  $f_n \in C_0^\infty(\mathcal{X})$  converging to  $f \in L^2(\mathcal{X})$  in  $L^2$ -norm, such that  $f_n$  restricted to  $\emptyset_X$  has unbounded derivatives  $\nabla f_n|_{\emptyset_X} \rightarrow \infty$ .

Because  $\mathfrak{L}(f_n) = f_n \mathfrak{L}(1) + \delta_N(f_n, 1)$ , the presence of the boundary defect  $\delta_N$  supported on the null boundary  $\emptyset_X$  forces  $\|\mathfrak{L}(f_n)\|_{L^2} \rightarrow \infty$  even as  $\|f_n\|_{L^2} \rightarrow 0$ . Thus, the graph of  $\mathfrak{L}$  is not closed in  $L^2(\mathcal{X}) \times L^2(\mathcal{X})$ .

The spectrum  $\sigma(\mathfrak{L})$  contains every  $\lambda \in \mathbb{C}$  because the operator  $(\mathfrak{L} - \lambda I)$  fails to have a bounded inverse on the singular boundary subspace  $\emptyset_X$ , making  $\sigma(\mathfrak{L}) = \mathbb{C}$ .

### Solution 2

Construct the sheaf  $\mathcal{F}_{\emptyset}$  over  $\mathcal{P}$  by associating to each open set  $U \subset \mathcal{P}$  the space of smooth functions with non-vanishing boundary conditions on  $\mathcal{B}_2 = \text{Infinity No-Exist}$ .

The long exact sequence in cohomology yields  $\dots \rightarrow H^k(\mathcal{P}) \rightarrow H^k(\mathcal{P} \setminus \mathcal{B}_2) \rightarrow H_{\mathcal{B}_2}^{k+1}(\mathcal{P}) \rightarrow \dots$



Since  $\mathcal{B}_2$  acts as a non-compact engulfing boundary,  $H_{\mathcal{B}_2}^{k+1}(\mathcal{P})$  is non-trivial for all  $k \geq 0$ . The spectral sequence  $E_2^{p,q} = H^p(\mathcal{P}, \mathcal{H}^q(\mathcal{F}_\emptyset)) \Rightarrow H^{p+q}(\mathcal{P}, \mathcal{F}_\emptyset)$  has non-zero differentials  $d_r$  for infinitely many  $r$ , preventing collapse and ensuring  $H^k(\mathcal{P}, \mathcal{F}_\emptyset) \neq 0$  for all  $k \geq 1$ .

### Solution 3

Suppose  $\omega$  is a KMS state at inverse temperature  $\beta > 0$  for the automorphism group generated by  $\mathfrak{L}_{\text{lie}}$ . The KMS condition requires that for all  $a, b \in \mathcal{D}(\mathfrak{L}_{\text{lie}})$ ,  $\omega(a\alpha_{i\beta}(b)) = \omega(ba)$ .

Taking the derivative at  $t = 0$  yields  $\omega(a\mathfrak{L}_{\text{lie}}(b)) = 0$ . Setting  $a = I$ , we get  $\omega(\mathfrak{L}_{\text{lie}}(b)) = 0$  for all  $b$ .

However, by hypothesis, no non-trivial state satisfies this invariant identity due to the non-closed derivation structure of the Lie basis. Thus, no such KMS state  $\omega$  can exist for any  $\beta \in (0, \infty)$ .

### Solution 4

The total operator  $H = H_0 + D_\Omega$  acts on  $\mathcal{H}_{\text{string}}$ . Since  $\lambda_n \sim e^{\gamma n}$ , the perturbation term  $D_\Omega = \sum \lambda_n (a_n^\dagger a_n)^2$  is unbounded and dominates  $H_0$  for all modes  $n \geq 1$ .

The resolvent  $R(z) = (H - zI)^{-1}$  has no point spectrum because any prospective eigenvector  $\psi = \sum c_n |\{n\}\rangle$  requires  $\sum |\{n\}c_n|^2 \lambda_n^2 < \infty$ , which forces  $c_n = 0$  for all  $n$  due to exponential growth of  $\lambda_n$ .

By the Birman-Schwinger principle, the operator spectrum collapses into a purely continuous spectrum on  $[0, \infty)$ , ruling out bound states.

### Solution 5

Consider the Radon-Nikodym derivative  $g = \frac{d(\nu \times \mu_\infty)}{d\lambda}$  relative to a reference Lebesgue measure  $\lambda$ . For any  $f \in L^1(X, \nu)$ , the integrated form over a measurable set  $E \times A \in \mathcal{M} \otimes \mathcal{N}_V$  becomes:

$$\int_{E \times A} f(x) d(\nu \times \mu_\infty) = \mu_\infty(A) \int_E f(x) d\nu$$

Since  $\mu_\infty(A) \in \{0, \infty\}$ , this integral evaluates strictly to 0 or  $\infty$ .

Consequently, no finite integrable function  $g \in L^1(X \times V_\infty)$  exists to satisfy the Radon-Nikodym theorem, invalidating  $L^1$  convergence under projection  $\pi_X$ .

### Solution 6

Calculate the Hilbert-Schmidt norm  $\|\Phi(T)\|_{\text{HS}}^2 = \text{Tr}(\Phi(T)^\dagger \Phi(T))$ .

Expanding:

$$\Phi(T)^\dagger \Phi(T) = (P_\Omega T^\dagger P_\epsilon + P_\epsilon T^\dagger P_\Omega)(P_\epsilon T P_\Omega + P_\Omega T P_\epsilon) = P_\Omega T^\dagger P_\epsilon T P_\Omega + P_\epsilon T^\dagger P_\Omega T P_\epsilon$$

Taking the trace yields  $\|P_\epsilon T P_\Omega\|_{\text{HS}}^2 + \|P_\Omega T P_\epsilon\|_{\text{HS}}^2 = \|\Phi(T)\|_{\text{HS}}^2$ , which holds for all  $T$  if and only if  $P_\epsilon + P_\Omega = I$  and  $P_\epsilon P_\Omega = 0$ .



However, for a non-zero compact operator  $T$ , the trace-class norm satisfies  $\|\Phi(T)\|_1 = \text{Tr}(|\Phi(T)|)$ . Since  $\Phi(T)$  has off-diagonal block structure with respect to  $\text{Im}(P_\epsilon) \oplus \text{Im}(P_\Omega)$ , its singular values do not sum to a finite value if  $P_\Omega$  is infinite-dimensional, making  $\Phi(T)$  non-trace-class.

### Solution 7

The modified differential boundary condition  $d\omega|_{\partial X} = \mathfrak{L}_{\text{lie}}(\omega|_{\partial X})$  replaces standard exterior differentiation at the boundary with an unbounded non-derational operator.

Applying the long exact sequence for pair cohomology:

$$\dots \rightarrow H^k(M, \partial X) \rightarrow H^k(M) \rightarrow H^k(\partial X) \rightarrow H^{k+1}(M, \partial X) \dots$$

Because  $\mathfrak{L}_{\text{lie}}$  introduces an infinite-dimensional kernel and range over  $C^\infty(\partial X)$ , the boundary cohomology  $H^k(\partial X)$  becomes infinite-dimensional.

By exactness,  $\dim H^k(M, \partial X) \geq \dim \text{coker}(H^{k-1}(M) \rightarrow H^{k-1}(\partial X)) = \infty$  for all  $1 < k < d - 1$ .

### Solution 8

Let  $f \in H^s(\mathbb{R}^n)$  with  $s > n/2$ . Consider the sequence of smooth vector fields  $\theta_k(x) = \frac{1}{k} \sin(k^2 x_1) \mathbf{e}_1$ . As  $k \rightarrow \infty$ ,  $\|\theta_k\|_{C^0} = \frac{1}{k} \rightarrow 0$ .

However, evaluating the operator norm  $\|\mathcal{S}_k - I\|_{\mathcal{B}(H^s)}$  on high-frequency testing functions  $f_\lambda(x) = e^{i\lambda x_1} e^{-|x|^2}$  with  $\lambda \sim k^2$ :

$$\|\{\!\!\{ \mathcal{S}_k f_\lambda - f_\lambda \}\!\!\|_{H^s} \sim k^{2s} \|\{\!\!\{ \theta_k \}\!\!\|_{C^0} = k^{2s-1} \rightarrow \infty \quad \text{for } s > 1/2$$

Since  $s > n/2 \geq 1/2$ , the map  $\theta \mapsto \mathcal{S}$  is discontinuous everywhere in the operator norm topology.

#### Solution 9

The operator  $\Delta_{\text{Parallax}} = \text{ad}_{D_\Omega} + \mathfrak{L}$  generates a formal perturbation series.

The commutator  $[D_\Omega, \mathfrak{L}] \neq 0$  implies that the operators do not commute on the domain  $\mathcal{D}(D_\Omega) \cap \mathcal{D}(\mathfrak{L})$ . For  $\tau = u + iv \in \mathbb{C} \setminus \mathbb{R}$  with  $v \neq 0$ , the operator norm satisfies:

$$\|\{\!\!\{ e^{\tau \Delta_{\text{Parallax}}} \}\!\!\|_{\{\!\!\{ \}\!\!\}} \geq e^{|v| \cdot \|[D_\Omega, \mathfrak{L}]\|}$$

Since  $[D_\Omega, \mathfrak{L}]$  is unbounded due to the continuous spectral properties of  $D_\Omega$ ,  $\|\{\!\!\{ e^{\tau \Delta_{\text{Parallax}}} \}\!\!\|_{\{\!\!\{ \}\!\!\}} = \infty$  for all  $v \neq 0$ .

Hence,  $e^{\tau \Delta_{\text{Parallax}}}$  fails to satisfy strong continuity  $\lim_{\tau \rightarrow 0} \|\{\!\!\{ e^{\tau \Delta_{\text{Parallax}}} A - A \}\!\!\|_{\{\!\!\{ \}\!\!\}} = 0$ .

#### Solution 10

Let  $x \in \mathbb{K}$ . The non-Archimedean valuation satisfies  $|\{\!\!\{ x \cdot y \}\!\!\|_v = |\{\!\!\{ x \}\!\!\|_v |\{\!\!\{ y \}\!\!\|_v$ . The valuation of the exaggerated image is:

$$|\{\!\!\{ \mathcal{E}_{ex}(x) \}\!\!\|_v = \left| x \cdot |\{\!\!\{ x \}\!\!\|_v^{-\alpha} \right|_v = |\{\!\!\{ x \}\!\!\|_v \cdot |\{\!\!\{ x \}\!\!\|_v^{-\alpha} = |\{\!\!\{ x \}\!\!\|_v^{1-\alpha}$$

Since  $\alpha > 1$ ,  $1 - \alpha < 0$ . As  $|\{\!\!\{ x \}\!\!\|_v \rightarrow 0$ ,  $|\{\!\!\{ \mathcal{E}_{ex}(x) \}\!\!\|_v \rightarrow \infty$ , and as  $|\{\!\!\{ x \}\!\!\|_v \rightarrow \infty$ ,  $|\{\!\!\{ \mathcal{E}_{ex}(x) \}\!\!\|_v \rightarrow 0$ .

### Solution 10

Let  $x \in \mathbb{K}$ . The non-Archimedean valuation satisfies  $|\{x \cdot y\}_v| = |\{x\}_v| |\{y\}_v|$ . The valuation of the exaggerated image is:

$$|\{\mathcal{E}_{ex}(x)\}_v| = |x \cdot |\{x\}_v|^{-\alpha}|_v = |\{x\}_v| \cdot |\{x\}_v|^{-\alpha} = |\{x\}_v|^{1-\alpha}$$

Since  $\alpha > 1$ ,  $1 - \alpha < 0$ . As  $|\{x\}_v| \rightarrow 0$ ,  $|\{\mathcal{E}_{ex}(x)\}_v| \rightarrow \infty$ , and as  $|\{x\}_v| \rightarrow \infty$ ,  $|\{\mathcal{E}_{ex}(x)\}_v| \rightarrow 0$ .

The image values form a discrete set of valuation spheres  $\{p^{k(\alpha-1)} : k \in \mathbb{Z}\}$  where  $p$  is the uniformizer of  $\mathbb{K}$ .

Because this set consists of isolated boundary spheres in the ultrametric topology, its closure  $\overline{\text{Range}(\mathcal{E}_{ex})}$  contains no open balls, making it nowhere dense. Its Hausdorff dimension relative to  $d(x, y) = |\{x - y\}_v|$  is identically 0.

### Solution 11

The commutator  $[F, \mathfrak{L}]$  applied to  $f \in L^2(X)$  is:

$$[F, \mathfrak{L}]f(x) = F(g(x)f(x)) - g(x)F(f(x)) \underset{f(Tx)}{=} g(Tx)f(Tx) - g(x)f(Tx) = (g(Tx) - g(x))f(Tx)$$

This acts as a composite operator  $M_{g \circ T - g} \circ F$ , where  $M_{g \circ T - g}$  is a multiplication operator.

Because  $F$  is an isometry on  $L^2(X)$  due to ergodicity and measure preservation of  $T$ ,  $[F, \mathfrak{L}]$  has a dense image if and only if  $M_{g \circ T - g}$  has a dense image.

The multiplication operator  $M_h$  has a dense image in  $L^2(X)$  if and only if  $h(x) \neq 0$  almost everywhere. Setting  $h(x) = g(Tx) - g(x)$ , the condition holds if and only if  $g(x) - g(Tx) \neq 0$  a.e.

### Solution 12

The total state operator is  $Q_{\text{tot}} = Q_{\text{BRST}} + D_{\Omega}$ .

The BRST operator satisfies  $Q_{\text{BRST}}^2 = 0$ , but the Discordance operator satisfies  $D_{\Omega}^2 \neq 0$  and  $\{Q_{\text{BRST}}, D_{\Omega}\} = \Lambda \neq 0$ . Compute  $Q_{\text{tot}}^2$ :

$$Q_{\text{tot}}^2 = (Q_{\text{BRST}} + D_{\Omega})^2 = Q_{\text{BRST}}^2 + \{Q_{\text{BRST}}, D_{\Omega}\} + D_{\Omega}^2 = \Lambda + D_{\Omega}^2 \neq 0$$

Because  $Q_{\text{tot}}^2 \neq 0$ , the nilpotent requirement for defining a physical cohomology  $H^* = \text{Ker}(Q_{\text{tot}})/\text{Im}(Q_{\text{tot}})$  fails.

Thus,  $\text{Ker}(Q_{\text{tot}}) = \{0\}$ , rendering the BRST physical cohomology completely trivial ( $\dim H^* = 0$ ). All physical states are projected out.

### Solution 13

Calculate  $\omega \wedge \omega$ :

$$\omega \wedge \omega = \left( \sum_{i=1}^{\infty} b_i \wedge b_{i+1} \right) \wedge \left( \sum_{j=1}^{\infty} b_j \wedge b_{j+1} \right) = \sum_{i,j=1}^{\infty} b_i \wedge b_{i+1} \wedge b_j \wedge b_{j+1}$$

For  $|i - j| \geq 2$ , terms like  $b_i \wedge b_{i+1} \wedge b_j \wedge b_{j+1} \neq 0$ , so  $\omega \wedge \omega \neq 0$ .

To find  $\text{Ker}(i_X \omega)$ , write  $X = \sum c_k b_k$ . The interior product gives:

$$i_X \omega = \sum_{k=1}^{\infty} (c_k b_{k+1} - c_{k+1} b_k)$$

Setting  $i_X \omega = 0$  requires  $c_k b_{k+1} - c_{k+1} b_k = 0$  for all  $k$ . Since  $\{b_k\}$  are linearly independent, this forces  $c_k = 0$  for all  $k$ .

Thus,  $\text{Ker}(i_X \omega) = \{0\}$ .

### Solution 14

The Parallax Resolvent is  $R_P(z) = (A - zI + D_{\Omega})^{-1}$ .

From Problem 4,  $D_{\Omega} = \sum \lambda_n (a_n^{\dagger} a_n)^2$  with  $\lambda_n \sim e^{\gamma n}$ . The operator  $A - zI + D_{\Omega}$  is unbounded with eigenvalues growing at least exponentially.

Its inverse  $R_P(z)$  has eigenvalues  $\mu_n = (\alpha_n - z + \lambda_n)^{-1} \sim e^{-\gamma n}$ .

Compute the Schatten- $p$  norm  $\| \mathfrak{I} R_p(z) \|_p^p = \sum_{n=1}^{\infty} |\mathfrak{I} \mu_n|^p \sim \sum_{n=1}^{\infty} e^{-p\gamma n} = \frac{e^{-p\gamma}}{1-e^{-p\gamma}} < \infty$  for all  $p > 0$ .

Thus,  $R_p(z)$  is compact and belongs to every Schatten class  $\mathcal{S}_p$  for  $p > 0$ .

### Solution 15

Let  $g_k = g_e + k \cdot h$ . In local coordinates  $(x^1, \dots, x^d)$  where  $h_{ij} \neq 0$ , the Riemann curvature tensor components are given by:

$$R^l_{ijk} = \partial_j \Gamma^l_{ik} - \partial_k \Gamma^l_{ij} + \Gamma^l_{jm} \Gamma^m_{ik} - \Gamma^l_{km} \Gamma^m_{ij}$$

The Christoffel symbols scale as  $\Gamma^l_{ij}(g_k) \sim \mathcal{O}(1)$  as  $k \rightarrow \infty$ , but their derivatives yield terms of order  $\mathcal{O}(k)$ .

The sectional curvature  $K(u, v) = \frac{g_k(R(u,v)v,u)}{g_k(u,u)g_k(v,v)-g_k(u,v)^2}$  evaluates to:

$$K_{g_k} \sim \frac{-k^2(\partial h)^2}{k^2 h^2} \sim -C \cdot k \rightarrow -\infty \quad \text{as } k \rightarrow \infty$$

Everywhere in  $\text{Supp}(h)$ , sectional curvature diverges to negative infinity, breaking metric smoothness into the engulfing Void.

### Solution 16

The wavefront set  $\text{WF}(u) \subset T^*\mathbb{R}^n \setminus \{0\}$  tracks singularity locations and directions.

For the Supposition transformation  $\mathcal{S}u(x) = u(x + \theta(x))$ , let  $y = \Phi(x) = x + \theta(x)$ . The pullback formula for distributions yields:

$$\text{WF}(\mathcal{S}u) = \Phi^* \text{WF}(u) = \{ (x, (I + \nabla\theta(x))^T \xi) \mid (x + \theta(x), \xi) \in \text{WF}(u) \}$$

If  $\det(I + \nabla\theta(x)) = 0$  at singular points  $x_0$ , the transformation matrix  $(I + \nabla\theta(x))^T$  becomes singular, causing the covector directions  $\xi$  to map to zero or collapse.

When  $\nabla\theta(x)$  has dense critical points where  $\text{Rank}(\nabla\theta(x))$  drops maximally, the directions  $(I + \nabla\theta(x))^T \xi$  sweep out all directions in  $\mathbb{R}^n \setminus \{0\}$ , expanding  $\text{WF}(\mathcal{S}u)$  to the full cotangent bundle  $T^*\mathbb{R}^n \setminus \{0\}$ .

### Solution 17

Under  $A \mapsto A' = gAg^{-1} + gdg^{-1} + \mathfrak{L}(g)$ , compute the field strength  $F' = dA' + A' \wedge A'$ :

$$F' = d(gAg^{-1} + gdg^{-1} + \mathfrak{L}(g)) + (gAg^{-1} + gdg^{-1} + \mathfrak{L}(g)) \wedge (gAg^{-1} + gdg^{-1} + \mathfrak{L}(g))$$

Using  $d(gAg^{-1}) = dgAg^{-1} + g dAg^{-1} - gA \wedge dg^{-1}$ :

$$F' = gFg^{-1} + d(\mathfrak{L}(g)) + [\mathfrak{L}(g), gAg^{-1} + gdg^{-1}] + \mathfrak{L}(g) \wedge \mathfrak{L}(g)$$

Let  $\Theta_{\mathfrak{g}} = d(\mathfrak{L}(g)) + [\mathfrak{L}(g), A'] + \mathfrak{L}(g) \wedge \mathfrak{L}(g)$ . Then  $F' = gFg^{-1} + \Theta_{\mathfrak{g}}$ .

The Yang-Mills action becomes:

$$S(A') = \int_M \text{Tr}(F' \wedge * F') = \int_M \text{Tr}(F \wedge * F) + 2 \int_M \text{Tr}(gFg^{-1} \wedge * \Theta_{\mathfrak{g}}) + \int_M \text{Tr}(\Theta_{\mathfrak{g}} \wedge * \Theta_{\mathfrak{g}})$$

The additional non-zero trace terms explicitly break the gauge invariance of  $S(A)$ .

### Solution 18

Let  $\mathcal{M} \subset \mathcal{B}(\mathcal{H})$  be a type  $\text{III}_1$  factor, and  $P_{\mathcal{H}_{\infty}}$  be a projection onto  $\mathcal{H}_{\infty}$ .

Since  $P_{\mathcal{H}_{\infty}} \notin \mathcal{M}$  and  $\mathcal{H}_{\infty}$  is decoupled from the modular flow of  $\mathcal{M}$ , the extension  $\mathcal{N} = \text{vN}(\mathcal{M}, P_{\mathcal{H}_{\infty}})$  breaks the central homogeneity of  $\mathcal{M}$ .

The center  $Z(\mathcal{N}) = \mathcal{N} \cap \mathcal{N}'$  contains non-trivial projections generated by  $P_{\mathcal{H}_{\infty}} \wedge I_{\mathcal{H}}$ , making  $\mathcal{N}$  non-factorable.

Its modular automorphism group  $\sigma_t^{\phi}$  decomposes into a direct sum  $\sigma_t^{\phi, \mathcal{M}} \oplus \text{id}_{\mathcal{H}_{\infty}}$ , scaling the modular flow kernel and shifting the flow spectrum to include an isolated discrete point  $\{0\}$ , rendering the extension a type  $\text{III}_0$  direct integral of factors.

### Solution 19

The topological entropy  $h_{\text{top}}(T_D)$  for a map on  $\mathbb{T}^d$  measures the exponential growth rate of distinct orbits.

Let  $D(x) = \sum_{j=1}^K c_j \chi_{E_j}(x)$  be a piecewise constant jump operator with discontinuity boundaries  $\bigcup \partial E_j$ .



For any  $\epsilon > 0$ , two points  $x, y$  separated by a discontinuity line  $\partial E_j$  are mapped to orbits  $(T_D^n(x), T_D^n(y))$  that diverge instantly by  $|\{c_i - c_j\}|$ .

The number of  $(n, \epsilon)$ -separated sets  $N(n, \epsilon)$  grows as  $K^n$  where  $K$  represents the number of jump domains. Taking the limit:

$$h_{\text{top}}(T_D) = \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \geq \lim_{K \rightarrow \infty} \log K = \infty$$

The discordance perturbation generates infinite topological entropy.

## Solution 20

Let  $\mathcal{T}_{\text{Parallax}}$  be a formal consistent theory containing Peano Arithmetic (PA) and the Void operators  $\{\mathfrak{L}, D_{\Omega}, \mathcal{S}, \emptyset_X\}$ .

By Gödel's First Incompleteness Theorem, we can construct a sentence  $G_{\text{Void}}$  encoding the statement: " $G_{\text{Void}}$  is not provable in  $\mathcal{T}_{\text{Parallax}}$ ."

Within the theory, represent "Supposition" as the structural capacity to assume sentences outside the axiomatic deduction sequence. If  $\mathcal{T}_{\text{Parallax}} \vdash \text{Consis}(\mathcal{T}_{\text{Parallax}})$ , then by Gödel's Second Incompleteness Theorem,  $\mathcal{T}_{\text{Parallax}}$  is inconsistent.

Thus, the formal validity of the theoretical framework relies on non-provable suppositions, proving mathematically that Supposition is a necessary basis for Void=No-Exist.

## CHAPTER 6 - N=BEYOND

Is there something beyond The Parallax of Existence?.

There IS something beyond The Parallax N=Beyond.

That IS the basis of N=Beyond.

Eternity. A Parallax opposed to Infinity.

By now - you think The Parallax N=Beyond reduces to Realms.

That is farther from The Truth.

We barely scratch the surface The Parallax N=Beyond.

The Parallax of Eternity is N=Nothing.

According to The United States Constitution: It says 'Nothing' of Eternity.

It is Timeless. A different concept from No-Time.

S = Timeless (state) {Eternity}.

Reads 'The state of Eternity is Timeless.'

So what can we do? Elementary building blocks - first.

From these proofs alone - we have a framework of The Parallax of Eternity N=Nothing.

**\*NOTE\***

**READ UP ON 'NOTHINGNESS AND BEING' IN EXISTENTIALISM PHILOSOPHY TO  
GET  
A SENSE OF ETERNITY**

Does The Parallax cross-over like a Venn diagram?.

**Can** The Parallax cross-over like a Venn diagram?.

You will have to disprove us if it wasn't.

This is just to get you comfortable with The Graduate Level Math Textbook of Existence.

N=Nothing is NOT null.

N=Nothing is NOT zero.

N=Nothing is NOT not.

We're slowly chipping away what is NOT like a sculptor.

The Parallax of Eternity N=Nothing has no dimensions.

Need more refreshments?.

This is an intro 101 to N=Beyond. More later.

## Problem Sets

### Problem 1 (Set Theory & Measure Theory: Non-Nullity of $N = \text{Nothing}$ )

Let  $(X, \mathcal{M}, \mu)$  be a non-atomic measure space. Let  $\emptyset$  denote the empty set and  $0 \in \mathbb{R}$ . We define  $N = \text{Nothing}$  as a state space object  $\mathcal{N} \in \mathcal{M}$  such that  $\mu(\mathcal{N}) > 0$  despite  $\mathcal{N} \cap A = \emptyset$  for all classic finite partitions  $A$ . Prove that  $\mathcal{N}$  cannot be isomorphic to the empty set  $\emptyset$  or the zero element  $0 \in \mathbb{R}$  in the category of measurable spaces **Meas**.

### Problem 2 (Topology: Dimensionality of Zero-Dimension vs. No-Dimension)

Let  $X$  be a topological space. Lebesgue covering dimension  $\dim(X) = 0$  implies every open cover has a refinement by disjoint open sets. Define an ultra-disconnected state space  $S_{\text{Eternity}}$  such that  $\dim(S_{\text{Eternity}}) = -\infty$  (or undefined under standard Inductive Dimension **ind**). Prove that if  $S_{\text{Eternity}}$  possesses no local neighborhood basis, it cannot embed into any  $\mathbb{R}^n$  for  $n \geq 0$ .

### Problem 3 (Category Theory: Parallax Cross-Over as Pullbacks)

Let  $\mathbf{C}$  be a category representing Existence spaces. Define two objects:  $P_1$  (Infinity) and  $P_2$  (Eternity). Suppose their cross-over is defined by a non-trivial Venn-like intersection modeled as a pullback object  $P_1 \times_X P_2$  over a base space  $X = N = \text{Beyond}$ . Prove that if the pullback object is non-zero, then  $P_1$  and  $P_2$  cannot be orthogonal in  $\mathbf{C}$ .

### Problem 4 (Differential Geometry: Timeless Metric Tensor)

Let  $M = \mathbb{R} \times M_{\text{space}}$  be a pseudo-Riemannian manifold with signature  $(-1, +, +, +)$ . Define the time state  $S_{\text{Timeless}}$  by a degeneracy condition on the metric tensor  $g: g(V, \cdot) = 0$  for all timelike vectors  $V \in TM$ . Show that the resulting manifold degenerates into a fiber bundle over space where the vector field associated with time vanishes identically without making the underlying spatial manifold trivial.

### Problem 5 (Non-Standard Analysis: Infinity vs. Eternity)

In the hyperreal field  ${}^*\mathbb{R}$ , let  $\omega \in {}^*\mathbb{R} \setminus \mathbb{R}$  be an infinite hyperreal number (representing Infinity). Define the state  $S_{\text{Eternity}}$  as the halo/monad  $\mu(0)$  or an infinitesimal hyperfinite quotient  $\mathbb{H}/\omega$ . Prove that  $S_{\text{Eternity}}$  is algebraically non-isomorphic to  $\mathbb{R}/\mathbb{Z}$  and cannot be mapped bijectively to the set of infinite hyperreals  ${}^*\mathbb{R}_\infty$ .

### Problem 6 (Functional Analysis: Spectral Theory of Timeless Operators)

Let  $\mathcal{H}$  be a Hilbert space and  $H$  a self-adjoint Hamiltonian operator generating time evolution via  $U(t) = e^{-iHt/\hbar}$ . Define a state  $\psi \in \mathcal{H}$  to be Timeless ( $S \in \{\text{Eternity}\}$ ) if  $U(t)\psi = \psi$  for all  $t \in \mathbb{R}$ . Prove that  $S$  corresponds strictly to the zero-eigenspace  $\ker(H)$ , and demonstrate that  $\dim(\ker(H))$  can be infinite without violating the boundedness of the resolvent  $R_z(H)$  for  $z \notin \sigma(H)$ .

### Problem 7 (Abstract Algebra: The Structure of $N = \text{Beyond}$ )

Let  $R$  be a non-commutative ring with unity. Define an element  $x \in R$  as "Not-Null" and "Not-Zero" ( $x \neq 0$ ,  $x^n \neq 0$  for any  $n$ ). Let  $I = (x)$  be the two-sided ideal generated by  $x$ . Prove that if  $R/I \cong R$ ,  $R$  must be non-Noetherian and  $x$  cannot be a unit or a classical zero-divisor.

### Problem 8 (Topological Quantum Field Theory: Boundary of Eternity)

Let  $M$  be a  $d$ -dimensional compact manifold representing a state of Existence. Let  $\partial M$  denote its boundary. Suppose the topological field theory action satisfies  $Z(\partial M) = N = \text{Nothing}$ . Prove that if the partition function  $Z(M) \neq 0$ , then  $\partial M$  cannot be homeomorphic to  $\emptyset$ .

### Problem 9 (Boolean Algebras: Venn Overlaps of Non-Not)

In a non-Boolean distributive lattice  $L$  (representing Parallax Logic), let  $a$  denote Infinity and  $b$  denote Eternity. Define the intuitionistic negation  $\neg a$ . Prove that if  $a \wedge b \neq \mathbf{0}$  and  $b \neq \neg a$ , then  $L$  cannot be a Heyting algebra where  $\neg a \vee a = \mathbf{1}$ .

### Problem 10 (Ergodic Theory: Invariance under Timeless Transformation)

Let  $(X, \mathcal{B}, \mu, T)$  be a measure-preserving dynamical system. Define  $\mathcal{S}_{\text{Timeless}}$  as the invariant  $\sigma$ -algebra  $\mathcal{I}_T = \{A \in \mathcal{B} : T^{-1}A = A\}$ . Prove that if  $T$  is non-ergodic,  $\mathcal{I}_T$  contains non-trivial elements that are neither  $\emptyset$  nor  $X$ , formalizing  $N = \text{Nothing} \neq \emptyset$ .

### Problem 11 (Axiomatic Set Theory: $N = \text{Nothing}$ and Non-Well-Founded Sets)

Under Zermelo-Fraenkel set theory with the Axiom of Foundation (WF), the empty set  $\emptyset$  is the unique set with von Neumann rank 0. If we admit the Anti-Foundation Axiom (AFA) of Aczel, define the set  $\Omega = \{\Omega\}$  ( $N = \text{Beyond}$ ). Prove that  $\Omega \neq \emptyset$ ,  $\Omega \neq 0$ , and  $\Omega \notin \text{WF}$ .

### Problem 12 (Differential Topology: De Rham Cohomology of $N = \text{Beyond}$ )

Let  $M$  be a smooth  $n$ -manifold representing  $N = \text{Beyond}$ . Suppose  $H_{dR}^k(M) = 0$  for all  $k \geq 1$ , but  $H_{dR}^0(M) \cong \mathbb{R}^{\mathbb{R}}$  (uncountably many connected components). Prove that  $M$  is non-compact and cannot be locally Euclidean if the topology on the components is non-discrete.

### Problem 13 (Lie Algebras: Timeless Symmetries)

Let  $\mathfrak{g}$  be a finite-dimensional semisimple Lie algebra acting on a state space  $V$ . Let  $T \in \mathfrak{g}$  be the time-generator element. Show that if the kernel of  $T$  under the adjoint representation  $\text{ad}_T : \mathfrak{g} \rightarrow \mathfrak{g}$  satisfies  $\ker(\text{ad}_T) = \mathfrak{g}$ , then  $\mathfrak{g}$  must be abelian, contradicting semisimplicity.

### Problem 14 (Complex Analysis: Singularities of The Parallax Function)

Let  $f(z)$  be a meromorphic function on  $\mathbb{C} \cup \{\infty\}$  defined as  $f(z) = \frac{\text{Infinity}(z)}{\text{Eternity}(z)}$ . Suppose  $z_0$  is an essential singularity of  $f(z)$ . Prove via the Big Picard Theorem that in any neighborhood of  $z_0$ ,  $f(z)$  takes every value in  $\mathbb{C}$  infinitely many times, with at most one exception, demonstrating the non-null cross-over of states.



### Problem 15 (Model Theory: Non-Standard Models of $N = \text{Nothing}$ )

Let  $T$  be the first-order theory of Peano Arithmetic (PA). Let  $\mathcal{M} \models T$  be a non-standard model of PA. Let  $c \in \mathcal{M}$  be a non-standard infinite integer. Define the interval  $I = [0, c] \subset \mathcal{M}$ . Prove that  $I$  cannot be mapped bijectively to  $\mathbb{N}$  within the object language of  $\mathcal{M}$ .

### Problem 16 (Symplectic Geometry: Timeless Submanifolds)

Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. Let  $N \subset M$  be a submanifold defined by the constraint of timelessness ( $S_{\text{Timeless}}$ ). If  $\omega|_N = 0$ ,  $N$  is isotropic. Show that the maximum dimension of a timeless submanifold  $N$  is  $n$  (Lagrangian), and prove that  $N$  cannot inherit a symplectic structure from  $M$ .

### Problem 17 (Algebraic Topology: Homotopy Groups of $N = \text{Beyond}$ )

Let  $X$  be a topological space modeling  $N = \text{Beyond}$ . Suppose  $\pi_0(X) \neq \{*\}$  and  $\pi_k(X) = 0$  for all  $k \geq 1$ . Prove that  $X$  is homotopically equivalent to a discrete space, but show that if  $X$  is connected and contractible, it cannot have non-trivial Venn cross-overs with a disconnected target space  $Y$ .

### Problem 18 (C\*-Algebras: Non-Commutative Eternity)

Let  $A$  be a C\*-algebra representing the observables of  $N = \text{Beyond}$ . Let  $\tau$  be a trace state on  $A$ . If  $p \in A$  is a projection ( $p^2 = p = p^*$ ) representing  $N = \text{Nothing}$ , prove that  $\tau(p) > 0$  implies  $p \neq 0$ , even if  $pAp \cong \mathbb{C}$  (a minimal projection).

### Problem 19 (Probability Theory: Martingale Persistence of Eternity)

Let  $(X_n)_{n \geq 0}$  be a continuous-time martingale on  $(\Omega, \mathcal{F}, \mathbb{P})$  representing states of Existence. Define the stopping time  $\tau = \inf\{t \geq 0 : X_t \in N = \text{Nothing}\}$ . If  $\mathbb{P}(\tau = \infty) > 0$ , prove that the state  $S_{\text{Eternity}}$  is persistent and non-evanescent under Optional Stopping Theorem conditions.

### Problem 20 (Sheaf Theory: Stalks of $N = \text{Nothing}$ )

Let  $\mathcal{F}$  be a sheaf of abelian groups on a topological space  $X$ . Suppose for a point  $p \in X$  (representing the singularity of  $N = \text{Beyond}$ ), the stalk  $\mathcal{F}_p \neq 0$ , but for every open neighborhood  $U \ni p$ , the section  $\mathcal{F}(U)$  contains no global non-zero element. Construct such a sheaf  $\mathcal{F}$  (e.g., a skyscraper sheaf or direct limit) and prove its stalk structure satisfies  $\mathcal{F}_p \neq 0$ .

## Solutions

### Solution 1

By definition in measure theory, the empty set  $\emptyset$  satisfies  $\mu(\emptyset) = 0$  for any measure  $\mu$ . Similarly, the zero element  $0 \in \mathbb{R}$  represented as a singleton set  $\{0\}$  in a non-atomic measure space satisfies  $\mu(\{0\}) = 0$ . Since  $\mu(\mathcal{N}) > 0$ , we have a direct inequality:

$$\mu(\mathcal{N}) > \mu(\emptyset) = \mu(\{0\}) = 0$$

In the category **Meas**, isomorphisms preserve measure zero sets. Because  $\mathcal{N}$  has strictly positive measure, no measure-preserving bijection can exist between  $\mathcal{N}$  and  $\emptyset$  or  $\{0\}$ . Hence,  $N = \text{Nothing}$  is strictly distinct from null and zero.

### Solution 2

The Lebesgue covering dimension  $\dim(X) = 0$  requires that every open cover has a disjoint open refinement. If  $\mathcal{S}_{\text{Eternity}}$  has no local neighborhood basis (e.g., equipped with the trivial topology or an indiscrete non-Hausdorff structure on a set with  $|\mathcal{S}| \geq 2$ ), open sets are restricted to  $\emptyset$  and  $\mathcal{S}_{\text{Eternity}}$ . Any subspace  $Y \subset \mathbb{R}^n$  with the induced Euclidean topology is  $T_2$  (Hausdorff) and regular. Since continuous embeddings preserve topological properties, a non-Hausdorff space without a neighborhood basis cannot embed into any Hausdorff space  $\mathbb{R}^n$ .



### Solution 3

In a category  $\mathbf{C}$ , a pullback  $P_1 \times_X P_2$  is defined by morphisms  $f: P_1 \rightarrow X$  and  $g: P_2 \rightarrow X$  such that the square commutes:

$$\begin{array}{ccc} P_1 \times_X P_2 & \xrightarrow{p_2} & P_2 \\ \downarrow p_1 & & \downarrow g \\ P_1 & \xrightarrow{f} & X \end{array}$$

Two objects are orthogonal if  $\text{Hom}(P_1, P_2) = \emptyset$  and their intersection in  $X$  yields the initial object 0. If  $P_1 \times_X P_2 \not\cong 0$ , then there exist non-trivial morphisms  $p_1: P_1 \times_X P_2 \rightarrow P_1$  and  $p_2: P_1 \times_X P_2 \rightarrow P_2$ . Thus,  $P_1$  and  $P_2$  share a common non-trivial subobject in  $\mathbf{C}$ , disproving orthogonality.

### Solution 4

Let  $M = \mathbb{R} \times M_{\text{space}}$ . The metric tensor  $g$  has components  $g_{\mu\nu}$ . The timeless condition implies  $g(V, \cdot) = 0$  for  $V = \partial/\partial t$ . In coordinate representation:

$$g_{00} = g\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = 0, \quad g_{0i} = g\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x^i}\right) = 0$$

The metric matrix collapses to:

$$g = \begin{pmatrix} 0 & 0 \\ 0 & g_{ij}^{\text{space}} \end{pmatrix}$$

$\det(g) = 0$ , meaning  $g$  is degenerate.  $M$  ceases to be Riemannian and reduces to a degenerate Galilean/Carrollian fiber bundle over  $M_{\text{space}}$ .  $M_{\text{space}}$  remains a valid  $(n - 1)$ -dimensional Riemannian manifold with non-degenerate metric  $g_{ij}^{\text{space}}$ , preserving spatial structure independently of time.

### Solution 5

In non-standard analysis,  ${}^*\mathbb{R}_\infty = \{x \in {}^*\mathbb{R} : |\{x\}| > n, \forall n \in \mathbb{N}\}$ . The halo of zero is  $\mu(0) = \{\epsilon \in {}^*\mathbb{R} : \forall n \in \mathbb{N}, |\{\epsilon\}| < 1/n\}$ .

1.  $\mu(0)$  is an additive subgroup of  ${}^*\mathbb{R}$  containing infinitesimals, whereas  ${}^*\mathbb{R}_\infty$  contains infinite numbers.
2.  $\mu(0)$  forms a non-archimedean subring without unit, whereas the quotient  $\mathbb{R}/\mathbb{Z}$  is a compact Lie group (the circle group  $S^1$ ).
3. Since  $|\{\mu(0)\}| = 2^{\aleph_0}$  (or higher cardinality depending on the ultrapower construction), and its algebraic structure lacks multiplicative units,  $S_{\text{Eternity}}$  modeled via  $\mu(0)$  cannot be isomorphic to  $S^1 \cong \mathbb{R}/\mathbb{Z}$  nor mapped bijectively to  ${}^*\mathbb{R}_\infty$  while preserving ordering/algebraic structure.

### Solution 6

Given  $U(t) = e^{-iHt/\hbar} \psi = \psi$  for all  $t \in \mathbb{R}$ :

Differentiating with respect to  $t$  at  $t = 0$ :

$$\left. \frac{d}{dt} U(t) \psi \right|_{t=0} = -\frac{i}{\hbar} H \psi = 0 \implies H \psi = 0$$

Thus,  $\psi \in \ker(H)$ .

Let  $\ker(H)$  be an infinite-dimensional closed subspace of  $\mathcal{H}$ .  $H$  acts as the zero operator on  $\ker(H)$ . For  $z \notin \sigma(H)$ , the resolvent  $R_z(H) = (H - zI)^{-1}$  restricted to  $\ker(H)$  is:

$$R_z(H)|_{\ker(H)} = (0 - zI)^{-1} = -\frac{1}{z} I_{\ker(H)}$$

The operator norm is  $\|R_z(H)|_{\ker(H)}\| = \frac{1}{|z|} < \infty$ .

Hence, infinite dimensionality of  $\ker(H)$  does not cause unboundedness of  $R_z(H)$  for any  $z \notin \sigma(H)$ .

### Solution 7

Let  $x \in R$  be non-zero and non-nilpotent, and let  $I = (x)$  be a proper two-sided ideal. If  $R/I \cong R$ , there exists a surjective ring homomorphism  $\phi: R \rightarrow R$  with  $\ker(\phi) = I \neq 0$ .

If  $R$  were Noetherian, every ascending chain of ideals would terminate. The chain  $\ker(\phi) \subset \ker(\phi^2) \subset \ker(\phi^3) \subset \dots$  must stabilize. However, since  $R/I \cong R$ , we can iterate  $\phi$  infinitely to form an infinite strictly increasing chain  $I_1 \subset I_2 \subset I_3 \dots$ , contradicting the Noetherian property. Furthermore, if  $x$  were a unit,  $I = (x) = R \implies R/I = \{0\} \not\cong R$ . Thus,  $x$  is neither a unit nor a classical zero divisor.

### Solution 8

In topological field theory, the partition function  $Z(M)$  assigns an element in a vector space (or complex number) to a manifold  $M$ . If  $M$  is a manifold with boundary  $\partial M$ ,  $Z(M)$  defines a state in the Hilbert space  $\mathcal{H}(\partial M)$  assigned to  $\partial M$ .

If  $\partial M = \emptyset$ ,  $\mathcal{H}(\emptyset) \cong \mathbb{C}$ , and  $Z(M) \in \mathbb{C}$  is a topological invariant.

If  $Z(\partial M) = N = \text{Nothing}$  where  $N = \text{Nothing}$  represents a non-trivial vector space with positive dimension  $\dim(\mathcal{H}(\partial M)) > 0$ , then  $\mathcal{H}(\partial M)$  cannot be the vector space associated with the empty boundary if  $Z(M) \neq 0$ . Thus,  $\partial M \neq \emptyset$ .

### Solution 9

In a Heyting algebra  $L$ , the law of excluded middle  $a \vee \neg a = \mathbf{1}$  does not hold universally. The pseudocomplement  $\neg a$  is defined as  $\max\{x \in L : a \wedge x = \mathbf{0}\}$ .

Given  $a \wedge b \neq \mathbf{0}$ ,  $b$  cannot be a lower bound for the complement condition of  $a$ , meaning  $b \not\leq \neg a$ .

If  $L$  were a Boolean algebra (where  $a \vee \neg a = \mathbf{1}$ ), every element  $b$  would satisfy  $b = (b \wedge a) \vee (b \wedge \neg a)$ .

Since  $b \not\leq \neg a$  and  $a \wedge b \neq \mathbf{0}$ ,  $b$  overlaps with  $a$  non-trivially. If  $a \vee \neg a = \mathbf{1}$  held for all  $a \in L$ ,  $L$  would collapse to a Boolean lattice, contradicting the intuitionistic non-classical logic premise of  $N = \text{Nothing} \neq \neg \text{Existence}$ .

### Solution 10

By definition,  $\mathcal{J}_T = \{A \in \mathcal{B} : T^{-1}A = A\}$  is a  $\sigma$ -algebra.

A system is ergodic if and only if every  $A \in \mathcal{J}_T$  satisfies  $\mu(A) = 0$  or  $\mu(A) = 1$ .

If  $T$  is non-ergodic, by definition there exists at least one set  $E \in \mathcal{J}_T$  such that  $0 < \mu(E) < 1$ .

Since  $\mu(E) \neq 0$ ,  $E \neq \emptyset \pmod{\mu}$ .

Since  $\mu(E) \neq 1$ ,  $E \neq X \pmod{\mu}$ .

Thus,  $E \in \mathcal{J}_T$  is a non-trivial invariant state, satisfying  $N = \text{Nothing} \neq \emptyset$  and  $N = \text{Nothing} \neq X$ .

### Solution 11

Under the Axiom of Foundation (WF), for every non-empty set  $x$ , there exists  $y \in x$  such that  $x \cap y = \emptyset$ .

Consider  $\Omega = \{\Omega\}$ .  $x = \{\Omega\}$ . The only element in  $x$  is  $\Omega$ .

$x \cap \Omega = \{\Omega\} \cap \Omega = \{\Omega\} \neq \emptyset$ . This violates Foundation, so  $\Omega \notin \mathbf{WF}$ .

Under Aczel's Anti-Foundation Axiom (AFA), every accessible pointed graph (APG) has a unique decoration. A graph with one node and one directed self-loop uniquely defines  $\Omega = \{\Omega\}$ .

$\Omega$  contains an element (itself), so  $\Omega \neq \emptyset$ . In von Neumann ordinals,  $0 = \emptyset$ , so  $\Omega \neq 0$ .

### Solution 12

The de Rham cohomology group  $H_{dR}^0(M)$  is isomorphic to  $\mathbb{R}^k$ , where  $k$  is the number of connected components of  $M$ .

If  $H_{dR}^0(M) \cong \mathbb{R}^{\mathbb{R}}$ ,  $M$  has uncountably many connected components ( $|\{C\}| = 2^{\aleph_0}$ ).

If  $M$  were compact, any open cover by disjoint component neighborhoods would yield a finite subcover, forcing  $k$  to be finite. Thus  $M$  is non-compact.

If the components have non-discrete topology,  $M$  cannot be locally Euclidean of fixed dimension  $n$  because the disjoint union of uncountably many non-discrete manifolds violates second-countability (a standard requirement for topological manifolds embedding in Euclidean space).

### Solution 13

The adjoint representation  $\text{ad} : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$  is defined by  $\text{ad}_T(X) = [T, X]$ .

If  $\ker(\text{ad}_T) = \mathfrak{g}$ , then for all  $X \in \mathfrak{g}$ :

$$\text{ad}_T(X) = [T, X] = 0$$

This means  $T$  belongs to the center  $Z(\mathfrak{g})$  of  $\mathfrak{g}$ .

For a Lie algebra to be semisimple, its center must be trivial:  $Z(\mathfrak{g}) = \{0\}$ .

If  $T \neq 0$  and  $T \in Z(\mathfrak{g})$ ,  $Z(\mathfrak{g}) \neq \{0\}$ , which contradicts the assumption that  $\mathfrak{g}$  is semisimple.

### Solution 14

Let  $f(z) = \frac{\text{Infinity}(z)}{\text{Eternity}(z)}$  have an essential singularity at  $z_0 \in \mathbb{C}$ .

By the Big Picard Theorem: If an analytic function  $f$  has an essential singularity at  $z_0$ , then in any punctured neighborhood  $U \setminus \{z_0\}$ ,  $f(z)$  takes on all possible complex values infinitely many times, with at most one exception  $c \in \mathbb{C}$ .

Therefore, for any value  $w \in \mathbb{C}$  (except possibly one exception), the equation:

$$\frac{\text{Infinity}(z)}{\text{Eternity}(z)} = w$$



has infinitely many solutions in  $U \setminus \{z_0\}$ . This proves the intersection/cross-over region in the complex plane contains an infinite spectrum of values, showing non-null overlap.

### Solution 15

Let  $\mathcal{M} \models T$  be a non-standard model of Peano Arithmetic, and let  $c \in \mathcal{M} \setminus \mathbb{N}$  be an infinite non-standard element.

Assume there exists an internal bijection  $f: [0, c]_{\mathcal{M}} \rightarrow \mathbb{N}$ .

In PA, every non-empty definable subset that is bounded above has a maximal element (or by mathematical induction, finite intervals cannot be in bijection with proper sub-initial segments).

$\mathbb{N}$  is an order-ideal in  $\mathcal{M}$  (an initial segment), but  $\mathbb{N}$  is NOT a definable subset of  $\mathcal{M}$  (by Overspill Lemma: if a property holds for all standard  $n \in \mathbb{N}$ , it holds for some non-standard  $x > \mathbb{N}$ ).

Thus, no internal bijection  $f$  can exist in  $\mathcal{M}$  mapping  $[0, c]$  to  $\mathbb{N}$ .

### Solution 16

Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold.  $N \subset M$  is a submanifold with  $\omega|_N = 0$  (isotropic).

For any  $p \in N$ ,  $T_p N \subset (T_p N)^\omega$ , where  $(T_p N)^\omega = \{v \in T_p M: \omega(v, w) = 0, \forall w \in T_p N\}$ .

By linear algebra of symplectic vector spaces:

$$\dim(T_p N) + \dim((T_p N)^\omega) = \dim(T_p M) = 2n$$

Since  $T_p N \subset (T_p N)^\omega$ , we have  $\dim(T_p N) \leq \dim((T_p N)^\omega)$ .

Thus:

$$2\dim(T_p N) \leq 2n \implies \dim(N) \leq n$$

Since  $\omega|_N = 0$ , the induced 2-form on  $N$  is identically zero. A symplectic form must be non-degenerate. Because  $\omega|_N = 0$ ,  $N$  cannot inherit a symplectic structure from  $M$ .

### Solution 17

1. If  $\pi_0(X) \neq \{*\}$  and  $\pi_k(X) = 0$  for all  $k \geq 1$ ,  $X$  has multiple path-connected components, each of which is weakly contractible. By Whitehead's Theorem (for CW complexes), each component is homotopically equivalent to a point  $*$ . Thus  $X \simeq \pi_0(X)$  (a discrete space).
2. If  $X$  is connected and contractible,  $X \simeq \{*\}$ .

Suppose  $Y$  is disconnected, so  $\pi_0(Y)$  has at least 2 points.

Any continuous map  $f: X \rightarrow Y$  must map the path-connected space  $X$  into a single path-component of  $Y$ .

Thus,  $f(X) \subset Y_i \subsetneq Y$ .  $X$  cannot cover or overlap with  $Y$  across multiple components simultaneously, proving that a contractible space cannot have a non-trivial topological cross-over spanning disconnected target states.

### Solution 18

Let  $A$  be a  $C^*$ -algebra and  $\tau: A \rightarrow \mathbb{C}$  a faithful trace state (meaning  $\tau(a^*a) \geq 0$  and  $\tau(a^*a) = 0 \iff a = 0$ ).

Let  $p \in A$  be a projection operator ( $p^2 = p = p^*$ ).

Since  $p = p^* p$ , we have:



$$\tau(p) = \tau(p^* p)$$

By the faithfulness of the trace state  $\tau$ :

$$\tau(p^* p) = 0 \iff p = 0$$

Given that  $\tau(p) > 0$ , it follows immediately that  $p \neq 0$ .

Even if  $pAp \cong \mathbb{C}$  (indicating that  $p$  is a minimal projection representing an indivisible state  $N = \text{Nothing}$ ),  $p$  remains a non-zero element in  $A$ .

### Solution 19

Let  $(X_t)_{t \geq 0}$  be a martingale on  $(\Omega, \mathcal{F}, \mathbb{P})$ . Let  $\tau = \inf\{t \geq 0: X_t \in N = \text{Nothing}\}$ .

We are given  $\mathbb{P}(\tau = \infty) > 0$ .

Define  $A = \{\omega \in \Omega: \tau(\omega) = \infty\}$ .

By definition,  $\mathbb{P}(A) > 0$ .

If the state  $\mathcal{S}_{\text{Eternity}}$  were transient or evanescent, the indicator variable  $I_{\{\tau < \infty\}} = 1$  almost surely, meaning  $\mathbb{P}(\tau < \infty) = 1 \implies \mathbb{P}(\tau = \infty) = 0$ .

Since  $\mathbb{P}(\tau = \infty) > 0$ , the set of paths that remain in the timeless state indefinitely without being stopped has positive probability measure. Thus,  $\mathcal{S}_{\text{Eternity}}$  is persistent and non-evanescent.

## Solution 20

Let  $X = \mathbb{R}$  with the standard topology. Let  $p = 0$ .

Define the skyscraper sheaf  $i_{0*}(\mathbb{A})$  supported at 0 for an abelian group  $A = \mathbb{Z}$ :

$$(i_{0*}(\mathbb{Z}))(U) = \begin{cases} \mathbb{Z} & \text{if } 0 \in U \\ 0 & \text{if } 0 \notin U \end{cases}$$

The stalk at  $x$  is:

$$\mathcal{F}_x = \varinjlim_{U \ni x} \mathcal{F}(U)$$

For  $x = 0$ , every open neighborhood  $U \ni 0$  satisfies  $\mathcal{F}(U) = \mathbb{Z}$ .

The direct limit of constant groups  $\mathbb{Z}$  over the directed set of open neighborhoods of 0 is:

$$\mathcal{F}_0 \cong \mathbb{Z} \neq 0$$

Now consider a subsheaf  $\mathcal{F}$  defined such that section restriction maps cancel global sections while retaining local persistence at 0:

Take  $\mathcal{F} = \bigoplus_{n=1}^{\infty} \mathcal{G}_n$  where  $\mathcal{G}_n$  are skyscraper sheaves at  $1/n$  converging to 0.

The stalk at 0 gains non-zero elements in the inductive limit  $\mathcal{F}_0 \neq 0$ , even if  $\Gamma(X, \mathcal{F}) = 0$ .

For  $x = 0$ , every open neighborhood  $U \ni 0$  satisfies  $\mathcal{F}(U) = \mathbb{Z}$ .

The direct limit of constant groups  $\mathbb{Z}$  over the directed set of open neighborhoods of  $0$  is:

$$\mathcal{F}_0 \cong \mathbb{Z} \neq 0$$

Now consider a subsheaf  $\mathcal{F}$  defined such that section restriction maps cancel global sections while retaining local persistence at  $0$ :

Take  $\mathcal{F} = \bigoplus_{n=1}^{\infty} \mathcal{G}_n$  where  $\mathcal{G}_n$  are skyscraper sheaves at  $1/n$  converging to  $0$ .

The stalk at  $0$  gains non-zero elements in the inductive limit  $\mathcal{F}_0 \neq 0$ , even if  $\Gamma(X, \mathcal{F}) = 0$ .

This confirms that the local state  $N = \text{Nothing}$  at point  $p$  satisfies  $\mathcal{F}_n \neq 0$ .