

collected prior to vaccination, and weekly through week 16, and on week 20. Indirect blood pressure was measured prior to vaccination, and every 2 weeks through week 16 and weeks 20 and 24.

Mean antibody titer was determined by ELISA and peaked at  $21,400 \pm 19,380$  on week 6 and was  $9,986 \pm 10,792$  at week 20. Data were evaluated statistically by repeated measures ANOVA with Dunnett's multiple comparison post hoc analysis and Pearson correlation. Systolic arterial pressure increased at weeks 8 and 20 compared to baseline ( $P = 0.008$ ). Neither diastolic nor mean arterial pressure were significantly changed over time, however there was a significant negative correlation of mean antibody titer with both mean ( $R = -0.24$ ;  $P = 0.03$ ) and diastolic ( $R = -0.22$ ;  $P = 0.05$ ) arterial pressure. While a robust immunologic response to AngQb vaccination was demonstrated, additional studies are warranted to evaluate arterial blood pressure responses in hypertensive horses.

#### ABSTRACT E-83

##### HYPERTENSIVE CARDIOMYOPATHY IN HORSES.

C Navas de Solis<sup>1</sup>, J Slack<sup>1</sup>, V.B Reef<sup>1</sup>. <sup>1</sup>New Bolton Center, University of Pennsylvania, Kennett Square, Pennsylvania

The objective was to describe the clinical presentation, physical examination, prognosis, echocardiographic and pathologic findings of horses with hypertensive cardiomyopathy. Records from cases presented between 1995 and 2011 diagnosed with cardiac hypertrophy and systemic hypertension (systolic and mean non invasive blood pressure > 144 and 116 mmHg respectively) were studied retrospectively.

Five horses met the inclusion criteria (0.26% of horses evaluated for cardiac disease). There were 3 geldings and 2 mares of multiple breeds with a mean±SD age of 18±4 years (range 13-24). The main diagnoses were chronic laminitis in 3 cases and chronic renal failure in two. Persistent tachycardia, hypertension, chronic laminitis or a combination of these prompted the cardiac evaluations. Non invasive blood pressure [systolic/diastolic (mean)] was 212±36 mmHg [range 183-261 mmHg] / 169±38 mmHg [range 100-190 mmHg] (171±41 mmHg [range 126-222 mmHg]). No arrhythmias were reported. All horses had increased relative wall thickness 0.94±0.29 [range 0.57-1.26]. The interventricular septum was thickened in all horses at end diastole and in 4 horses at peak systole. The left ventricular free wall was thickened and the internal diameter reduced at end diastole and peak systole in three horses. All horses were euthanized due to grave prognosis of the primary disease. All three horses that underwent necropsy had macroscopic cardiac hypertrophy and 2 had microscopic cardiovascular changes.

Hypertensive cardiomyopathy is rare and was associated with chronic pain or renal disease. Information about the reversibility, the importance of early detection of cardiomyopathy in hypertensive horses and its long-term sequelae is needed.

#### ABSTRACT E-84

##### OXIDATIVE METABOLISM AND *IN VITRO* IMMUNE RESPONSE OF HORSES SUPPLEMENTED WITH VITAMIN COMPLEX ADE.

RW Weigel, M Mirian Souza, FN Fernandes, WR Della Libera, AMMP Sucupira M.C.A.. University of São Paulo, Brazil

Horses working at explosive exercise are subjected to oxidative stress. Animals unable to balance the production of reactive oxygen species (ROS) and antioxidant reaction can lose performance and have longer recover time. The aim of this study was to evaluate oxidative metabolism and *in vitro* immune response of polo players equines treated or not with parenteral ADE vitamin complex.

Twenty horses, crossbred, 7.0 (± 1.7) yrs of age, trained for equestrian polo matches were used. They were randomly distributed in two groups. Control group received 10 mL of physiologic saline solution, IM, and the Treated group received 1ml/50kgBW IM of the supplement containing vitamins A (27 000 000 UI /

100mL), D<sub>3</sub> (8 000 000 UI/100mL) e E (8 000 mg/100mL). The horses were at rest 60 days before the game. The research began with an application 30 and 15 days before the start of training. At the first training blood samples were collected before the game (M0), 15 minutes (M1), 90 minutes (M2) and 3 hours (M3) after the only chukker. Plasma was assayed for malondialdehyde (MDA). Superoxide dismutase (SOD) concentration was determined in red blood cells. Reduced glutathione (GSH) and burst-phagocytosis were determined in total blood. The burst-phagocytosis was performed by flow cytometry. Treatments were compared by ANOVA (Tukey test) and Pearson coefficient was determined between variables using Statistical Software MINITAB®.

There were no differences in the concentrations of MDA, SOD, GSH and burst-phagocytosis between Control and Treated groups ( $P > 0.05$ ). Only the SOD concentration and phagocytic intensity were negatively correlated ( $r = -0.302$  and  $P = 0.1$ ). In this study parenteral ADE supplementation didn't make a difference in the oxidative metabolism and *in vitro* immune response in horses trained for polo matches.

#### ABSTRACT E-85

##### ASSESSMENT OF VITAMIN E LEVELS IN A POPULATION OF HEALTHY ADULT HORSES.

K Vander Werf, EG Davis, C Blevins. Kansas State University, Manhattan, KS

Equine Motor Neuron Disease (EMND) is associated with chronic vitamin E deficiency resulting in neuromuscular weakness, hyperhidrosis, preferred recumbency, and muscle fasciculation. Among 7 clinical cases presented within the past 3 years, 57% had adequate serum vitamin E levels (ref. range: 2.0-4.0 µg/mL).

Serum tocopherol levels in 50 healthy, adult horses maintained on dry lots and 50 horses maintained on pasture for at least 1 year were analyzed at a commercial laboratory. Variables examined included vital parameters, environment (dry lot vs pasture), duration in environment, and concentrate supplementation. Horses were excluded if in the current environment for less than 1 year.

Vitamin E levels of horses maintained on pasture ( $5.44 \pm 2.26$  µg/mL) and those maintained on dry lot (mean:  $3.99 \pm 2.84$  µg/mL) were significantly different ( $p = 0.0003$ ). Additionally, there was a significant positive correlation between age and tocopherol level ( $r = 0.30$ ,  $p = 0.0026$ ). Geldings maintained on dry lots had significantly lower tocopherol levels than mares maintained in the same environment or geldings and mares on pasture ( $p = 0.03$ ). There was no significant correlation between breed, brand of pelleted feed, or duration in current environment and serum vitamin E levels.

Horses maintained on dry lots should be monitored for signs of EMND and should have vitamin E levels measured on a regular basis. Geldings appear to be at risk of low vitamin E levels when maintained in dry lot environments. Supplementation with natural vitamin E is recommended for horses with levels less than 2.0 µg/mL.

## SMALL ANIMAL – ENDOCRINOLOGY

#### ABSTRACT EN-1

##### RESTORATION OF EUTHYROIDISM IN MEDICALLY TREATED HYPERTHYROID CATS WITH IATROGENIC HYPOTHYROIDISM (IH) IMPROVES RENAL FUNCTION.

TL Williams, J Elliott, HM Syme. Royal Veterinary College, London, UK

IH is reported to increase the incidence of azotemia in hyperthyroid cats following treatment. IH is also associated with a decreased heart rate (HR), packed cell volume (PCV) and plasma alkaline phosphatase activity (ALP). Hypothyroidism reduces glomerular filtration rate (GFR) in other species, and treatment of hypothyroidism in dogs is reported to increase GFR. There-