Chapter 5 Rational Exponents and Radical Functions

Section 5-1 nth Roots and Rational Exponents

nth Roots

You can extend the concept of a square root to other types of roots. For example, 2 is a cube root of 8 because $2^3 = 8$. In general, for an integer n greater than 1, if $b^n = a$, then b is an **nth root of a**. An nth root of a is written as $\sqrt[n]{a}$, where n is the **index** of the radical.

You can also write an nth root of a as a power of a. If you assume the Power of a Power Property applies to rational exponents, then the following is true.

$$(a^{1/2})^2 = a^{(1/2) \cdot 2} = a^1 = a$$

$$(a^{1/3})^3 = a^{(1/3) \cdot 3} = a^1 = a$$

$$(a^{1/4})^4 = a^{(1/4) \cdot 4} = a^1 = a$$

Because $a^{1/2}$ is a number whose square is a, you can write $\sqrt{a} = a^{1/2}$. Similarly, $\sqrt[3]{a} = a^{1/3}$ and $\sqrt[4]{a} = a^{1/4}$. In general, $\sqrt[n]{a} = a^{1/n}$ for any integer n greater than 1.

1

Rational Exponents

A rational exponent does not have to be of the form $\frac{1}{n}$. Other rational numbers, such as $\frac{3}{2}$ and $-\frac{1}{2}$, can also be used as exponents. Two properties of rational exponents are shown below.

🔄 Core Concept

Rational Exponents
Let
$$a^{1/n}$$
 be an n th root of a , and let m be a positive integer.
$$a^{m/n} = (a^{1/n})^m = (\sqrt[n]{a})^m$$

$$a^{-m/n} = \frac{1}{a^{m/n}} = \frac{1}{(a^{1/n})^m} = \frac{1}{(\sqrt[n]{a})^m}, a \neq 0$$

Evaluate each expression.

b.
$$32^{-3/5}$$

SOLUTION

Radical Form

Approximating Expressions with Rational Exponents

Evaluate each expression using a calculator. Round your answer to two decimal places.

b.
$$12^{3/8}$$

e.
$$(\sqrt[4]{7})^3$$

SOLUTION

a.
$$9^{1/5} \approx 1.55$$

b.
$$12^{3/8} \approx 2.54$$

c. Before evaluating
$$(\sqrt[4]{7})^3$$
, rewrite the expression in rational exponent form.

$$(\sqrt[4]{7})^3 = 7^{3/4} \approx 4.30$$

Evaluate the expression without using a calculator.

- **5.** 45/2
- 6. 9^{-1/2}
- **7.** 81^{3/4}
- 8. 17/8

Evaluate the expression using a calculator. Round your answer to two decimal places when appropriate.

- **9.** 62/5
- **10.** $64^{-2/3}$ **11.** $(\sqrt[4]{16})^5$ **12.** $(\sqrt[3]{-30})^2$

Solving Equations Using nth Roots

To solve an equation of the form $u^n = d$, where u is an algebraic expression, take the nth root of each side.

EXAMPLE 4 Solving Equations Using *n*th Roots

Find the real solution(s) of (a) $4x^5 = 128$ and (b) $(x - 3)^4 = 21$.

COMMON ERROR

When n is even and a > 0, be sure to consider both the positive and negative nth roots of a.



EXAMPLE 5

Real-Life Application

A hospital purchases an ultrasound machine for \$50,000. The hospital expects the useful life of the machine to be 10 years, at which time its value will have depreciated to \$8000. The hospital uses the declining balances method for depreciation, so the annual depreciation rate r (in decimal form) is given by the formula

$$r = 1 - \left(\frac{S}{C}\right)^{1/n}.$$

In the formula, n is the useful life of the item (in years), S is the salvage value (in dollars), and C is the original cost (in dollars). What annual depreciation rate did the hospital use?

SOLUTION