

## Brief Annotated Bibliography for My Pain Alert® Scale Communication Tool

Bahan, Ben and Joe Dannis. *Signs for Me Basic Sign Vocabulary for Children, Parents & Teachers*. San Diego, CA: Dawn Sign Press, 1990. Print.

City of Hope Pain & Palliative Care Resource Center. *Categories of Materials. IV. Pain and Symptom Management. A. Pain Assessment Tools*. Modified February 1, 2016. [www.prc.coh/pain\\_assessment.asp](http://www.prc.coh/pain_assessment.asp) Web. Accessed March 30, 2016.

Drager, Kathryn D. R. "Aided Modeling Interventions for Children with Autism Spectrum Disorders Who Require AAC." *SIG 12 Perspectives on Augmentative and Alternative Communication*. Volume 18 (December 2009): 114-120. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "This article discusses the research evidence suggesting that aided modeling interventions may be effective for children with autism spectrum disorders (ASD)."

Dunn, Lloyd and Dunn, Leota. *Peabody Picture Vocabulary Test- Revised Manual for forms L and M*. Circle Pines, Minnesota: American Guidance Service, 1981. Print.

Findley, G. Allen. "Pain in children." *Pain Research and Management*: 2006 Autumn; 11(3); 156. [www.ncbi.nlm.nih.gov/pmc/articles/PMC2539003/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2539003/) Web. Accessed May 10, 2017. "In my opinion, failure to recognize children's pain remains the greatest barrier to adequate care, so this work is especially relevant." (referring to von Baeyer article).

Flodin, Mickey. *Signing for Kids*. New York, NY: Perigee Books, 1991. Print.

Hasselkus, Amy. "Health Communication: Implications for Diverse Populations." *SIG 14 Perspectives on Communication Disorders and Sciences in Culturally and Linguistically Diverse Populations*. Volume 18 (March 2011): 12-19. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "...poor health literacy has been estimated to cost between \$106-238 billion each year. Health literacy also contributes to health disparities."

Komesidou, Rouzana and Holly L. Storkel. "Learning and Remembering New Words: Clinical Illustrations from Children with Specific Language Impairment." *SIG 1 Perspectives on Language Learning and Education*. Volume 22 (November 2015): 138-146. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "...at least two neurocognitive processes involved in word learning: a) learning from input and b) memory evolution in the absence of input."

McGrath, P., Findley, G. Ritchie, J. and S. Dowden *Pain, pain go away: Helping children with Pain 2<sup>nd</sup> Ed*. 2003 [Pediatric-pain.ca/wp-content/uploads/2-13/04/PPGA2003.pdf](http://pediatric-pain.ca/wp-content/uploads/2-13/04/PPGA2003.pdf). Web. Accessed January 5, 2017.

Mesko, Phyllis J. and Jennifer L. Clark. "Commentary - The Importance of pictures to communicate pain location in children." *Pediatric Pain Letter*: February 2019. Vol. 21 No. 1: 11-14. <http://www.childpain.org/ppl/>. Web. Accessed February 23, 2019.

"Measuring your child's pain using the pain scale." *Pain Management for Children*. Health-First.Org. [www.health-first.org/health\\_info/your\\_health\\_first/kids/pain.cfm](http://www.health-first.org/health_info/your_health_first/kids/pain.cfm). Web. Accessed 5/21/2017.

Moseley, Lorimer, editor. "Can Preschool-age children reliably report the intensity of their pain?" *Body in Mind*. (22 January 2018). <http://www.bodyinmind.org/children-reporting-pain>. Web. Accessed July 1, 2019. "The only studies that do provide separate, age-specific data for preschoolers show that most 3-year-olds are not able to provide a graded estimate of the intensity of their pain using published scales."

Rao, Paul R. "Our Role in Effective Parent-Provider Communication." *The Asha Leader*. Volume 16 (November 2011): 17. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "Based on data from NIH and MarkeTrak VII estimates of "communication-vulnerable individuals...in health care settings in the United States...is more than two-thirds of the entire population." "Anyone entering the health care system should have services and tools available to allow them to communicate with health care professionals."

Roberts, Megan Y. and Ann P. Kaiser. "The Effectiveness of Parent-Implemented Language Interventions: A Meta-Analysis." *American Journal of Speech-Language Pathology*. Volume 20 (August 2011): 180-199. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "The results of this review indicate that parent-implemented language interventions are an effective approach to early language intervention for young children with language impairments."

Rossetti, Louis. *The Rosetti Infant-Toddler Language Scale*. East Moline, Illinois: LinguiSystems, Inc. 1990. Print.

Spencer, Trina D., Douglas B. Petersen and John L. Adams. "Tier 2 Language Intervention for Diverse Preschoolers: An Early Stage Randomized Control Group Study Following an Analysis of Response to Intervention." *American Journal of Speech Language Pathology*. Volume 24 (November 2015): 619-635. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. "...evidence that narrative language intervention is an effective approach to improving language skills of preschoolers with diverse language needs."

Stransky, Michelle L. and Megan A. Morris. "Communication: It's Critical to Care." *The ASHA Leader*, Vol. 24 Issue 3 (March 2019) 46-55. Print.

Steinberg, Martin L. *American Sign Language Concise Dictionary*. New York, NY: Harper and Row Publishers, 1990. Print.

“Tell me how it hurts.” *Consumer Reports Magazine* (June 2016): 30-31. Print.

“The Child’s Understanding of Pain, Behavior Responses, and Verbal Descriptions by Developmental Stage” *Management in Children’s Pain Assessment*. Chap. 42, pp1104-1105, 1109. M42\_LOND8464\_03\_SE\_CH42.qxd 11/24/09 Web. Accessed July 31, 2017.

Verghise, Susan T. and Raafat S. Hannallah. “Acute Pain Management in Children.” *Journal of Pain Management* (July 2010): 105-123. [www.ncbi.nlm.nih.gov >PMC3004641](http://www.ncbi.nlm.nih.gov/PMC3004641) Web. Accessed May 13, 2016. “...factors such as fear, anxiety, coping style, and lack of social support can further aggravate physical pain in children.”

Voelme, Krista and Holly L. Storkel. “Teaching New Words to Children with Specific Language Impairment.” *SIG 1 Perspectives on Language Learning and Education*. Volume 22 (November 2015): 131-137. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. “...review of evidence base for interactive book reading to facilitate new word learning for preschool and school age children,” “children with Specific Language Impairment need 2-3 times as many exposures as typically developing children to learn new words.” “Current results suggest that 36 story retelling exposures is the most promising intensity for children with SLI.”

von Baeyer, Carl. “Children’s self-reports of pain intensity: Scale selection, limitations and interpretation.” *Pain Research and Management*: 2006 Autumn 11 (3) 157-162. [www.ncbi.nlm.gov/pmc/articles/PMC2539005/](http://www.ncbi.nlm.gov/pmc/articles/PMC2539005/) Web. Accessed May 10 2017.

Walsh, Trudi M., Patrick J. McGrath and Douglas K. Symons. “Attachment dimensions and young children’s response to pain.” *Pain Research and Management* Volume 13 #1 (January/February 2008): 33-40. [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) Web . Accessed March 13, 2016.

Wilner, L. Scott and Robert Arnold. *Fast Facts and Concepts: #126 Pain Assessment in the Cognitively Impaired*. Palliative Care Network of Wisconsin. [www.mypcnw.org](http://www.mypcnw.org) Web. Accessed March 27, 2016.

Wong, Donna L. and Connie M. Baker. “Pain in Children: Comparison of Assessment Scales.” *Pediatric Nursing* Volume 14 #1 (January-February 1988) [www.wongbakerfaces.org](http://www.wongbakerfaces.org) Web. Accessed March 29, 2016.

Yorkston, Kathryn M., Carolyn R. Baylor and Michael I. Burns. “Stimulating Patient Communication Strategies.” *The ASHA Leader* Volume 21 (March 2016): 46-51. [www.asha.org](http://www.asha.org) Web. Accessed March 27, 2016. “Medical students practice ‘FRAME’ strategies” paraphrased: F = familiarize with communication strategies patient prefers or require, R= reduce speaking rate, A= assist with communication, M= mix communication methods, and E= engage the patient.