

## Recommended Data Standardized Collection

MPAS+ needs to have consistency across data sets from different studies. This consistency will enable decisive meta-studies to be published clarifying this tool's validity and reliability, as well as, the cost impact of improving patient pain communication. A second goal is to facilitate single subject studies as there are many people with unique combinations of disorders needing scientific attention, and advanced students in need of single subject studies to perfect their clinical and research skills. We encourage therapeutic professionals in training to use the following in their data sets.

Question for ADMITTING, start of new year, or change of personnel dealing primarily with this limited communicator are, mark as many as apply:

How does patient express pain at home? \_\_\_\_\_ tells caregiver words like "hurt", "pain", "ouch"  
\_\_\_\_\_ cries. \_\_\_\_\_ points \_\_\_\_\_ squirms, body actions. \_\_\_\_\_ face shows distress  
\_\_\_\_\_ uses a pain scale \_\_\_\_\_ uses assisted communication device  
\_\_\_\_\_ meets own needs without other's help. \_\_\_\_\_ this has been a problem

Use line below to tell which pain scale used, or describe communication effort or device:

---

Questions for RELEASE from facility, end of year, or end of personnel's responsibility (signed):

1. Patient's communication of pain is: \_\_\_\_\_ clear and consistent, \_\_\_\_\_ adequate, \_\_\_\_\_ inconsistent or questionable: \_\_\_\_\_
2. If questionable or inconsistent, was care effected: \_\_\_\_\_ patient required more staff effort to get needs met; \_\_\_\_\_ patient's pain reaction resulted in injury (\_\_\_ patient or\_\_ staff);  
\_\_\_\_\_ patient's reaction to pain resulted in material waste; \_\_\_\_\_ patient care required more staff time.
3. If patient response required more staff effort what helped? \_\_\_\_\_  
\_\_\_\_\_ What hindered? \_\_\_\_\_
4. Practice or use of a pain scale prior to use with pain: \_\_\_\_\_ none, \_\_\_\_\_ 1-3 sessions, \_\_\_\_\_ 4-6 sessions, \_\_\_\_\_ 7-10 sessions, \_\_\_\_\_ 11-35 sessions, \_\_\_\_\_ 36 or more sessions, \_\_\_\_\_ unknown.
5. Family/caregivers were present during \_\_\_\_\_ percent of signee's interaction with patient.  
Family efforts supported patient pain scale training: \_\_\_\_\_actively \_\_\_\_\_sometimes \_\_\_\_\_not observed

Comments: \_\_\_\_\_

---

Signature/Title \_\_\_\_\_ Date \_\_\_\_\_