PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

NAME:	DATE:				
Over the last 2 weeks, how often have you been bothered by any of the following problems?					
(use "✓" to indicate your answer)	Not at all	Several days	More than half the days	Nearly every day	
1. Little interest or pleasure in doing things	0	1	2, 3	3	
2. Feeling down, depressed, or hopeless	0	1	2	3	
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3	
4. Feeling tired or having little energy	0	·1	2	3	
5. Poor appetite or overeating	0	1	2	3	
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	2	3	
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2, 2	3	
8. Moving or speaking so slowly that other people could have noticed. Or the opposite — being so figety or restless that you have been moving around a lot more than usual	0	1	2	3	
9. Thoughts that you would be better off dead, or of hurting yourself	0	2 1 - 1 - 2	2	3	
	add columns		+	+	
(Healthcare professional: For interpretation of TOT. please refer to accompanying scoring card).	AL, TOTAL:				
10. If you checked off any problems, how difficult	Not difficult at all				
have these problems made it for you to do	Somewhat difficult				
your work, take care of things at home, or get	Very difficult				
along with other people?	Extremely difficult				

Copyright © 1999 Pfizer Inc. All rights reserved. Reproduced with permission. PRIME-MD© is a trademark of Pfizer Inc. A2663B 10-04-2005

Generalized Anxiety Disorder 7-item (GAD-7) scale

Over the last 2 weeks, how often have you been bothered by the following problems?	Not at all sure	Several days	Over half the days	Nearly every day
1. Feeling nervous, anxious, or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it's hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	l	2	3
Add the score for each column	+	+	+	
Total Score (add your column scores) =				

If you checked off any problems, how difficult have these made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all
Somewhat difficult
Very difficult
Extremely difficult

Scoring

Scores of 5, 10, and 15 are taken as the cut-off points for mild, moderate and severe anxiety, respectively. When used as a screening tool, further evaluation is recommended when the score is 10 or greater.

Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. It is moderately good at screening three other common anxiety disorders - panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%) and post-traumatic stress disorder (sensitivity 66%, specificity 81%).

Source: Spitzer RL, Kroenke K, Williams JBW, Lowe B. A brief measure for assessing generalized anxiety disorder. *Arch Inern Med.* 2006;166:1092-1097.

THE MOOD DISORDER QUESTIONNAIRE

Instructions: Please answer each question to the best of your ability.

1. Has there ever been a period of time when you were not your usual self and	YES	NO
you felt so good or so hyper that other people thought you were not your normal self or you were so hyper that you got into trouble?	0	0
you were so irritable that you shouted at people or started fights or arguments?	0	0
you felt much more self-confident than usual?	0	0
you got much less sleep than usual and found you didn't really miss it?	0	0
you were much more talkative or spoke much faster than usual?	0	0
thoughts raced through your head or you couldn't slow your mind down?	0	0
you were so easily distracted by things around you that you had trouble concentrating or staying on track?	0	0
you had much more energy than usual?	0	0
you were much more active or did many more things than usual?	0	0
you were much more social or outgoing than usual, for example, you telephoned friends in the middle of the night?	0	0
you were much more interested in sex than usual?	0	0
you did things that were unusual for you or that other people might have thought were excessive, foolish, or risky?	0	0
spending money got you or your family into trouble?	0	0
2. If you checked YES to more than one of the above, have several of these ever happened during the same period of time?	0	0
3. How much of a problem did any of these cause you – like being unable to work; having family, money or legal troubles; getting into arguments or fights? Please circle one response only. No Problem Minor Problem Moderate Problem Serious Problem		
4. Have any of your blood relatives (i.e. children, siblings, parents, grandparents, aunts, uncles) had manic-depressive illness or bipolar disorder?	0	0
5. Has a health professional ever told you that you have manic-depressive illness or bipolar disorder?	0	0

Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

Patient Name		Today's I	Date				
scale on the right side of the p best describes how you have for	elow, rating yourself on each of the criteria sloage. As you answer each question, place an a elt and conducted yourself over the past 6 m ur healthcare professional to discuss during to	(in the box that onths. Please give	Never	Rarely	Sometimes	Often	Very Often
How often do you have tro once the challenging parts	ouble wrapping up the final details of a proj have been done?	ect,					
How often do you have dif a task that requires organize	ficulty getting things in order when you havation?	ve to do					
3. How often do you have pro	oblems remembering appointments or obli	gations?					
4. When you have a task that or delay getting started?	requires a lot of thought, how often do yo	ou avoid					
5. How often do you fidget o to sit down for a long time	or squirm with your hands or feet when you	u have					
6. How often do you feel ove were driven by a motor?	erly active and compelled to do things, like	you					
						F	art A
How often do you make of difficult project?	careless mistakes when you have to work o	on a boring or					
8. How often do you have do or repetitive work?	ifficulty keeping your attention when you a	re doing boring					
How often do you have di even when they are speak	ifficulty concentrating on what people say to ting to you directly?	o you,					
10. How often do you misplac	ce or have difficulty finding things at home	or at work?					
II. How often are you distrac	cted by activity or noise around you?						
12. How often do you leave y you are expected to rema	your seat in meetings or other situations in ain seated?	which					
13. How often do you feel re	estless or fidgety?						
14. How often do you have d to yourself?	lifficulty unwinding and relaxing when you h	nave time					
15. How often do you find yo	ourself talking too much when you are in s	ocial situations?					
16. When you're in a convers the sentences of the peop them themselves?	sation, how often do you find yourself finishole you are talking to, before they can finish	ning 1					
17. How often do you have of turn taking is required?	difficulty waiting your turn in situations whe	en					
18. How often do you interru	upt others when they are busy?						
							Part B

Please circle any that you have experienced

Low mood for >2 weeks Grandiose Hallucinations/illusions Trembling Less sleep Increased activity Delusions Palpitations goal-directed/high risk Low interest Self-reference: Nausea/chills Guilt/worthlessness Choking/chest pain Impulsivity people watching you Low or high energy Incr sex drive talking about you Sweating Low concentration Irritability messages from media Fear: Need less sleep Appetite/weight changes Thought blocking/insertion dying/going crazy Elevated mood Anticipatory anxiety Psychomotor slowing Disorganization: Suicide: Speedy talking speech/behavior Avoidance Hopelessness/Plan/Access Speedy thoughts Excess worry Intrusive/persistent thoughts Experienced/witness event Restless/edgy Recognized as excessive/irrational Persistent re-experiencing Easily fatigued Repetitive behaviors: Dreams/flashbacks Washing/cleaning Muscle tension Avoidance behavior Counting/checking ⊥ sleep Hyper-arousal: concentration Organizing/praying ↑ vigilance/↑ startle Fear abandonment/rejection Forensic history: Performance situations: arrests/imprisonment Fear of embarrassment Unstable relationships Fear of humiliation Chronic emptiness Aggressiveness/violence self esteem Criticism Lack of empathy/remorse Intense anger/outbursts Lack of concern for safety: Self-damaging behavior self or others Labile mood and impulsivity Childhood conduct disorder Heights/crowds/animals Excess concern with appearance Binging/purging/restriction/amenorrhea Mood changes unrelated to circumstances Perception of body image or weight or certain part of body Agitation Avoidance behavior Cutting, scratching, burning, purging

Is there someone in your family who has had depression, anxiety, bipolar, schizophrenia? If so, please list illness and relation to you:

Name/date